

Policy Number	DME104.001
Policy Effective Date	08/01/2024

Upper-Limb Prosthesis, Including Myoelectric and Orthotic Components, and Other Prosthetics Except for Lower-Limb Prosthesis

Table of Contents
Coverage
Policy Guidelines
Description
Rationale
Coding
References
Policy History

Related Policies (if applicable)
None

Disclaimer

Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

Legislative Mandates

EXCEPTION: For HCSC members residing in the state of Arkansas, § 23-99-417 relating to orthotic devices, orthotic services, prosthetic devices, and prosthetic services, requires coverage for an orthotic device or service, a prosthetic device or service, prosthetic device for athletics or recreation, or a prosthetic device for showering or bathing. "Prosthetic device for athletics or recreation" means a device that provides an individual with the ability or potential for prosthesis ambulation that exceeds basic ambulation skills, exhibiting high impact, stress, or energy levels including the use of a blade-type foot designed for running and other high activity or high-impact endeavors. A candidate for a recreational prosthesis shall qualify in the Medicare functional level status as a K-3 or K-4 functional level as a user who: 1) Can achieve any high-level activity pursuits; and 2) Exhibits an ability to perform above and beyond normal ambulation. Coverage is not required for a device or service more than once every three [3] years unless medically necessary. This applies to the following: Fully Insured Group, Student, Small Group, Mid-Market, Large Group, HMO, EPO, PPO, POS. Unless indicated by the group, this mandate or coverage will not apply to ASO groups.

EXCEPTION: For HCSC members residing in the state of Arkansas, § 23-99-405 related to coverage of mastectomy and reconstruction services, should an enrollee elect reconstruction after a mastectomy, requires coverage for surgery and reconstruction of the breast on which the mastectomy has been performed, surgery and reconstruction of the other breast to produce a symmetrical appearance, and prostheses and coverage for physical complications at all stages of a mastectomy, including lymphedema. This applies to the following: Fully Insured Group, Student, Small Group, Mid-Market, Large Group, HMO, EPO, PPO, POS. Unless indicated by the group, this mandate or coverage will not apply to ASO groups.

Coverage

Myoelectric Upper-Limb Prosthetic Components

Myoelectric upper-limb prosthetic components **may be considered medically necessary** when **all** of the following conditions are met:

- The individual has an amputation or missing limb at the wrist or above (e.g., forearm, elbow, etc.); **and**
- Standard body-powered prosthetic devices cannot be used or are insufficient to meet the functional needs of the individual in performing activities of daily living; **and**
- The remaining musculature of the arm(s) contains the minimum microvolt threshold to allow operation of a myoelectric prosthetic device; **and**
- The individual has demonstrated sufficient neurological and cognitive function to operate the prosthesis effectively; **and**
- The individual is free of comorbidities that could interfere with function of the prosthesis (e.g., neuromuscular disease, etc.); **and**
- Functional evaluation indicates that with training, use of a myoelectric prosthesis is likely to meet the functional needs of the individual (e.g., gripping, releasing, holding, and coordinating movement of the prosthesis) when performing activities of daily living. This evaluation should consider the individual's needs for control, durability (maintenance), function (speed, work capability), and usability.

Advanced upper-limb prosthetic components with both sensor (input device options such as a pressure sensor, rocker switch or linear transducer) and myoelectric control (e.g., LUKE Arm) **are considered experimental, investigational and/or unproven.**

A prosthesis with individually powered digits, including but not limited to a partial hand prosthesis, **is considered experimental, investigational and/or unproven.**

Myoelectric upper-limb prosthetic components **are considered not medically necessary** under all other conditions.

NOTE 1: Upper-limb amputees should be evaluated by an independent qualified professional to determine the most appropriate prosthetic components and control mechanism (e.g., body-powered, myoelectric, or combination of body-powered and myoelectric). A trial period may be indicated to evaluate the tolerability and efficacy of the prosthesis in a real-life setting.

Myoelectric Upper-Limb Orthoses

Myoelectric controlled upper-limb orthoses **are considered experimental, investigational and/or unproven.**

Prosthetic Appliances and Other Devices

Prosthetic appliances (except myoelectric prosthetics) **may be considered medically necessary** when prescribed by a qualified provider to replace absent or nonfunctioning parts of the body by an artificial substitute. Prosthetic appliances include, but are not limited to:

- **Surgical Prostheses:**
 1. Artificial joints necessary for joint repair and/or reconstructive surgery, except ankle replacement and intervertebral disc replacement (refer to **NOTE 4** below);
 2. Artificial urethral sphincters for urinary incontinence;
 3. Facial, ear, and nose prostheses, when necessary to correct absence or deformity that is congenital or acquired as a result of trauma or disease (refer to **NOTE 4** below).
- **Nonsurgical Prostheses:**
 1. Artificial eyes;
 2. External breast prosthesis, including surgical brassiere, for individuals who have had a mastectomy;
 3. Standard mechanical and manual control prosthetic components replacing all or part of absent upper-limbs (refer to **NOTE 4** below).

NOTE 2: Member contract benefit may vary. **CHECK CONTRACTS FOR COVERAGE ELIGIBILITY** of prosthetics, including, but not limited to the following appliances or devices:

- Dentures replacing teeth or structures directly supporting teeth;
- Electrical incontinence aids or alarms, for either bowel or bladder incontinence;
- Hairpieces or wigs for alopecia;
- Hearing aids (refer to **NOTE 4** below);
- Implants for cosmetic purposes (refer to **NOTE 3** below);
- Penile prostheses (refer to **NOTE 4** below).

NOTE 3: The above list is not all-inclusive. Please refer to the individual contract for exclusions related to prosthetic appliances.

NOTE 4: Many of the topics listed in coverage may be addressed in detail on one or more specific medical policies, which can be identified by a keyword(s) search of Medical Policies on the internet.

Policy Guidelines

Coverage should include supplies necessary for the effective use of a covered prosthetic appliance (i.e., harnesses, stump socks, etc., essential to use of an artificial limb).

Adjustments and repairs to the device(s) are covered when they are necessary to make the equipment functional for as long as the equipment continues to be medically required.

Replacement is provided for the device(s) only if:

- Wear and tear has made them ineffective; and/or
- The device malfunctions; and/or
- The patient's size changes and adjustments to the device can no longer be made (i.e., a child grows and artificial limb no longer fits or functions).

Hearing Aids

LT (left) or RT (right) modifiers must be used on monaural codes to identify which ear the hearing aid is to be used. LT or RT modifiers should not be used on bilateral or binaural codes as the “bi” indicates that it is for two ears.

Description

Among those living with limb loss, the main causes are vascular disease (54%) including diabetes and peripheral arterial disease, trauma (45%), and cancer (less than 2%). There are nearly 2 million people living with limb loss in the U.S. There are more than 1 million annual limb amputations globally, one every 30 seconds. (1)

Background

The need for a prosthesis can occur for a number of reasons, including trauma, surgery, or congenital anomalies. Prosthetics are artificial substitutes that replace all or part of a body organ or replace all or part of the function of a permanently inoperative, absent, or malfunctioning body part. Not all body or organ loss will be addressed in this description, only upper-limb prosthetics will be included below.

Upper-Limb Prosthetics

Myoelectric prostheses are powered by electric motors with an external power source. The joint movement of an upper-limb prosthesis (e.g., hand, wrist, and/or elbow) is driven by microchip-processed electrical activity in the muscles of the remaining limb or limb stump.

Treatment

The primary goals of the upper-limb prostheses are to restore function and natural appearance. Achieving these goals also requires sufficient comfort and ease of use for continued acceptance by the wearer. The difficulty of achieving these diverse goals with an upper-limb prosthesis increases with the level of amputation (digits, hand, wrist, elbow, shoulder), and thus the complexity of joint movement increases.

Upper-limb prostheses are classified into 3 categories depending on the means of generating movement at the joints: passive, body-powered, and electrically powered movement. All 3

types of prostheses have been in use for more than 30 years; each possesses unique advantages and disadvantages.

Passive Prostheses:

The passive prostheses rely on manual repositioning, typically using the opposite arm and cannot restore function. This unit is the lightest of the 3 prosthetic types and is thus generally the most comfortable.

Body-Powered Prostheses:

The body-powered prostheses use a body harness and cable system to provide functional manipulation of the elbow and hand. Voluntary movement of the shoulder and/or limb stump extends the cable and transmits the force to the terminal device. Prosthetic hand attachments, which may be claw-like devices that allow good grip strength and visual control of objects or latex-gloved devices that provide a more natural appearance at the expense of control, can be opened and closed by the cable system. Patient complaints with body-powered prostheses include harness discomfort, particularly the wear temperature, wire failure, and the unattractive appearance.

Myoelectric Prostheses:

- Myoelectric prostheses use muscle activity from the remaining limb for control of joint movement. Electromyographic signals from the limb stump are detected by surface electrodes, amplified, and then processed by a controller to drive battery-powered motors that move the hand, wrist, or elbow. Although upper-arm movement may be slow and limited to 1 joint at a time, myoelectric control of movement may be considered the most physiologically natural.
- Myoelectric hand attachments are similar in form to those offered with the body-powered prosthesis but are battery-powered. Commercially available examples are listed in the Regulatory Status of the Description section.
- A hybrid system, a combination of body-powered and myoelectric components, may be used for high-level amputations (at or above the elbow). Hybrid systems allow for control of 2 joints at once (i.e., 1 body-powered, 1 myoelectric) and are generally lighter and less expensive than a prosthesis composed entirely of myoelectric components.

Technology in this area is rapidly changing, driven by advances in biomedical engineering and by the U.S. Department of Defense (DoD) Advanced Research Projects Agency (DARPA), which is funding a public and private collaborative effort on prosthetic research and development. Areas of development include the use of skin-like silicone elastomer gloves, “artificial muscles,” and sensory feedback. Smaller motors, microcontrollers, implantable myoelectric sensors, and reinnervation of remaining muscle fibers are being developed to allow fine movement control. Lighter batteries and newer materials are being incorporated into myoelectric prostheses to improve comfort.

The LUKE Arm (previously known as the DEKA Arm System) was developed in a joint effort between DEKA Research & Development and the Defense Advanced Research Projects Agency’s

(DARPA) Revolutionizing Prosthetics program. It is the first commercially available myoelectric upper-limb that can perform complex tasks with multiple simultaneous powered movements (e.g., movement of the elbow, wrist, and hand at the same time). In addition to the electromyographic electrodes, the LUKE Arm contains a combination of mechanisms, including switches, movement sensors, and force sensors. The primary control resides with inertial measurement sensors on top of the feet. This control directs the LUKE arm movement when rolling, tapping, etc., the foot. The prosthesis includes vibration pressure and grip sensors.

Myoelectric Orthoses

The MyoPro® (Myomo) is a myoelectric powered upper-extremity orthotic. This orthotic device weighs about 1.8 kilograms (kg; 4 pounds), has manual wrist articulation, and myoelectric initiated bi-directional elbow movement. The MyoPro® detects weak muscle activity from the affected muscle groups. A therapist or prosthetist/orthoptist can adjust the gain (amount of assistance), signal boost, thresholds, and range of motion. Potential users include individuals with traumatic brain injury, spinal cord injury, brachial plexus injury, amyotrophic lateral sclerosis, and multiple sclerosis. Use of robotic devices for therapy has been reported. The MyoPro® is the first myoelectric orthotic available for home use.

Additional Information and Terminology

- *Terminal devices*: these devices are separated into four categories: passive hands, split hooks, mechanical hands, and specialized tools.
- *Passive hand*: no gripping action but does provide functional value to the unilateral amputee as a holding and positioning aid.
- *Split hook*: the most popular terminal device because it provides gripping and precise manipulation at its tip.
- *Mechanical hands*: more cosmetic than hooks but provide little real functional benefit.
- *Special tools*: can be designed for a variety of activities, such as sports, or allowing attachment of hand tools or kitchen utensils.
- *Electronic terminal devices*: come in hook styles but are usually requested as hands; these have two functions: fast clasping and firm gripping.
- *Exoskeletal*: a hard shell built up into the shape of the limb with a hollow center.
- *Endoskeletal*: component parts covered with an external soft foam covering.
- *Typical below-elbow prosthesis*: usually includes the following components: voluntary opening split hook, friction wrist, double-walled plastic laminate socket, flexible elbow hinge, single-control cable system, biceps or triceps cuff, and figure-of-8 harness.
- *Typical above-elbow prosthesis*: is similar but substitutes an internal-locking elbow for the flexible elbow hinge, uses a dual control cable (instead of single control), and does not have a biceps or triceps cuff.

Regulatory Status

Manufacturers must register prostheses with the Restorative and Repair Devices Branch of the U.S. Food and Drug Administration (FDA) and keep a record of any complaints, but do not have to undergo a full FDA review.

Available myoelectric devices include, but are not limited to, ProDigits™ and i-limb™ (Touch Bionics), the SensorHand™ Speed and Michelangelo® Hand (Otto Bock), the LTI Boston Digital Arm™ System (Liberating Technologies), the Utah Arm Systems (Motion Control), and bebionic (Ottobock).

In 2014, the DEKA Arm System (DEKA Integrated Solutions, now DEKA Research & Development), now called the LUKE™ Arm (Mobius Bionics), was cleared for marketing by the FDA through the *de novo* 513(f)(2) classification process for novel low- to moderate-risk medical devices that are first-of-a-kind.

FDA product codes: GXY, IQZ.

The MyoPro® (Myomo) is registered with the FDA as a class 1 limb orthosis.

Rationale

This medical policy was created in 1990 and since then updated periodically using the PubMed database. The most recent literature review was performed through January 29, 2024.

Medical policies assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function—including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Prospective comparative studies with objective and subjective outcome measures would provide the most informative data on which to compare different prostheses, but little evidence was identified that directly addresses whether standard myoelectric prostheses

improve function and health-related quality of life. This may be true for non-myoelectric prosthetics.

The available indirect evidence is based on 2 assumptions: 1) use of any prosthesis confers a clinical benefit and 2) self-selected use is an acceptable measure of the perceived benefit (combination of utility, comfort, appearance) of a particular prosthesis for that person. Most studies identified have described amputees' self-selected use or rejection rates. The results are usually presented as hours worn at work, hours worn at home, and hours worn in social situations. Amputees' self-reported reasons for use and abandonment are also frequently reported. Upper-limb amputee's needs may depend on the particular situation; e.g., the increased functional capability may be needed with heavy work or domestic duties, while a more naturally appearing prosthesis with reduced functional capability may be acceptable for an office, school, or other social environment.

Myoelectric Proximal Upper-Limb Prostheses

Clinical Context and Therapy Purpose

The purpose of myoelectric upper-limb prosthesis components at or proximal to the wrist is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with a missing limb at the wrist or higher.

The following PICO was used to select literature to inform this policy.

Population

Individuals with a missing limb at the wrist or higher.

Intervention

Myoelectric upper-limb prosthesis components at or proximal to the wrist.

Comparator(s)

The body-powered prosthesis.

Outcomes

Relevant outcomes include: Functional outcomes in the use of the Myoelectric upper limb prosthesis and impact on quality of life. Follow-up ranged on average between 2 years and 4 years.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;

- Studies with duplicative or overlapping populations were excluded.

Systematic Reviews

A 2007 systematic review of 40 articles published over the previous 25 years assessed upper-limb prosthesis acceptance and abandonment (see Table 1). (2) For pediatric patients, the mean rejection rate was 38% for passive prostheses (1 study), 45% for body-powered prostheses (3 studies), and 32% for myoelectric prostheses (12 studies) (see Table 2). For adults, there was considerable variation between studies, with mean rejection rates of 39% for passive (6 studies), 26% for body-powered (8 studies), and 23% for myoelectric (10 studies) prostheses. Reviewers found no evidence that the acceptability of passive prostheses had declined over the period from 1983 to 2004, “despite the advent of myoelectric devices with functional as well as cosmetic appeal.” Body-powered prostheses were also found to have remained a popular choice, with the type of hand attachment being the major factor in acceptance. Body-powered hooks were considered acceptable by many users, but body-powered hands were frequently rejected (80%-87% rejection rates) due to slowness in movement, awkward use, maintenance issues, excessive weight, insufficient grip strength, and the energy needed to operate. Rejection rates of myoelectric prostheses tended to increase with longer follow-up. There was no evidence of a change in rejection rates over the 25 years of study, but the results were limited by sampling bias from isolated populations and the generally poor quality of studies selected.

Within-Subject Comparisons

One prospective controlled study (1993) compared preferences for body-powered with myoelectric hands in children. (3) Juvenile amputees (toddlers to teenagers) were fitted in a randomized order with one of the 2 types of prostheses; after a 3-month period, the terminal devices were switched, and the children selected one of the prostheses to use. At the time of follow-up, more than a third of children were wearing the myoelectric prosthesis, a third were wearing a body-powered prosthesis, and 22% were not using a prosthesis (see Table 2). There was no difference in the children’s ratings of the myoelectric and body-powered devices.

Silcox et al. (1993) conducted a within-subject comparison of preference for body-powered or myoelectric prostheses in adults. (4) Of 44 patients fitted with a myoelectric prosthesis, 91% also owned a body-powered prosthesis, and 20% owned a passive prosthesis. Rejection rates of these prostheses are shown in Table 2. Use of a body-powered prosthesis was unaffected by the type of work; good-to-excellent use was reported in 35% of patients with heavy work demands and 39% of patients with light work demands. In contrast, the proportion of patients using a myoelectric prosthesis was higher in the group with light work demands (44%) than in those with heavy work demands (26%). There was also a trend toward the higher use of the myoelectric prosthesis compared with a body-powered prosthesis in social situations. Appearance was cited more frequently as a reason for using a myoelectric prosthesis than any other factor. Weight and speed were more frequently cited than any other factors as reasons for nonuse of the myoelectric prosthesis.

McFarland et al. (2010) conducted a cross-sectional survey of major combat-related upper-limb loss in veterans and service members from Vietnam (n=47) and Iraq (n=50) recruited through a national survey. (5) In the first year of limb loss, the Vietnam group received a mean of 1.2 devices (usually body-powered), while the Iraq group received a mean of 3.0 devices (typically 1 myoelectric/hybrid, 1 body-powered, 1 cosmetic). Preferences in the Iraq group are shown in Table 2. At the time of the survey, upper-limb prosthetic devices were used by 70% of the Vietnam group and 76% of the Iraq group. The most common reasons for rejection included short residual limbs, pain, poor comfort (e.g., the weight of the device), and lack of functionality.

Table 1. Summary of Key Study Characteristics

Author	Study Type	N	Dates	Participants	Intervention	FU
Rejection Rates						
Biddiss et al. (2007) (2)	Systematic review	40 articles	1983-2004	Pediatric and adult		25 years
Silcox et al. (1993) (4)	Within-subject comparison	44		Adult	All fitted with a myoelectric prosthesis	
Sjoberg et al. (2017) (6)	Prospective case-control	<ul style="list-style-type: none"> • 9 children <2.5 years • 27 children >2.5 to 4 years 	1994-2002	Pediatric	Training with a myoelectric prosthesis	Until 12 years of age
Acceptance Rates						
Kruger and Fishman (1993) (3)	Randomized within-subject comparison	78		Pediatric	Trial period for both myoelectric and body-powered	2 years
McFarland et al. (2010) (5)	Cross-sectional survey	50		Veterans and service members	Provided with all 3 device types	
Egermann et al. (2009) (7)	Parental questionnaire	41		Pediatric (2-5 years)	Training with a myoelectric prosthesis	2 years (range, 0.7-5 years)

N: number; FU: follow-up.

Table 2. Summary of Key Study Outcomes

Author	Outcomes	Adult and Pediatric	Myoelectric	Body-Powered	Passive	None
Rejection Rates						
Biddiss et al. (2007) (2)	Mean rejection rates	Pediatric	32%	45%	38%	
		Adult	23%	26%	39%	
Silcox et al. (1993) (4)	Rejection of own prosthesis	Adult	22 (50%)	13 (32%)	5 (55%)	
Sjoberg et al. (2017) (6)	Rejection of a myoelectric prosthesis	<2.5 years	3 (33%)			
		2.5 to 4 years	4 (15%)			
Acceptance and Preference Rates						
Kruger and Fishman (1993) (3)	Preference rates		34 (44%)	26 (34%)		18 (22%)
McFarland et al. (2010) (5)	Preference rates	Iraq Veterans	18 (36%)	15 (30%)		11 (22%)
Egermann et al. (2009) (7)	Acceptance	Pediatric	31 (76%)			

Values are percent or number (%).

Acceptance Rates in Children

Sjoberg et al. (2018) conducted a prospective long-term case-control study to determine whether fitting a myoelectric prosthesis before 2.5 years of age improved prosthesis acceptance rates compared with the current Scandinavian standard of fitting between 2.5 and 4 years old. (6) All children had a congenital amputation and had used a passive hand prosthesis from 6 months of age, and both groups (case, n=9; control, n=27) were fitted with the same type of prosthetic hand and received structured training beginning at 3 years of age. They were followed every 6 months between 3 and 6 years of age and then as needed for service or training for a total of 17 years. Prosthetic skill measured by the Skills Index Ranking Scale (SIRS) increased over time, however, there were no statistically significant differences between groups. By 12 years of age, all but one child in the case group and all but 2 children in the control group achieved maximum performance on the Skills Index Ranking Scale (SIRS) (level 14, the ability to throw objects from above the shoulder). To note, 3 (33%) children in the case group and 4 (15%) in the control group were lost to follow-up at after 9 years of age due to prosthetic rejection. This difference was not statistically significant in this small study. Overall, study results did not favor earlier intervention with a myoelectric prosthesis.

Egermann et al. (2009) evaluated the acceptance rate of a myoelectric prosthesis in 41 children between 2 and 5 years of age. (7) To be fitted with a myoelectric prosthesis, the children had to communicate well and follow instructions from strangers, have interest in an artificial limb, have bimanual handling (use of both limbs in handling objects), and have a supportive family setting. A 1- to 2-week interdisciplinary training program (inpatient or outpatient) was provided for the child and parents. At a mean 2-year follow-up (range, 0.7-5.1 years), a questionnaire was distributed to evaluate acceptance and use during daily life (100% return rate). Successful use, defined as a mean daily wearing time of more than 2 hours, was achieved in 76% of the study group. The average daily use was 5.8 hours per day (h/d; range, 0-14 h/d). The level of amputation significantly influenced the daily wearing time, with above elbow amputees wearing the prosthesis for longer periods than children with below-elbow amputations. Three (60%) of 5 children with amputations at or below the wrist refused use of any prosthetic device. There were statistically nonsignificant trends for increased use in younger children, in those who had inpatient occupational training, and in children who had a previous passive (versus body-powered) prosthesis. During the follow-up period, maintenance averaged 1.9 times per year (range, 0-8 repairs); this was correlated with the daily wearing time. The authors noted that more important selection criteria than age were the activity and temperament of the child (e.g., a myoelectric prosthesis would more likely be used in a calm child interested in quiet bimanual play, whereas a body-powered prosthesis would be more durable for outdoor sports, and in sand or water).

Section Summary: Myoelectric Upper-Limb Prosthesis

The identified literature focuses primarily on patient acceptance and rejection; data are limited or lacking in the areas of function and functional status. The limited evidence suggests that the percentage of amputees who accept a myoelectric prosthesis is approximately the same as those who prefer to use a body-powered prosthesis, and that self-selected use depends partly on the individual's activities of daily living. When compared with body-powered prostheses, myoelectric components possess similar capability to perform light work, and myoelectric components may improve range of motion. The literature has also indicated that appearance is most frequently cited as an advantage of myoelectric prostheses, and for individuals who desire a restorative appearance, the myoelectric prosthesis can provide greater function than a passive prosthesis, with equivalent function to a body-powered prosthesis for light work.

Sensor and Myoelectric Upper-Limb Components

Clinical Context and Therapy Purpose

The purpose of implantation of sensor and myoelectric controlled upper-limb prosthetic components is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with a missing limb at the wrist or higher who receive sensor and myoelectric controlled upper-limb prosthetic components.

The following PICO was used to select literature to inform this policy.

Population

Individuals with a missing limb at the wrist or higher who receive sensor and myoelectric controlled upper-limb prosthetic components.

Intervention

Implantation of sensor and myoelectric controlled upper-limb prosthetic components.

Comparator(s)

Use of a conventional prosthesis.

Outcomes

Relevant outcomes include: Functional outcomes in the use of the myoelectric upper limb prosthesis and impact on quality of life. Outcomes were both performance-based and self-reported measures.

Follow-up ranged on average between 2 years and 4 years.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Investigators from 3 Veterans Administration (VA) medical centers and the Center for the Intrepid at Brooke Army Medical Center published a series of reports on home use of the LUKE prototype (DEKA Gen 2 and DEKA Gen 3) in 2017 and 2018. (8-13) Participants were included in the in-laboratory training if they met criteria and had sufficient control options (e.g., myoelectric and/or active control over one or both feet) to operate the device. In-lab training included a virtual reality training component. At the completion of the in-lab training, the investigators determined, using a priori criteria, which participants were eligible to continue to the 12-week home trial. The criteria included the independent use of the prosthesis in the laboratory and community setting, fair, functional performance, and sound judgment when operating or troubleshooting minor technical issues. On ClinicalTrials.gov, the total enrollment target is listed as 100 patients with study completion by February 2018 (NCT01551420).

Resnick et al. (2017) reported on the acceptance of the LUKE prototype before and after a 12-week trial of home use. (8) Of 42 participants enrolled at the time, 32 (76%) participants completed the in-laboratory training, 22 (52%) wanted to receive a LUKE Arm and proceeded to the home trial, 18 (43%) completed the home trial, and 14 (33%) expressed a desire to receive the prototype at the end of the home trial. Over 80% of those who completed the home trial

preferred the prototype arm for hand and wrist function, but as many preferred the weight and look of their own prosthesis. One-third of those who completed the home training thought that the arm was not ready for commercialization. Participants who completed the trial were more likely to be prosthesis users at study onset ($p=0.03$), and less likely to have musculoskeletal problems ($p=0.047$). (9) Reasons for attrition during the in-laboratory training were reported in a separate publication by Resnik and Klinger (2017). (10) Attrition was related to the prosthesis entirely or in part by 67% of the participants, leading to a recommendation to provide patients with an opportunity to train with the prosthesis before a final decision about the appropriateness of the device.

Functional outcomes of the Gen 2 and Gen 3 arms, as compared with participants' prostheses, were reported by Resnick et al. (2018). (11) At the time of the report, 23 regular prosthesis users had completed the in-lab training, and 15 had gone on to complete the home use portion of the study. Outcomes were both performance-based and self-reported measures. At the end of the lab training, dexterity was similar, but performance was slower with the LUKE prototype than with their conventional prosthesis. At the end of the home study, activity speed was similar to the conventional prostheses, and one of the performance measures (Activities Measure for Upper-Limb Amputees) was improved. Participants also reported that they were able to perform more activities, had less perceived disability, and less difficulty in activities, but there were no differences between the 2 prostheses on many of the outcome measures including dexterity, prosthetic skill, spontaneity, pain, community integration, or quality of life. Post hoc power analysis suggested that evaluation of some outcomes might not have been sufficiently powered to detect a difference.

In a separate publication, Resnick et al. (2017) reported that participants continued to use their prosthesis (average, 2.7 h/d) in addition to the LUKE prototype, concluding that availability of both prostheses would have the greatest utility. (12) This conclusion is similar to those from earlier prosthesis surveys, which found that the selection of a specific prosthesis type (myoelectric, powered, or passive) could differ depending on the specific activity during the day. In the DEKA Gen 2 and Gen 3 study reported here, 29% of participants had a body-powered device, and 71% had a conventional myoelectric prosthesis.

Section Summary: Sensor and Myoelectric Upper-Limb Components

The LUKE Arm was cleared for marketing in 2014 and is now commercially available. The prototypes for the LUKE Arm, the DEKA Gen 2 and Gen 3, were evaluated by the U.S. military and VA in a 12-week home study, with study results reported in a series of publications. Acceptance of the advanced prosthesis in this trial was mixed, with one-third of enrolled participants desiring to receive the prototype at the end of the trial. Demonstration of improvement in function has also been mixed. After several months of home use, activity speed was shown to be similar to the conventional prosthesis. There was an improvement in the performance of some, but not all activities. Participants continued to use their prosthesis for part of the day, and some commented that the prosthesis was not ready for commercialization. There were no differences between the LUKE Arm prototype and the participants' prostheses for many outcome measures. Additional study of the current generation of the LUKE Arm is

needed to determine whether the newer models of this advanced prosthesis lead to consistent improvements in function and quality of life.

Myoelectric Hand with Individual Digit Control

Clinical Context and Therapy Purpose

The purpose of a myoelectric upper-limb prosthesis with individually powered digits is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals with a missing hand distal to the wrist.

The following PICO was used to select literature to inform this policy.

Population

Individuals with a missing hand distal to the wrist.

Intervention

A myoelectric upper-limb prosthesis with individually powered digits.

Comparator(s)

Body-powered prosthesis.

Outcomes

Generally, the outcomes were functional status and quality of life.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Although the availability of a myoelectric hand with individual control of digits has been widely reported in lay technology reports, video clips, and basic science reports, no peer-reviewed publications were found to evaluate functional outcomes of individual digit control in amputees.

Myoelectric Orthotic

Clinical Context and Therapy Purpose

The purpose of a myoelectric powered upper-limb orthotic device is to provide a treatment option that is an alternative to or an improvement on existing therapies for individuals who are stable post-stroke, who have upper-limb weakness or paresis.

The following PICO was used to select literature to inform this policy.

Population

Individuals who are stable post-stroke, who have upper-limb weakness or paresis.

Intervention

A myoelectric powered upper-limb orthotic device.

Comparator(s)

Usual care post-stroke.

Outcomes

The functional status and movement of the upper-limb with and without the orthotic in stable post-stroke participants who had no prior experience with the device. Impact on quality of life was also measured.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies;
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought;
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Peters et al. (2017) evaluated the immediate effect (no training) of a myoelectric elbow-wrist-hand orthosis on paretic upper-extremity impairment. (14) Participants (n=18) were stable and moderately impaired with a single stroke, 12 months or later before study enrollment. They were tested using a battery of measures without, and then with the device; the order of testing was not counterbalanced. The primary measure was the upper-extremity section of the Fugl-Meyer Assessment, a validated scale that determines active movement. Upper-extremity movement on the Fugl-Meyer Assessment was significantly improved while wearing the orthotic (a clinically significant increase of 8.71 points, $p < 0.001$). The most commonly observed gains were in elbow extension, finger extension, grasping a tennis ball, and grasping a pencil. The Box and Block test (moving blocks from one side of a box to another) also improved ($p < 0.001$). Clinically significant improvements were observed for raising a spoon and cup, and there were significant decreases in the time taken to grasp a cup and gross manual dexterity. Performance on these tests changed from unable to able to complete. The functional outcome measures (raising a spoon and cup, turning on a light switch, and picking up a laundry basket with 2 hands) were developed by the investigators to assess these moderately impaired participants. The authors noted that performance on these tasks was inconsistent and

proposed a future study that would include training with the myoelectric orthosis before testing.

Section Summary: Myoelectric Orthotic

The largest study identified tested participants with and without the orthosis. This study evaluated the function with and without the orthotic in stable poststroke participants who had no prior experience with the device. Outcomes were inconsistent. Studies are needed that show consistent improvements in relevant outcome measures. Results should also be replicated in a larger number of patients.

Prosthetic Appliances and Other Devices

Determinations of whether a proposed prosthetic would be considered a replacement of an absent or nonfunctioning part of the body should always be interpreted in the context of the specific benefit language.

Section Summary: Prosthetic Appliances and Other Devices

The requirement for the replacement of an absent or nonfunctioning body part for a specific patient condition may vary depending on the individual patient. It should be noted that the presence of a functional impairment or body part absence would render a prosthetic therapy medically necessary when meeting specific criteria listed in coverage.

Summary of Evidence

For individuals who have a missing limb at the wrist or higher who receive myoelectric upper-limb prosthesis components at or proximal to the wrist, the evidence includes a systematic review and comparative studies. Relevant outcomes are functional outcomes and quality of life. The goals of upper-limb prostheses relate to restoration of both appearance and function while maintaining sufficient comfort for continued use. The identified literature focuses primarily on patient acceptance and rejection; data are limited or lacking in the areas of function and functional status. The limited evidence suggests that, when compared with body-powered prostheses, myoelectric components possess the similar capability to perform light work; however, myoelectric components could also suffer a reduction in performance when operating under heavy working conditions. The literature has also indicated that the percentage of amputees who accept the use of a myoelectric prosthesis is approximately the same as those who prefer to use a body-powered prosthesis, and that self-selected use depends partly on the individual's activities of daily living. Appearance is most frequently cited as an advantage of myoelectric prostheses, and for patients who desire a restorative appearance, the myoelectric prosthesis can provide greater function than a passive prosthesis--with equivalent function to a body-powered prosthesis for light work. Because of the different advantages and disadvantages of currently available prostheses, myoelectric components for persons with an amputation at the wrist or above may be considered when passive or body-powered prostheses cannot be used or are insufficient to meet the functional needs of the patient in activities of daily living. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have a missing limb at the wrist or higher who receive sensor and myoelectric controlled upper-limb prosthetic components, the evidence includes a series of publications from a 12-week home study. Relevant outcomes are functional outcomes and quality of life. The prototypes for the advanced prosthesis were evaluated by the U.S. Military and Veterans Administration (VA). Demonstration of improvement in function has been mixed. After several months of home use, activity speed was shown to be similar to the conventional prosthesis, and there were improvements in the performance of some activities, but not all. There were no differences between the prototype and the participants' prostheses for outcomes of dexterity, prosthetic skill, spontaneity, pain, community integration, or quality of life. Study of the current generation of the sensor (input device options such as a pressure sensor, rocker switch or linear transducer) and myoelectric controlled prosthesis is needed to determine whether newer models of this advanced prosthesis lead to consistent improvements in function and quality of life. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have a missing limb distal to the wrist who receive a myoelectric prosthesis with individually powered digits, no peer-reviewed publications evaluating functional outcomes in amputees were identified. Relevant outcomes are functional outcomes and quality of life. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with upper-extremity weakness or paresis who receive a myoelectric powered upper-limb orthosis, the evidence includes a small within-subject study. Relevant outcomes are functional outcomes and quality of life. The largest study (N=18) identified tested participants with and without the orthosis but did not provide any training with the device. Performance on the tests was inconsistent. Studies are needed that show consistent improvements in relevant outcome measures. Results should also be replicated in a larger number of patients. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals requiring a replacement of an absent or nonfunctioning part of the body, determinations should be viewed in the context of the specific benefit language for the individual. Relevant outcomes are functional outcomes and quality of life. The evidence is sufficient to determine the effects of the technology on health outcomes.

Professional Guidelines and Position Statements

There are no professional guidelines and position statements that would likely influence this policy.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this policy are listed in Table 3.

Table 3. Summary of Key Trials

NCT Number	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT03401762	Wearable MCI [myoelectric computer interface] to Reduce Muscle Co-activation in Acute and Chronic Stroke	96	Aug 2024
NCT05768802	Evaluation of Myoelectric Implantable Recording Array (MIRA) in Participants With Transradial Amputation (MIRA)	5	Dec 2029
NCT03178890 ^a	The Osseointegrated Human-machine Gateway	18	May 2024
Unpublished			
NCT02274532	Myoelectric SoftHand Pro to Improve Prosthetic Function for People with Below-elbow Amputations: A Feasibility Study	18	May 2016 (completed)

NCT: National Clinical Trial.

^a: Denotes industry-sponsored or cosponsored trial.

Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member’s benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

CPT Codes	97763
HCPCS Codes	A4280, A9282, L6000, L6010, L6020, L6026, L6050, L6055, L6100, L6110, L6120, L6130, L6200, L6205, L6250, L6300, L6310, L6320, L6350, L6360, L6370, L6380, L6382, L6384, L6386, L6388, L6400, L6450, L6500, L6550, L6570, L6580, L6582, L6584, L6586, L6588, L6590, L6600, L6605, L6610, L6611, L6615, L6616, L6620, L6621, L6623, L6624, L6625, L6628, L6629, L6630, L6632, L6635, L6637, L6638, L6640, L6641, L6642, L6645, L6646, L6647, L6648, L6650, L6655, L6660, L6665, L6670, L6672, L6675, L6676, L6677, L6680, L6682, L6684, L6686, L6687, L6688, L6689, L6690, L6691, L6692, L6693, L6694, L6695, L6696, L6697, L6698, L6703, L6704, L6706, L6707, L6708, L6709, L6711, L6712, L6713, L6714, L6715, L6721, L6722, L6805, L6810, L6880, L6881, L6882, L6883, L6884, L6885, L6890, L6895, L6900, L6905, L6910, L6915, L6920, L6925, L6930, L6935, L6940, L6945, L6950, L6955, L6960, L6965, L6970, L6975, L7007, L7008, L7009, L7040, L7045, L7170, L7180, L7181, L7185, L7186, L7190, L7191, L7259, L7360,

	L7362, L7364, L7366, L7367, L7368, L7400, L7401, L7402, L7403, L7404, L7405, L7499, L7510, L7520, L7600, L7700, L8000, L8001, L8002, L8010, L8015, L8020, L8030, L8031, L8032, L8035, L8039, L8040, L8041, L8042, L8043, L8044, L8045, L8046, L8047, L8048, L8049, L8500, L8501, L8505, L8507, L8509, L8510, L8511, L8512, L8513, L8514, L8515, L8600, L8613, L8630, L8631, L8641, L8642, L8658, L8659, L8692, L8699, L8701, L8702, L9900, S8270, V2623, V2624, V2625, V2626, V2628, V2629, V5011, V5014, V5020, V5030, V5040, V5050, V5060, V5070, V5080, V5090, V5100, V5110, V5120, V5130, V5140, V5150, V5160, V5170, V5180, V5190, V5200, V5210, V5220, V5230, V5240, V5241, V5242, V5243, V5244, V5245, V5246, V5247, V5248, V5249, V5250, V5251, V5252, V5253, V5254, V5255, V5256, V5257, V5258, V5259, V5260, V5261, V5262, V5263, V5264, V5265, V5266, V5267, V5268, V5269, V5270, V5271, V5272, V5273, V5274, V5275, V5281, V5282, V5283, V5284, V5285, V5286, V5287, V5288, V5289, V5290, V5298, V5299
--	--

*Current Procedural Terminology (CPT®) ©2023 American Medical Association: Chicago, IL.

References

1. AAS – Amputee Statistics You Ought to Know (2022) Media Release. Advanced Amputee Solutions, LLC. Available at: <<https://www.advancedamputees.com>> (accessed May 8, 2024).
2. Biddiss EA, Chau TT. Upper-limb prosthesis use and abandonment: a survey of the last 25 years. *Prosthet Orthot Int.* Sep 2007; 31(3):236-257. PMID 17979010
3. Kruger LM, Fishman S. Myoelectric and body-powered prostheses. *J Pediatr Orthop.* Jan-Feb 1993; 13(1):68-75. PMID 8416358
4. Silcox DH 3rd, Rooks MD, Vogel RR, et al. Myoelectric prostheses. A long-term follow-up and a study of the use of alternate prostheses. *J Bone Joint Surg Am.* Dec 1993; 75(12):1781-1789. PMID 8258548
5. McFarland LV, Hubbard Winkler SL, Heinemann AW, et al. Unilateral upper-limb loss: satisfaction and prosthetic-device use in veterans and service-members from Vietnam and OIF/OEF conflicts. *J Rehabil Res Dev.* 2010; 47(4):299-316. PMID 20803400
6. Sjoberg L, Lindner H, Hermansson L. Long-term results of early myoelectric prosthesis fittings: A prospective case-control study. *Prosthet Orthot Int.* Oct 2018; 42(5):527-533. PMID 28905686
7. Egermann M, Kasten P, Thomsen M. Myoelectric hand prostheses in very young children. *Int Orthop.* Aug 2009; 33(4):1101-1105. PMID 18636257
8. Resnik LJ, Borgia ML, Acluche F. Perceptions of satisfaction, usability and desirability of the DEKA Arm before and after a trial of home use. *PLoS One.* Jun 2017; 12(6):e0178640. PMID 28575025
9. Resnik L, Cancio J, Klinger S, et al. Predictors of retention and attrition in a study of an advanced upper limb prosthesis: implications for adoption of the DEKA Arm. *Disabil Rehabil Assist Technol.* Feb 2018; 13(2):206-210. PMID 28375687

10. Resnik L, Klinger S. Attrition and retention in upper limb prosthetics research: experience of the VA home study of the DEKA arm. *Disabil Rehabil Assist Technol*. Nov 2017; 12(8):816-821. PMID 28098513
11. Resnik LJ, Borgia ML, Acluche F, et al. How do the outcomes of the DEKA Arm compare to conventional prostheses? *PLoS One*. Jan 2018; 13(1):e0191326. PMID 29342217
12. Resnik L, Acluche F, Lieberman Klinger S, et al. Does the DEKA Arm substitute for or supplement conventional prostheses. *Prosthet Orthot Int*. Oct 2018; 42(5):534-543. PMID 28905665
13. Resnik L, Acluche F, Borgia M. The DEKA hand: A multifunction prosthetic terminal device-patterns of grip usage at home. *Prosthet Orthot Int*. Aug 2018; 42(4):446-454. PMID 28914583
14. Peters HT, Page SJ, Persch A. Giving them a hand: wearing a myoelectric elbow-wrist-hand orthosis reduces upper extremity impairment in chronic stroke. *Ann Rehabil Med*. Sep 2017; 98(9):1821-1827. PMID 28130084

Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been changed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

Policy History/Revision

Date	Description of Change
08/01/2024	Document updated with literature review. Coverage unchanged. No new references added.
08/15/2023	Reviewed. No changes.
06/15/2022	Document updated with literature review. Coverage unchanged. No new references added.
06/15/2021	Document updated with literature review. Coverage unchanged. No new references added.
10/15/2020	Reviewed. No changes.
11/01/2019	Document updated with literature review. The following coverage statements were added: 1) Upper-limb prosthetic components that utilize both sensor (input device options such as a pressure sensor, rocker switch or linear transducer) and myoelectric control are considered experimental, investigational and/or unproven; and 2) Myoelectric controlled upper-limb orthoses are considered experimental, investigational and/or unproven.

	NOTE 1 added; others renumbered. References 5 and 7-13 were added, several removed. Title changed from Prosthetics, Except Lower Limb Prosthetics.
04/15/2017	Reviewed. No changes.
12/01/2016	Document updated with literature review. Coverage unchanged.
03/15/2015	Reviewed. No changes.
02/15/2014	Document updated with literature review. The following was added to the Coverage section: A prosthesis with individually powered digits, including but not limited to a partial hand prosthesis, is considered experimental, investigational and unproven.
12/01/2011	Document updated with literature review. The following medical necessity criteria for myoelectric upper-limb prosthetic components have been added: 1) The patient has an amputation or missing limb at the wrist or above (forearm, elbow, etc.); and 2) The remaining musculature of the arm(s) contains the minimum microvolt threshold to allow operation of a myoelectric prosthetic device; and 3) The patient is free of comorbidities that could interfere with function of the prosthesis (neuromuscular disease, etc.). The following was also added: Myoelectric upper-limb prosthetic components are considered not medically necessary under all other conditions. CPT/HCPCS code(s) updated.
04/01/2009	Revised/updated entire document
01/01/2009	Policy reviewed without literature review; new review date only. This policy is no longer scheduled for routine literature review and update.
06/01/2006	Revised/updated entire document
07/01/2005	CPT/HCPCS code(s) updated
04/01/2005	CPT/HCPCS code(s) updated
01/01/2005	CPT/HCPCS code(s) updated
10/01/2003	CPT/HCPCS code(s) updated
02/01/2002	CPT/HCPCS code(s) updated
06/01/2001	CPT/HCPCS code(s) updated
03/01/2000	Revised/updated entire document
09/01/1999	Revised/updated entire document
04/01/1999	CPT/HCPCS code(s) updated
10/01/1998	Revised/updated entire document
05/01/1996	Revised/updated entire document
05/01/1990	New medical document