

Policy Number	OTH903.029
Policy Effective Date	07/15/2025

Endothelial Keratoplasty

Table of Contents	Related Policies (if applicable)
Coverage	None
Policy Guidelines	
Description	
Rationale	
Coding	
References	
Policy History	

Disclaimer

Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

Coverage

Endothelial keratoplasty (Descemet stripping endothelial keratoplasty, Descemet stripping automated endothelial keratoplasty, Descemet membrane endothelial keratoplasty, or Descemet membrane automated endothelial keratoplasty) **may be considered medically necessary** for the treatment of endothelial dysfunction, including but not limited to:

- Ruptures in Descemet membrane,
- Endothelial dystrophy,
- Aphakic and pseudophakic bullous keratopathy,
- Iridocorneal endothelial syndrome,
- Corneal edema attributed to endothelial failure, and
- Failure or rejection of a previous corneal transplant.

Endothelial keratoplasty **is considered not medically necessary** when endothelial dysfunction is not the primary cause of decreased corneal clarity.

Femtosecond laser-assisted endothelial keratoplasty or femtosecond and excimer laser-assisted endothelial keratoplasty **are considered experimental, investigational, and/or unproven.**

Policy Guidelines

Endothelial keratoplasty should be performed by surgeons adequately trained and experienced in the specific techniques and devices used.

Description

Endothelial keratoplasty, also referred to as posterior lamellar keratoplasty, is a form of corneal transplantation in which the diseased inner layer of the cornea, the endothelium, is replaced with healthy donor tissue. Specific techniques include Descemet stripping endothelial keratoplasty, Descemet stripping automated endothelial keratoplasty, Descemet membrane endothelial keratoplasty, and Descemet membrane automated endothelial keratoplasty. Endothelial keratoplasty, and particularly the specific techniques mentioned, are becoming standard procedures. Femtosecond laser-assisted endothelial keratoplasty and femtosecond and excimer laser-assisted endothelial keratoplasty have also been reported as alternatives to prepare the donor endothelium.

Background

Corneal Disease

The cornea, a clear, dome-shaped membrane that covers the front of the eye, is a key refractive element for vision. Layers of the cornea consist of the epithelium (outermost layer); Bowman layer; the stroma, which comprises approximately 90% of the cornea; Descemet membrane; and the endothelium. The endothelium removes fluid from and limits fluid into the stroma, thereby maintaining the ordered arrangement of collagen and preserving the cornea's transparency. Diseases that affect the endothelial layer include Fuchs endothelial dystrophy, aphakic and pseudophakic bullous keratopathy (corneal edema following cataract extraction), and failure or rejection of a previous corneal transplant.

Treatment

The established surgical treatment for corneal disease is penetrating keratoplasty, which involves the creation of a large central opening through the cornea and then filling the opening with a full-thickness donor cornea that is sutured in place. Visual recovery after penetrating keratoplasty may take 1 year or more due to slow wound healing of the avascular full-thickness incision, and the procedure frequently results in irregular astigmatism due to sutures and the full-thickness vertical corneal wound. Penetrating keratoplasty is associated with an increased risk of wound dehiscence, endophthalmitis, and total visual loss after relatively minor trauma for years after the index procedure. There is also the risk of severe, sight-threatening complications such as explosive suprachoroidal hemorrhage, in which the ocular contents are expelled during the operative procedure, as well as postoperative catastrophic wound failure.

A number of related techniques have been, or are being, developed to selectively replace the diseased endothelial layer. One of the first endothelial keratoplasty techniques was termed *deep lamellar endothelial keratoplasty*, which used a smaller incision than penetrating keratoplasty, allowed more rapid visual rehabilitation, and reduced postoperative irregular astigmatism and suture complications. Modified endothelial keratoplasty techniques include endothelial lamellar keratoplasty, endokeratoplasty, posterior corneal grafting, and microkeratome-assisted posterior keratoplasty. Most frequently used at this time are Descemet stripping endothelial keratoplasty, which uses hand-dissected donor tissue, and Descemet stripping automated endothelial keratoplasty, which uses an automated microkeratome to assist in donor tissue dissection. These techniques include donor stroma along with the endothelium and Descemet membrane, which results in a thickened stromal layer after transplantation. If the donor tissue comprises the Descemet membrane and endothelium alone, the technique is known as Descemet membrane endothelial keratoplasty. By eliminating the stroma on the donor tissue and possibly reducing stromal interface haze, Descemet membrane endothelial keratoplasty is considered a potential improvement over Descemet stripping endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty. A variation of Descemet membrane endothelial keratoplasty is Descemet membrane automated endothelial keratoplasty. Descemet membrane automated endothelial keratoplasty contains a stromal rim of tissue at the periphery of the Descemet membrane endothelial keratoplasty graft to improve adherence and improve handling of the donor tissue. A laser may also be used for stripping in a procedure called femtosecond laser-assisted endothelial keratoplasty and femtosecond and excimer laser-assisted endothelial keratoplasty.

Endothelial keratoplasty involves removal of the diseased host endothelium and Descemet membrane with special instruments through a small peripheral incision. A donor tissue button is prepared from the corneoscleral tissue after removing the anterior donor corneal stroma by hand (e.g., Descemet stripping endothelial keratoplasty) or with the assistance of an automated microkeratome (e.g., Descemet stripping automated endothelial keratoplasty) or laser (femtosecond laser-assisted endothelial keratoplasty or femtosecond and excimer laser-assisted endothelial keratoplasty). Donor tissue preparation may be performed by the surgeon in the operating room or by the eye bank and then transported to the operating room for the final punch out of the donor tissue button. For minimal endothelial damage, the donor tissue must be carefully positioned in the anterior chamber. An air bubble is frequently used to center the donor tissue and facilitate adhesion between the stromal side of the donor lenticule and the host posterior corneal stroma. Repositioning of the donor tissue with the application of another air bubble may be required in the first week if the donor tissue dislocates. The small corneal incision is closed with 1 or more sutures, and steroids or immune-suppressants may be provided topically or orally to reduce the potential for graft rejection. Visual recovery following endothelial keratoplasty is typically 4 to 8 weeks.

Eye Bank Association of America statistics have shown the number of endothelial keratoplasty cases in the United States increased from 30,710 in 2015 to 35,555 in 2019. (1) The Eye Bank Association of America estimated that, as of 2016, nearly 40% of corneal transplants performed

in the United States were endothelial grafts. As with any new surgical technique, questions have been posed about long-term efficacy and risk of complications. Endothelial keratoplasty-specific complications include graft dislocations, endothelial cell loss, and rate of failed grafts. Long-term complications include increased intraocular pressure, graft rejection, and late endothelial failure.

Regulatory Status

Endothelial keratoplasty is a surgical procedure and, as such, is not subject to regulation by the U.S. Food and Drug Administration (FDA). Several microkeratomes have been cleared for marketing by the FDA through the 510(k) process.

Rationale

Medical policies assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function—including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and the quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Comparative Studies

Woo et al. (2019) published the results of a retrospective comparative cohort study comparing long-term graft survival outcomes and complications of patients enrolled in the Singapore Corneal Transplant Registry. (2) Patients with Fuchs endothelial corneal dystrophy and bullous keratopathy underwent Descemet membrane endothelial keratoplasty (121 eyes), Descemet stripping automated endothelial keratoplasty (423 eyes), or penetrating keratoplasty (405 eyes). Descemet membrane endothelial keratoplasty demonstrated better graft survival compared to Descemet stripping automated endothelial keratoplasty or penetrating keratoplasty in both Fuchs endothelial corneal dystrophy and bullous keratopathy. Overall

cumulative graft survival was 97.4%, 78.4%, and 54.6% ($p<.001$) in Descemet membrane endothelial keratoplasty, Descemet stripping automated endothelial keratoplasty, and penetrating keratoplasty groups, respectively. In eyes with Fuchs endothelial corneal dystrophy, the graft survival was 98.7%, 96.2%, and 73.5% ($p=.009$) in Descemet membrane endothelial keratoplasty, Descemet stripping automated endothelial keratoplasty, and penetrating keratoplasty groups, respectively. In eyes with bullous keratopathy, the graft survival was 94.7%, 65.1%, and 47.0% ($p=.001$) in Descemet membrane endothelial keratoplasty, Descemet stripping automated endothelial keratoplasty, and penetrating keratoplasty groups, respectively. Graft rejection was lowest in eyes undergoing Descemet membrane endothelial keratoplasty (1.7% vs. Descemet stripping automated endothelial keratoplasty 5.0% vs. penetrating keratoplasty 14.1%; $p=.001$).

Descemet Stripping Endothelial Keratoplasty and Descemet Stripping Automated Endothelial Keratoplasty

Clinical Context and Therapy Purpose

The purpose of Descemet stripping endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as penetrating keratoplasty, in individuals with endothelial disease of the cornea.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with endothelial disease of the cornea. Diseases that affect the endothelial layer include Fuchs endothelial corneal dystrophy, aphakic and pseudophakic bullous keratopathy (corneal edema following cataract extraction), and failure or rejection of a previous corneal transplant.

Interventions

The therapy being considered is Descemet stripping endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty.

Comparators

Comparators of interest include penetrating keratoplasty.

Outcomes

The general outcomes of interest are change in disease status, morbid events, and functional outcomes. Relevant outcome measures include visual acuity, endothelial cell densities, patient satisfaction or quality of life, and complications including graft rejection, graft dislocation, and need for rebubble procedures. Follow-up generally occurs through 1 to 2 years post-surgery.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Systematic Reviews

In 2009, the American Academy of Ophthalmology performed a review of the safety and efficacy of Descemet stripping automated endothelial keratoplasty, identifying a level I study (RCT of pre-cut vs. surgeon dissected) along with 9 level II (well-designed observational studies) and 21 level III studies (mostly retrospective case series). (3) Although more than 2000 eyes treated with Descemet stripping automated endothelial keratoplasty were reported in different publications, most were reported by the same research group with some overlap in patients.

The main results of this review are as follows:

- Descemet stripping automated endothelial keratoplasty-induced hyperopia ranged from 0.7 to 1.5 diopters (D), with minimal induction of astigmatism (range, -0.4 to 0.6 D).
- The reporting of visual acuity was not standardized in studies reviewed. The average best-corrected visual acuity (BCVA) ranged from 20/34 to 20/66, and the percentage of patients seeing 20/40 or better ranged from 38% to 100%.
- The most common complication from Descemet stripping automated endothelial keratoplasty was posterior graft dislocation (mean, 14%; range, 0% to 82%), with a lack of adhesion of the donor posterior lenticule to the recipient stroma, typically occurring within the first week. It was noted that this percentage might have been skewed by multiple publications from a single research group with low complication rates. Graft dislocation required additional surgical procedures (rebubble procedures) but did not lead to sight-threatening vision loss in the articles reviewed.
- Endothelial graft rejection occurred in a mean of 10% of patients (range, 0% to 45%); most were reversed with topical or oral immunosuppression, with some cases progressing to graft failure. Primary graft failure, defined as unhealthy tissue that has not cleared within 2 months, occurred in a mean of 5% of patients (range, 0% to 29%). Iatrogenic glaucoma occurred in a mean of 3% of patients (range, 0% to 15%) due to a pupil block induced from the air bubble in the immediate postoperative period or delayed glaucoma from topical corticosteroid adverse events.
- Mean endothelial cell loss, which provides an estimate of long-term graft survival, was 37% at 6 months and 41% at 12 months. These percentages of cell loss were reported to be similar to those observed with penetrating keratoplasty.

Reviewers concluded that Descemet stripping automated endothelial keratoplasty appeared to be at least equivalent to penetrating keratoplasty regarding safety, efficacy, surgical risks, and complication rates, although long-term results were not yet available. The evidence also

indicated that endothelial keratoplasty is superior to penetrating keratoplasty regarding refractive stability, postoperative refractive outcomes, wound- and suture-related complications, and risk of intraoperative choroidal hemorrhage. The reduction in serious and occasionally catastrophic adverse events associated with penetrating keratoplasty has led to the rapid adoption of endothelial keratoplasty for treatment of corneal endothelial failure.

A Cochrane review of Descemet stripping automated endothelial keratoplasty compared to Descemet membrane endothelial keratoplasty for corneal endothelial failure was published in 2018. (4) The literature search identified 4 nonrandomized trials including 72 adult participants (144 eyes) who received Descemet stripping automated endothelial keratoplasty in the first eye followed by Descemet membrane endothelial keratoplasty in the fellow eye published between 2011 and 2015. All participants met criteria for Fuchs endothelial dystrophy and endothelial failure requiring a corneal transplant. Studies reported outcomes at various time points, including at 6, 12, and 6 to 24 months. At 1-year post-procedure, Descemet membrane endothelial keratoplasty resulted in better BCVA compared to Descemet stripping automated endothelial keratoplasty (mean difference, -0.14; 95% confidence interval [CI], -0.18 to -0.10 Logarithm of the Minimum Angle of Resolution [logMar]; low-certainty evidence). Two studies reported that Descemet membrane endothelial keratoplasty provided a higher cell density at 1 year. Graft dislocations requiring rebubbling were more common using Descemet membrane endothelial keratoplasty, although this difference could not be precisely estimated (relative risk [RR], 5.40; 95% CI, 1.51 to 19.3; very low-certainty evidence). The paired, contralateral eye studies in which Descemet stripping automated endothelial keratoplasty in 1 eye preceded Descemet membrane endothelial keratoplasty in the fellow eye for all patients was found to be at high-risk for bias due to potential unknown confounding factors.

Marques et al. (2019) conducted a meta-analysis of Descemet membrane endothelial keratoplasty compared to Descemet stripping automated endothelial keratoplasty for Fuchs endothelial dystrophy. (5) A literature search through August 2017 identified 10 retrospective studies of moderate methodological quality (N=947 eyes; 646 Descemet membrane endothelial keratoplasty). The primary outcome consisted of the mean difference in BCVA at 3-, 6-, and 12-months post-procedure. Secondary outcomes included rates of graft failure, rejection, rebubbling, endothelial cell density, subjective visual outcomes, and patient satisfaction. Best-corrected visual acuity was improved with Descemet membrane endothelial keratoplasty at all time points compared to Descemet stripping automated endothelial keratoplasty (12 months: 0.16 logMAR vs. 0.30 logMAR; $p<.001$). Descemet membrane endothelial keratoplasty had a 60% reduced rate of rejection (RR, 0.4; 95% CI, 0.24 to 0.67; $p=.0005$) but required more rebubbings (RR, 2.48; 95% CI, 1.32 to 4.64; $p=.005$). Descemet membrane endothelial keratoplasty had an increased number of primary graft failures and less endothelial cell density loss; however, these differences did not reach statistical significance. More patients reported being satisfied after Descemet membrane endothelial keratoplasty (odds ratio [OR], 10.29; 95% CI, 3.55 to 29.80; $p<.0001$).

Randomized Controlled Trials

Chamberlain et al. (2018) compared clinical outcomes of ultrathin-Descemet stripping automated endothelial keratoplasty with Descemet membrane endothelial keratoplasty in patients with damaged or diseased endothelium from Fuchs endothelial dystrophy or pseudophakic bullous keratopathy in the Descemet Endothelial Thickness Comparison Trial (DETECT). (6) The primary outcome measure was best spectacle-corrected visual acuity (BSCVA) at 6 months. Secondary outcomes included 3- and 12-month BSCVA, endothelial cell counts, and complications. The study included 50 eyes from 38 patients with 25 eyes randomized to each treatment arm. Compared to ultrathin Descemet stripping automated endothelial keratoplasty, Descemet membrane endothelial keratoplasty had superior visual acuity results. Best spectacle-corrected visual acuity was 1.5 lines better at 3 months (95% CI, 2.5 to 0.6 lines better; $p=.002$), 1.8 lines better at 6 months (95% CI, 2.8 to 1.0 lines better; $p<.001$), and 1.4 lines better at 12 months (95% CI, 2.2 to 0.7 lines better; $p<.001$). Average endothelial cell counts were 1855 cells/mm² in Descemet membrane endothelial keratoplasty and 2070 cells/mm² in ultrathin Descemet stripping automated endothelial keratoplasty at 12 months ($p=.051$). Intraoperative and postoperative complications rates were not statistically different between groups. Duggan et al. (2019) reported an update on corneal higher-order aberrations after ultrathin Descemet stripping automated endothelial keratoplasty versus Descemet membrane endothelial keratoplasty in DETECT. (7) In patients receiving Descemet membrane endothelial keratoplasty, the posterior corneal surface had significantly fewer coma aberrations ($p\le.003$) and total higher-order aberrations ($p\le.001$) at 3-, 6-, and 12-months post-surgery compared to ultrathin Descemet stripping automated endothelial keratoplasty. Descemet membrane endothelial keratoplasty was found to decrease whereas ultrathin Descemet stripping automated endothelial keratoplasty was found to increase posterior corneal higher-order aberrations compared with presurgical values, potentially accounting for the better visual acuity observed with Descemet membrane endothelial keratoplasty. Further, Lin et al. (2024) conducted a secondary analysis of DETECT data to investigate mediators of visual acuity with Descemet membrane endothelial keratoplasty and ultrathin-Descemet stripping automated endothelial keratoplasty. (8) Results showed that 64% of the observed differences in visual acuity at 24 months were mediated by reductions in posterior higher-order aberrations with Descemet membrane endothelial keratoplasty group vs ultrathin-Descemet stripping automated endothelial keratoplasty, emphasizing the importance of posterior optical quality in determining visual outcomes. Hirabayashi et al. (2020) reported on an update of corneal light scatter outcomes as measured by densitometry in DETECT. (9) Both Descemet membrane endothelial keratoplasty and ultrathin Descemet stripping automated endothelial keratoplasty were found to improve the degree of corneal light scatter after surgery, with no differences between groups observed at 12 months post-surgery. Similar results were shown by Lin et al. (2024) at 24 months. (8)

Dunker et al. (2020) published the results of a prospective, multicenter RCT comparing the efficacy of ultrathin Descemet stripping automated endothelial keratoplasty (n=25) versus Descemet membrane endothelial keratoplasty (n=29) in patients with Fuchs endothelial corneal dystrophy. (10) Fifty-four patients were enrolled from 6 corneal centers in the Netherlands. There was no significant difference in BSCVA at 3 ($p=.15$), 6 ($p=.20$), or 12 months post-surgery ($p=.06$), between study arms. However, the percentage of eyes achieving 20/25 Snellen vision

was significantly higher with Descemet membrane endothelial keratoplasty at 12 months (p=.02).

Observational Studies

Wilhelm et al. (2025) conducted a retrospective study involving 2956 first-time keratoplasty procedures over a 10-year period to compare outcomes among Descemet membrane endothelial keratoplasty (n=20 at 10-year follow-up), Descemet stripping automated endothelial keratoplasty (n=9 at 10-year follow up), and penetrating keratoplasty (n=51 at 10-year follow up) for patients with Fuchs endothelial corneal dystrophy. (11) The study showed that Descemet membrane endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty provided faster visual recovery, with median times to achieve a BSCVA of $\geq 6/12$ being 7.8 months and 12.4 months, respectively, compared to 37.9 months for penetrating keratoplasty. However, penetrating keratoplasty demonstrated superior long-term graft survival, with a 92% survival rate at 10 years, compared to 75% for Descemet membrane endothelial keratoplasty and 73% for Descemet stripping automated endothelial keratoplasty. Endothelial cell density declined more rapidly following Descemet membrane endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty, which contributed to lower graft survival. The probability of maintaining an endothelial cell density greater than 1,000 cells per square millimeter at 10 years was 18% for penetrating keratoplasty, 8% for Descemet stripping automated endothelial keratoplasty, and 3% for Descemet membrane endothelial keratoplasty. Additionally, Descemet membrane endothelial keratoplasty exhibited the lowest rejection rate, with 10% at 10 years, compared to 13% for penetrating keratoplasty and 19% for Descemet stripping automated endothelial keratoplasty.

Fuest et al. (2017) compared 5-year visual acuity outcomes in patients receiving Descemet stripping automated endothelial keratoplasty (n=423) or penetrating keratoplasty (n=405) in the Singapore Cornea Transplant Registry. (12) Mean age of patients was 67 years. The Descemet stripping automated endothelial keratoplasty group had a higher percentage of Chinese patients, a higher percentage of patients with Fuchs endothelial dystrophy, and a lower percentage of patients with bullous keratopathy than the penetrating keratoplasty group. Controlling for preoperative BSCVA, which differed significantly between groups, patients receiving Descemet stripping automated endothelial keratoplasty experienced significantly better vision through 3 years of follow-up than patients undergoing penetrating keratoplasty. Four- and 5-year follow-up measures showed similar BSCVA among both treatment groups. Subgroup analyses by Fuchs endothelial dystrophy and bullous keratopathy showed similar patterns of significantly better vision through the first 3 years of follow-up in patients receiving Descemet stripping automated endothelial keratoplasty than in patients receiving penetrating keratoplasty.

Heinzelmann et al. (2016) reported on 2-year outcomes in patients who underwent endothelial keratoplasty or penetrating keratoplasty for Fuchs endothelial dystrophy or bullous keratopathy. (13) The study included 89 eyes undergoing Descemet stripping automated endothelial keratoplasty and 329 eyes undergoing penetrating keratoplasty. The postoperative visual improvement was faster after endothelial keratoplasty than after penetrating

keratoplasty. For example, among patients with Fuchs endothelial dystrophy, 50% of patients achieved a BSCVA of Snellen 6/12 or more 18 months after Descemet stripping automated endothelial keratoplasty versus more than 24 months after penetrating keratoplasty.

Endothelial cell loss was similar after endothelial keratoplasty and penetrating keratoplasty in the early postoperative period. However, after an early decrease, endothelial cell loss stabilized in patients who received endothelial keratoplasty whereas the decrease continued in those who had penetrating keratoplasty. Among patients with Fuchs endothelial dystrophy, there was a slightly increased risk of late endothelial failure in the first 2 years with endothelial keratoplasty than with penetrating keratoplasty. Graft failure was reported to be lower among patients with bullous keratopathy compared with patients with Fuchs endothelial dystrophy (numbers not reported).

Longer-term outcomes have been reported in several studies. Five-year outcomes from a prospective study conducted at the Mayo Clinic were published by Wacker et al. (2016). (14) The study included 45 participants (52 eyes) with Fuchs endothelial dystrophy who underwent Descemet stripping endothelial keratoplasty. Five-year follow-up was available for 34 (65%) eyes. Mean high-contrast BSCVA was 20/56 Snellen equivalent pre-surgery and decreased to 20/25 Snellen equivalent at 60 months. The difference in high-contrast BSCVA at 5 years versus pre-surgery was statistically significant ($p<.001$). Similarly, the proportion of patients with BSCVA of 20/25 Snellen equivalent or better increased from 26% at 1-year post-surgery to 56% at 5 years ($p<.001$). There were 6 graft failures during the study period (4 failed to clear after surgery, 2 failed during follow-up). All patients with graft failures were regrafted.

Previously, 3-year outcomes after Descemet stripping automated endothelial keratoplasty were reported by an eye institute. (15) This retrospective analysis (2012) included 108 patients who underwent Descemet stripping automated endothelial keratoplasty for Fuchs endothelial dystrophy or pseudophakic bullous keratopathy and had no other ocular comorbidities. Best spectacle-corrected visual acuity was measured at 6 months and 1, 2, and 3 years. Best spectacle-corrected visual acuity after Descemet stripping automated endothelial keratoplasty improved over 3 years of follow-up. The percentage of patients who reached a BSCVA of 20/20 or greater was 0.9% at baseline, 11.1% at 6 months, 13.9% at 1 year, 34.3% at 2 years, and 47.2% at 3 years. Ninety-eight percent of patients reached a BSCVA of 20/40 or greater by 3 years. Tables 1 and 2 describe the characteristics and results of key nonrandomized trials.

Table 1. Summary of Key Nonrandomized Trial Characteristics

Study	Study Type	Country	Dates	Participant N Eyes	DSAEK, N	PK, N	DMEK, N	Follow-Up
Wilhelm et al. (2025) (11)	Retrospective	Germany	2003-2023	Total 2956	80	204	2672	10 yrs (n=80)
Fuest et al. (2017) (12)	Prospective	Singapore	1991-2011	Total 828	423	405	NR	5 yrs

Heinzelman et al. (2016) (13)	Cohort	Germany	2011-2014	Total 868	89	329	450	2 yrs
Wacker et al. (2016) (14)	Prospective	U.S.	2006-2010	Total 52	34	NR	NR	5 yrs (n=34, 65%)
Li et al. (2012) (15)	Retro-spective	U.S.	2005-2007	Total 207	108	NR	NR	3 yrs

DMEK: Descemet membrane endothelial keratoplasty; DSAEK: Descemet stripping automated endothelial keratoplasty; NR: not reported; PK=penetrating keratoplasty; U.S.: United States; yrs: years; N: number.

Table 2. Summary of Key Nonrandomized Trial Results

Study	BSCVA < 6/12 Snellen at 10 years	Endothelial cell density > 1000 cells/mm ² at 10 years	Graft Survival at 10 years			
Wilhelm et al. (2025) (11)						
DMEK (n=20 at 10 years)	1%	3%	75%			
DSAEK (n=9 at 10 years)	NR	8%	73%			
PK (n=51 at 10 years)	22%	18%	92%			
Study	BSCVA	SE		Cylinder		
Fuest et al. (2017) (12)	at 5-years (n); mean (SD); p-value	at 5-years (n); mean (SD); p-value		at 5-years (n); mean (SD); p-value		
Total	(N=89); 0.62 (0.6); p=0.037	(N=62); -1.7 (2.7); P=0.017		(N=62); -3.1 (2.1); P<0.001		
DSAEK	(n=25); 0.46(0.5); p=0.037	(n=18); -0.8 (1.7); P=0.017		(n=18); -1.6 (1.1); P<0.001		
PK	(n=25); 0.63 (0.6); p=0.037	(n=44); -2.1 (2.9); P=0.017		(n=44); -3.75 (2.1); P=0.001		

Study	% of BSCVA of Snellen 6/7.5 or better at 24-months		Chronic endothelial cell loss >500 cells/mm ² at 15 months		Chronic endothelial cell loss >500 cells/mm ² at 24 months	
Heinzelmann et al. (2016) (13)						
FED DMEK	53%		95%		NR	
FED DSAEK	15%		93%			
FED PK	10%		99%		NR	
BK DMEK	NR		NR		NR	
BK DSAEK	NR		NR		NR	
BK PK	NR		NR		90%	
Study	Mean high-contrast BSCVA presurgery	Mean high-contrast BSCVA at 5-years				
Wacker et al. (2016) (14)						
FECD DSEK	20/56	20/25				
Study	% of eyes achieving a BSCVA of 20/40 at 3-years	% of eyes achieving a BSCVA of 20/30 at 3-years		% of eyes achieving a BSCVA of 20/25 at 3-years		% of eyes achieving a BSCVA of 20/20 at 3-years
Li et al. (2012) (15)						
FED+BK DSAEK	98.1% (n=106)	90.7% (n=98)		70.4% (n=76)		47.2% (n=51)

BK: bullous keratopathy; BSCVA: best spectacle-corrected visual acuity; DMEK: Descemet membrane endothelial keratoplasty; DSAEK: Descemet stripping automated endothelial keratoplasty; DSEK: Descemet stripping endothelial keratoplasty; FED/FECD: Fuchs endothelial corneal dystrophy; NR: not reported; N: number of eyes; PK: penetrating keratoplasty; SD: standard deviation; SE: spherical equivalent.

Section Summary: Descemet Stripping Endothelial Keratoplasty and Descemet Stripping Automated Endothelial Keratoplasty

Evidence for the use of Descemet stripping endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty consists of systematic reviews and several large observational studies with follow-up extending from 2 to 10 years. The review and the studies showed that patients undergoing Descemet stripping endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty experience greater improvements in visual acuity than patients undergoing penetrating keratoplasty. Also, patients undergoing Descemet stripping endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty

experienced significantly fewer serious adverse events than patients undergoing penetrating keratoplasty.

Descemet Membrane Endothelial Keratoplasty and Descemet Membrane Automated Endothelial Keratoplasty

Clinical Context and Therapy Purpose

The purpose of Descemet membrane endothelial keratoplasty and Descemet membrane automated endothelial keratoplasty is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as penetrating keratoplasty, in individuals with endothelial disease of the cornea.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with endothelial disease of the cornea. Diseases that affect the endothelial layer include Fuchs endothelial dystrophy, aphakic and pseudophakic bullous keratopathy (corneal edema following cataract extraction), and failure or rejection of a previous corneal transplant.

Interventions

The therapy being considered is Descemet membrane endothelial keratoplasty and Descemet membrane automated endothelial keratoplasty. It has been suggested that by eliminating the stroma on the donor tissue, Descemet membrane endothelial keratoplasty and Descemet membrane automated endothelial keratoplasty may reduce stromal interface haze and provide better visual acuity outcomes than Descemet stripping endothelial keratoplasty or Descemet stripping automated endothelial keratoplasty. (16, 17)

Comparators

Comparators of interest include penetrating keratoplasty.

Outcomes

The general outcomes of interest are change in disease status, morbid events, and functional outcomes. Relevant outcome measures include visual acuity, endothelial cell densities, patient satisfaction or quality-of-life, and complications including graft rejection, graft dislocation, and need for rebubble procedures. Follow-up generally occurs through 1 to 2 years post-surgery.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.

- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Systematic Reviews

The American Academy of Ophthalmology conducted a systematic review of the safety and outcomes of Descemet membrane endothelial keratoplasty and investigated whether Descemet membrane endothelial keratoplasty offered any advantages over Descemet stripping endothelial keratoplasty (Deng et al. [2018]). (18) The literature search, conducted through May 2017, identified 47 studies for inclusion. Quality was assessed using a scale from the Oxford Centre for Evidence-Based Medicine. Two studies were rated level I evidence (well-designed and well-conducted RCTs), 15 studies were level II (well-designed case-control or cohort studies or RCTs with methodologic deficits), and 30 studies were level III (case series, case reports, or poor-quality cohort or case-control). Mean length of follow-up among the studies ranged from 5 to 68 months. A BSCVA of 20/25 was achieved by 33% to 67% of patients (5 studies). A BSCVA of 20/20 was achieved by 29% to 32% (3 studies) at 3 months postsurgery and by 17% to 67% at 6 months postsurgery. Seven studies, 6 of which were rated as level II evidence, directly compared Descemet membrane endothelial keratoplasty with Descemet stripping endothelial keratoplasty and all 7 showed a faster visual recovery and a better visual outcome after Descemet membrane endothelial keratoplasty compared with Descemet stripping endothelial keratoplasty. The rate of endothelial cell loss, graft failure, and intraoperative and postoperative complications was similar between Descemet membrane endothelial keratoplasty and Descemet stripping endothelial keratoplasty.

Singh et al. (2017) conducted a systematic review and meta-analysis of studies comparing Descemet membrane endothelial keratoplasty with Descemet stripping endothelial keratoplasty or Descemet stripping automated endothelial keratoplasty. (19) The literature search, conducted through May 2016, identified 9 studies for inclusion in the qualitative analysis and 7 studies for inclusion in the meta-analysis. A quality assessment of studies was not presented. Meta-analyses of 343 eyes showed that the 6-month mean difference in BSCVA was significantly better in patients undergoing Descemet membrane endothelial keratoplasty than in patients undergoing Descemet stripping endothelial keratoplasty (-0.13; 95% CI, -0.16 to -0.09). The 6-month mean difference in endothelial cell density (n=348) did not differ significantly between groups (76.8; 95% CI, -79.8 to 233.4), though the interpretation of this result is limited due to high heterogeneity. A higher rate of air injection/rebubbling was reported among patients in the Descemet membrane endothelial keratoplasty group compared with the Descemet stripping endothelial keratoplasty group.

Pavlovic et al. (2017) conducted a meta-analysis of 11 studies comparing Descemet membrane endothelial keratoplasty (n=350) with Descemet stripping automated endothelial keratoplasty (n=373). (20) The date of the literature search and quality assessment methods were not reported. The mean difference in BSCVA did not differ significantly at the 3-month follow-up (-0.12; 95% CI, -0.28 to 0.04), but was significantly better in the Descemet membrane endothelial keratoplasty group than in the Descemet stripping automated endothelial

keratoplasty group at both the 6-month (-0.12; 95% CI, -0.15 to -0.10) and at the 6-month and beyond follow-ups (-0.13; 95% CI, -0.17 to -0.09). There were no statistical differences in endothelial cell loss between the 2 procedures at 6 (mean difference, 0.2; 95% CI, -5.6 to 6.1) or 12 months (mean difference, 3.6; 95% CI, -3.7 to 10.9). There were more graft rejections reported among patients in the Descemet stripping automated endothelial keratoplasty group compared with those in the Descemet membrane endothelial keratoplasty group, but the difference was not significant (OR, 2.7; 95% CI, 0.6 to 11.9). There were more graft failures reported in the Descemet membrane endothelial keratoplasty group compared with the Descemet stripping automated endothelial keratoplasty group, but this difference, too, was not significant (OR, 2.8; 95% CI, 0.7 to 10.6).

Li et al. (2017) conducted a systematic review and meta-analysis comparing Descemet membrane endothelial keratoplasty and Descemet stripping endothelial keratoplasty. (21) The literature search, conducted through January 2017, identified 19 studies for inclusion: 15 retrospective control studies, a prospective nonrandomized case series, and 3 for which the study designs could not be determined from the meeting abstracts. A modified version of the Newcastle-Ottawa Scale was used to assess the quality of the studies. Eight items relating to selection, comparability, and outcome were assessed, and if a study received a score greater than 6, it was considered relatively high quality. Two studies had a score of 7, 8 studies had a score of 6, 3 studies had a score of 5, and 6 studies had a score of 4. A total of 2378 eyes were included in the studies, 1124 receiving Descemet membrane endothelial keratoplasty and 1254 receiving Descemet stripping endothelial keratoplasty. Meta-analyses of 13 studies showed an overall mean difference in BSCVA that was significantly improved in the Descemet membrane endothelial keratoplasty group compared with the Descemet stripping endothelial keratoplasty group (-0.15; 95% CI, -0.19 to -0.11). This significant mean difference in BSCVA was seen at the 3-, 6-, and 12-month follow-ups. Meta-analyses, which included 354 Descemet membrane endothelial keratoplasty and 313 Descemet stripping endothelial keratoplasty eyes (N=667), showed no significant difference in endothelial cell density between groups (mean difference, 14.9; 95% CI, -181.5 to 211.3). The most common complication in both procedures was partial or total graft detachment, with significantly more occurrences in the Descemet membrane endothelial keratoplasty group than in the Descemet stripping endothelial keratoplasty group (OR, 4.6; 95% CI, 2.4 to 8.6).

Wu et al. (2021) conducted a systematic review and meta-analysis comparing Descemet membrane endothelial keratoplasty and Descemet stripping endothelial keratoplasty after failed penetrating keratoplasty. (22) A literature search was conducted through July 10, 2020, and included 25 studies (16 Descemet stripping endothelial keratoplasty; 9 Descemet membrane endothelial keratoplasty) for inclusion: 22 retrospective cohort studies and 3 prospective cohort studies. There was a total of 970 patients enrolled with 989 total eyes included in this review. The mean visual acuity of the Descemet stripping endothelial keratoplasty and Descemet membrane endothelial keratoplasty groups were 0.65 ± 0.18 and 0.43 ± 0.23 logMAR, respectively, at 6 months postoperatively. This shows a general trend for improved visual acuity following both Descemet stripping endothelial keratoplasty and

Descemet membrane endothelial keratoplasty after failed penetrating keratoplasty. Graft survival and rejection rates were comparable between the two groups.

Maier et al. (2023) conducted a systematic review and meta-analysis comparing Descemet membrane endothelial keratoplasty and ultrathin Descemet stripping automated endothelial keratoplasty. (23) A literature search was conducted through June 2022, and included 7 studies: 3 RCTs, 1 prospective case series, 1 retrospective comparative study, and 2 retrospective cohort studies. The primary outcome assessed was BSCVA and secondary outcomes included endothelial cell density and postoperative complications. Baseline BSCVA data consisted of 163 eyes treated with Descemet membrane endothelial keratoplasty and 165 eyes treated with ultrathin Descemet stripping automated endothelial keratoplasty. The BSCVA standardized mean difference (SMD) between groups after 3 months was 0.49 (95% CI, 0.22 to 0.76; $p=.0004$) and after 12 months was 0.50 (95% CI, 0.27 to 0.74; $p=.0001$); this favored Descemet membrane endothelial keratoplasty. Data at 6 months could not be evaluated due to high heterogeneity of the studies. Another significant outcome between groups was the re-bubbling rate after Descemet membrane endothelial keratoplasty compared to ultrathin Descemet stripping automated endothelial keratoplasty (RR, 0.33; 95% CI, 0.15 to 0.67; $p=.0025$). All other measured outcomes were not significantly different between groups. Tables 3 and 4 describe the characteristics and results of key systematic reviews and meta-analyses.

Table 3. SR & M-A Characteristics

Study	Dates	Trials	N (Eyes)	Intervention	N (Range)	Design	Duration
Deng et al. (2018) (18)	NR-05/2017	47	9046; patients with corneal endothelial dysfunction	DMEK	9046 (25-905)	RCT; case control and cohort; case series, case reports	5.3-68 months
Singh et al. (2017) (19)	NR-05/2016	9	586	DMEK, DSAEK	586 (20-155)	NR	NR
Pavlovic et al. (2017) (20)	NR	11	723	DMEK (n=350); DSAEK (n=373)	NR	NR	NR
Li et al. (2017) (21)	NR-01/2017	19	2378	DMEK; DSEK	2378 (20-739)	NR	3.1-22.55 months
Wu et al. (2021) (22)	NR-07/2020	25	989	DMEK, DSAEK	989 (7-246)	prospective and	6-36.1 months

						retrospective cohorts	
Maier et al. (2023) (23)	NR-06/2022	7	328	DMEK; UT-DSAEK	NR	RCT; case series; retrospective cohorts	NR

DMEK: Descemet membrane endothelial keratoplasty; DSAEK: Descemet stripping automated endothelial keratoplasty; DSEK: Descemet stripping endothelial keratoplasty; M-A: meta-analysis; N: number; NR: not reported; RCT: randomized controlled trial; SR: systematic review; UT-DSAEK: ultrathin Descemet stripping automated endothelial keratoplasty.

Table 4. SR & M-A Results

Study	Mean BCVA	Mean endothelial cell loss at time	Change in SE	Minimal induced astigmatism
Deng et al. (2018) (18)				
Total N*=9046	Range: 20/21 to 20/31	33% (range, 25% to 47%) [6-months]	+0.43 D (range, -1.17 to +1.2 D)	+0.03 D (range, -0.03 to +1.11 D)
	BCVA at 6 months	ECD at 6 months	Graft Detachment overall	Graft Rejection
Singh et al. (2017) (19)				
After DMEK, mean; SD, p-value	0.161; 0.129; p<0.0001; n=184	1855; 442; p=0.708	NR	NR
After DSAEK, mean; SD, p-value	0.293; 0.153 P<0.0001; n=159	1872; 429; p=0.708	NR	NR
Pooled mean difference (CI, SD)	-0.13 (95% CI, 0.16 to 0.09); N=343	Could not be interpreted due to high statistical heterogeneity	NR	NR
Pavlovic et al. (2017) (20)				
MD between DSAEK and DMEK group	-0.12; 95% CI, -0.15 to -0.10	0.2; 95% CI, -5.6 to 6.1	Not available	Not available
Li et al. (2017) (21)				
Comparison between DMEK and DSEK (MD [95% CI] % weight)	-0.13 (-0.17 to -0.08) 51.29	25.59 (-183.15 to 234.32) p=0.810	4.56 (2.43 to 8.58)	-0.04 (-0.08 to -0.002)

Pooled mean difference (CI, SD) overall	-0.15 (-0.19 to -0.11) P<0.001	14.88 (-181.5 to 211.27) P=0.882	NR	NR
Wu et al. (2021) (22)				
After DMEK, mean; SD	0.43; 0.23; n=243	47.6% (range 37.1% to 61.4%) [12-months]	NR	NR
After DSAEK, mean; SD	0.65; 0.18; n=746	NR	NR	NR
	BCVA at 12 months	ECD at 6 months	Graft detachment overall	Graft rejection
Maier et al. (2023) (23)				
Comparison between DMEK and UT-DSAEK groups	MD, 0.50; (95% CI, 0.27 to 0.74); p<.0001	Could not be interpreted due to high statistical heterogeneity	RR, 0.33 (95% CI, 0.16 to 0.67); p=.0025	RR, 1.4 (95% CI, 0.27 to 7.30); p=.69

*N=eyes

BCVA: best corrected visual acuity; CI: confidence interval; D: diopters; DMEK: Descemet membrane endothelial keratoplasty; DSAEK: Descemet stripping automated endothelial keratoplasty; ECD: endothelial corneal dystrophy; ECL: endothelial cell loss; MD: mean difference; NR: not reported; N: number of eyes; SD: standard deviation; SE: spherical equivalent; SR: systematic review; M-A: meta-analysis; UT-DSAEK: ultrathin Descemet stripping automated endothelial keratoplasty; DSEK: Descemet stripping endothelial keratoplasty.

Observational Studies

Oellerich et al. (2017) reported on 6-month outcomes of a large cohort of patients undergoing Descemet membrane endothelial keratoplasty by 55 surgeons from 23 countries. (24) Outcomes of interest were BSCVA, a decrease in endothelial cell density, and complications. Subgroup analyses were conducted by the number of procedures performed by the surgeon (1 to 24 [39%], 25 to 99 [38%], and ≥ 100 [23%]). In the total population, 91% of patients achieved BSCVA improvement, with 5% experiencing no change and 5% experiencing deterioration in visual acuity. Subgroup analyses showed that the proportion of patients achieving BSCVA improvement did not differ significantly between patients whose surgeons had performed 100 or more procedures and those whose surgeons had performed fewer than 25 procedures. Nine percent of patients experienced intraoperative complications, with the rate decreasing significantly as the surgeon performed more procedures. The most frequent postoperative complication was partial graft detachment (27%), which also decreased significantly with surgeon experience. Rates of other postoperative complications such as graft failure, cataract, and glaucoma did not differ based on surgeon experience.

Tourtas et al. (2012) conducted a retrospective comparison of 38 consecutive patients/eyes that underwent Descemet membrane endothelial keratoplasty and 35 consecutive patients/eyes who had undergone Descemet stripping automated endothelial keratoplasty. (25) Only patients with Fuchs endothelial dystrophy or pseudophakic bullous keratopathy were included. After Descemet membrane endothelial keratoplasty, 82% of eyes required rebubbling. After Descemet stripping automated endothelial keratoplasty, 20% of eyes required rebubbling. Best spectacle-corrected visual acuity in both groups was comparable at baseline (Descemet membrane endothelial keratoplasty=0.70 logMAR; Descemet stripping automated endothelial keratoplasty=0.75 logMAR). At 6-month follow-up, mean visual acuity improved to 0.17 logMAR after Descemet membrane endothelial keratoplasty and 0.36 logMAR after Descemet stripping automated endothelial keratoplasty. This difference was statistically significant. At 6 months following surgery, 95% of Descemet membrane endothelial keratoplasty treated eyes reached a visual acuity of 20/40 or better, and 43% of Descemet stripping automated endothelial keratoplasty treated eyes reached a visual acuity of 20/40 or better. Endothelial cell density decreased by a similar amount after both procedures (41% after Descemet membrane endothelial keratoplasty, 39% after Descemet stripping automated endothelial keratoplasty).

van Dijk et al. (2013) reported on outcomes of their first 300 consecutive eyes treated with Descemet membrane endothelial keratoplasty. (26) Indications for Descemet membrane endothelial keratoplasty were Fuchs endothelial dystrophy, pseudophakic bullous keratopathy, failed penetrating keratoplasty, or failed endothelial keratoplasty. Of the 142 eyes evaluated for visual outcomes at 6 months, 79% reached a BSCVA of 20/25 or more, and 46% reached a BSCVA of 20/20 or more. Endothelial cell density measurements at 6 months were available in 251 eyes. Average cell density was 1674 cells/mm², representing a decrease of 34.6% from preoperative donor cell density. The major postoperative complication in this series was graft detachment requiring rebubbling or regraft, which occurred in 10.3% of eyes. Allograft rejection occurred in 3 eyes (1%), and intraocular pressure was increased in 20 (6.7%) eyes. Except for 3 early cases that may have been prematurely regrafted, all but 1 eye with an attached graft cleared in 1 to 12 weeks.

A 2009 review of cases from another group in Europe suggested that a greater number of patients achieve 20/25 vision or better with Descemet membrane endothelial keratoplasty. (27) Of the first 50 consecutive eyes, 10 (20%) required a secondary Descemet stripping endothelial keratoplasty for failed Descemet membrane endothelial keratoplasty. For the remaining 40 eyes, 95% had a BSCVA of 20/40 or better, and 75% had a BSCVA of 20/25 or better. Donor detachments and primary graft failure with Descemet membrane endothelial keratoplasty were problematic. In 2011, this group reported on the surgical learning curve for Descemet membrane endothelial keratoplasty, with their first 135 consecutive cases retrospectively divided into 3 subgroups of 45 eyes each. (28) Graft detachment was the most common complication, which decreased with surgeon experience. In their first 45 cases, a complete or partial graft detachment occurred in 20% of cases, compared with 13.3% in the second group, and 4.4% in the third group. Clinical outcomes in eyes with normal visual potential and a functional graft (n=110) were similar across the 3 groups, with an average

endothelial cell density of 1747 cells/mm² and 73% of cases achieving a BSCVA of 20/25 or better at 6 months.

A North American group reported on 3-month outcomes from a prospective consecutive series of 60 cases of Descemet membrane endothelial keratoplasty in 2009, and in 2011, they reported on 1-year outcomes from these 60 cases plus an additional 76 cases of Descemet membrane endothelial keratoplasty. (29, 30) Preoperative BSCVA averaged 20/65 (range of 20/20 to counting fingers). Sixteen eyes were lost to follow-up, and 12 (8.8%) grafts had failed. For the 108 grafts examined and found to be clear at 1 year, 98% achieved a BSCVA of 20/30 or better. Endothelial cell loss was 31% at 3 months and 36% at 1 year. Although visual acuity outcomes appeared to be improved over a Descemet stripping automated endothelial keratoplasty series from the same investigators, preparation of the donor tissue and attachment of the endothelial graft were more challenging. A 2012 cohort study by this group found reduced transplant rejection with Descemet membrane endothelial keratoplasty. (31) One (0.7%) of 141 patients in the Descemet membrane endothelial keratoplasty group had a documented episode of rejection compared with 54 (9%) of 598 in the Descemet stripping endothelial keratoplasty group and 5 (17%) of 30 in the penetrating keratoplasty group.

The same group also reported on a prospective consecutive series (2011) of their initial 40 cases (36 patients) of Descemet membrane automated endothelial keratoplasty (microkeratome dissection and a stromal ring). (32) Indications for endothelial keratoplasty were Fuchs endothelial dystrophy (87.5%), pseudophakic bullous keratopathy (7.5%), and failed endothelial keratoplasty (5%). Air was reinjected in 10 (25%) eyes to promote graft attachment; 2 (5%) grafts failed to clear and were successfully regrafted. Compared with a median BSCVA of 20/40 at baseline (range, 20/25 to 20/400), median BSCVA at 1 month was 20/30 (range, 20/15 to 20/50). At 6 months, 48% of eyes had 20/20 vision or better, and 100% had 20/40 or better. Mean endothelial cell loss at 6 months relative to baseline donor cell density was 31%.

Tables 5 and 6 describe key characteristics and results of these observational studies.

Table 5a. Summary of Key Observational Study Characteristics

Study	Study Design	Country	Dates	Participants
Oellerich et al. (2017) (24)	Retrospective cohort	Europe, Asia, Africa, North America, South America, Australia	Aug 2008-July 2015	Mean age, 69.8 yrs ± 11.0 (range, 16-99 yrs); 37% male, 57.9% female, 5.2% not specified; 74.4% FED, 16.8% BK; 7.6% failed transplant, 0.9% other; 0.3% not specified
van Dijk et al. (2013) (26)	Prospective	Netherlands	NR	Mean age 67 yrs ± 13 (range, 30-93 yrs); 166 female/134 male; FED=272 patients; BK=17 patients; Failed DSEK/PK=9/1 patients

Tourtas et al. (2012) (25)	Retrospective cohort	Germany	Aug 2009-Dec 2009; DSAEK: Aug 2008-Mar 2009	DMEK: mean age, 68.3 yrs \pm 9 (range, 42-85 yrs), 16 female/22 male; DSAEK: mean age, 68.1 \pm 11 (range, 48-87 yrs), 20 female/15 male
Ham et al. (2009) (27)	Prospective case	Netherlands	NR	Patients with FED; 23 men, 27 women; age range, 41-88 yrs;
Dapenda et al. (2011) (28)	Retrospective	Netherlands	Feb 2005-Dec 2010	118 patients with FED, 49 male, 69 female; age range, 33-93 yrs
Price et al. (2009) (29)	Prospective	U.S.	Feb 2009-Oct 2009	58 patients with FED, PK, or failed previous graft; mean age, 68 yrs \pm 9.9 (range, 48-85 yrs); 34 female/26 male
Guerra et al. (2011) (30)	Prospective	U.S.	Feb 2009-Oct 2009	112 patients with FED, PK, or failed previous graft; mean age, 78 yrs \pm 10.36; 72 female/40 male
McCauley et al. (2011) (32)	Prospective	U.S.	NR	36 patients treated with DMAEK. Mean age, 69 yrs (range, 48-88 yrs); 53% female
Anshu et al. (2012) (31)	Comparative	U.S.	Feb 2009-Oct 2009	Patients undergoing DMEK compared retrospectively with matched cohort undergoing DSEK and PK, treated at same center, with similar demographics, follow-up, duration, indications for surgery

BK: bullous keratopathy; DMAEK: Descemet membrane automated endothelial keratoplasty; DMEK: Descemet membrane endothelial keratoplasty; DSAEK: Descemet stripping automated endothelial keratoplasty; DSEK: Descemet stripping endothelial keratoplasty; FED/FECD: Fuchs endothelial corneal dystrophy; N=eyes except where indicated otherwise; NR: not reported; PK: penetrating keratoplasty; SD: standard deviation; yr(s) years; U.S.: United States.

Table 5b. Summary of Key Observational Study Characteristics

Study	Treatment 1	Treatment 2	Follow- Up
Oellerich et al. (2017) (24)	DMEK (N=2448)	NA	6 mos
van Dijk et al. (2013) (26)	DMEK (N=300)	NA	6 mos

Tourtas et al. (2012) (25)	DMEK (n=38)	DSAEK (n=35)	6 mos
Ham et al. (2009) (27)	DMEK (n=40)	DMEK followed by DSEK as a back-up procedure in the event of DMEK graft failure (n=10)	6 mos
Dapenda et al. (2011) (28)	DMEK (N=135)	NA	6 mos
Price et al. (2009) (29)	DMEK (N=60)	NA	3 mos
Guerra et al. (2011) (30)	DMEK (N=136)	NA	1 yr
McCauley et al. (2011) (32)	DMAEK (N=40)	NA	6 mos
Anshu et al. (2012) (31)	DMEK/DSEK (n=739)	PK (n=30)	2 yrs

DMAEK: Descemet membrane automated endothelial keratoplasty; DMEK: Descemet membrane endothelial keratoplasty; DSAEK: Descemet stripping automated endothelial keratoplasty; DSEK: Descemet stripping endothelial keratoplasty; N=eyes except where indicated otherwise; NA: not applicable; PK: penetrating keratoplasty; mo(s): months; yr(s): years.

Table 6. Summary of Key Observational Study Results

Study	BCVA pre-operative	BCVA 6 mos FU	ECD pre-operative mean +/- SD (cells/mm ²)	ECD 6 mos FU mean +/- SD (cells/mm ²)	Post-operative complications
Oellerich et al. (2017) (24)	N=2430	N=1959	N=1956	N=1405	N=2363
DMEK	n (%) ≥20/25 Snellen=46.17 (1.9%)	n (%) ≥20/25 Snellen=889 (45.4%)	2635 +/- 294	1575 +/- 489	647 (27.4%) (for all types of post-operative complications)
van Dijk et al. (2013) (26)	N=221	N=221	N=251	N=251	N=300
DMEK	n (%) ≥20/25 Snellen=16 (7%)	n (%) ≥20/25 Snellen=175 (79%)	2561 +/- 198	1674 +/- 518	31 (10%) for most frequent complication, (partial) graft detachment

Tourtas et al. (2012) (25)	N=73	N=73	N=73	N=73	N=73
DMEK (n=38)	Mean +/- SD; 0.70 +/- 0.48 logMAR	Mean +/- SD; 0.17 +/- 0.12 logMAR (n=38)	2575 +/- 260	1520 +/- 299	31 (82%) required air injections for partial dehiscence of the EDM
DSAEK (n=35)	n +/- SD; 0.75 +/- 0.32 logMAR	n +/- SD; 0.36 +/- 0.15 logMAR (n=35)	2502 +/- 220	1532 +/- 495	7 (20%) required air injections for partial dehiscence of the EDM
Ham et al. (2009) (27)					
Pooled (N=50)	NR	n (%) ≥20/25 Snellen=47 (66%)	2623 +/- 193 (n=43)	1815 +/- 578 (n=43)	All complications, n=14 (28%)
DMEK only (n=40)	NR	n (%) ≥20/25 Snellen=30 (75%)	2618 +/- 201 (n=35)	1876 +/- 522 (n=35)	NR
Dapena et al. (2011) (28)	N=135	N=110	N=135	174 +/- 527 (n=106)	Primary graft failure (2.2%, 3/135)
DMEK (N=135)	NR	n (%) ≥20/25 Snellen=80 (73%)	NR		
Price et al. (2009) (29)	N=60	N=57 at 3-mos	N=60	N=57 at 3-mos	NR
DMEK	Median preoperative BSCVA=20/50	n (%) ≥20/25 Snellen=36 (63%)	3010 +/- 200 (range, 2520-3430)	30% +/- 20% (range, 2.7%-78%)	NR
	BSCVA	BSCVA FU [time]	ECD pre-operative (mean +/- SD, cells/mm²)	ECD 6 mos FU (mean +/- SD, cells/mm²)	Donor Tissue Loss (N=corneas)
Guerra et al. (2011) (30)	N=108				

DMEK	0.51 +/- 0.44 logMar of the minimum angle of resolution units (20/65; range, 20/20-20/2000)	1-year: 0.07 1 +/- 0.09 logMar of the minimum angle of resolution units (20/24; range, 20/15 - 20/40); p<0.001	2980+/-252 (2514-3706) at 1 yr	1911+/-593 (range, 347-2976) at 1 yr	N=6 (4.2%)
McCauley et al. (2011) (32)					
DMAEK (N=40)	Median pre-op BSCVA was 20/40 (range: 20/25-20/400)	6-mos: median BSCVA was 20/25 (range: 20/15-20/40); 48% 20/20; 74% 20/25; 93% 20/30; all \geq 20/40	The median donor ECD=3140 cells/mm ² (range: 2695-4630 cells/mm ²)	6 mos FU, median ECD was 2121 cells/mm ² (range: 1204-4268 cells/mm ² , n=30)	Not statistically significant
	Probability of Rejection % at 1 yr	Probability of Rejection % at 2 yrs	Eyes still followed without rejection (n) at 1 yr	Eyes still followed without rejection (n) at 2 yrs	
Anshu et al. (2012) (31)					
DMEK (n=141)	1	1	82	35	
DSEK (n=598)	8	12	246	79	
PK (n=30)	14	18	23	11	

BCVA: best-corrected visual acuity; BSCVA: best spectacle-corrected visual acuity; CI: confidence interval; DMEK: Descemet membrane endothelial keratoplasty; DSAEK: Descemet stripping automated endothelial keratoplasty; DSEK: Descemet stripping endothelial keratoplasty; ECD: endothelial corneal dystrophy; EDM: endothelium-Descemet's membrane; FU: follow up; mo(s): month(s); N: number; NR: not reported; OS: overall survival; PK: penetrating keratoplasty; SD: standard deviation; yr(s): years; logMar: Logarithm of the Minimum Angle of Resolution.

Section Summary: Descemet Membrane Endothelial Keratoplasty and Descemet Membrane Automated Endothelial Keratoplasty

Evidence for the use of Descemet membrane endothelial keratoplasty or Descemet membrane automated endothelial keratoplasty consists of several systematic reviews with overlapping studies, and several observational studies, some of which had no comparators and some of which compared Descemet membrane endothelial keratoplasty or Descemet membrane automated endothelial keratoplasty with Descemet stripping endothelial keratoplasty or Descemet stripping automated endothelial keratoplasty. Analyses in the individual studies and the meta-analyses consistently showed that patients receiving Descemet membrane endothelial keratoplasty or Descemet membrane automated endothelial keratoplasty experienced significantly better visual acuity outcomes postprocedure than patients receiving Descemet stripping endothelial keratoplasty or Descemet stripping automated endothelial keratoplasty, both short-term and through 1 year of follow-up. A large cohort study showed that intraoperative complications decreased as surgeon experience increased. Some studies reported similar complication rates between the procedures, some reported more complications with Descemet membrane endothelial keratoplasty than Descemet stripping endothelial keratoplasty, though the complications were not considered severe.

Femtosecond Laser-Assisted Endothelial Keratoplasty and Femtosecond and Excimer Laser-Assisted Endothelial Keratoplasty

Clinical Context and Therapy Purpose

The purpose of femtosecond laser-assisted endothelial keratoplasty and femtosecond and excimer laser-assisted endothelial keratoplasty is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as penetrating keratoplasty, in individuals with endothelial disease of the cornea.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with endothelial disease of the cornea. Diseases that affect the endothelial layer include Fuchs endothelial dystrophy, aphakic and pseudophakic bullous keratopathy (corneal edema following cataract extraction), and failure or rejection of a previous corneal transplant.

Interventions

The therapy being considered is femtosecond laser-assisted endothelial keratoplasty and femtosecond and excimer laser-assisted endothelial keratoplasty. Variations of femtosecond laser-assisted endothelial keratoplasty include femtosecond laser-assisted Descemet stripping automated endothelial keratoplasty.

Comparators

Comparators of interest include penetrating keratoplasty.

Outcomes

The general outcomes of interest are change in disease status, morbid events, and functional outcomes. Relevant outcome measures include visual acuity, endothelial cell densities, patient satisfaction or quality-of-life, and complications including graft rejection, graft dislocation, and need for rebubble procedures. Follow-up generally occurs through 1 to 2 years post-surgery.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Systemic Review

Liu et al. (2021) conducted a systematic review and meta-analysis of studies comparing femtosecond laser-enabled keratoplasty with conventional penetrating keratoplasty. (33) The literature search was conducted through April 2018 and identified 7 comparative studies for inclusion. Follow-up periods of the included studies spanned from 6 months to 3.5 years, with the majority of patients having up to 1 year of follow-up. The meta-analyses of 1855 eyes illustrated that mean BSCVA after femtosecond laser-enabled keratoplasty was significantly better than after penetrating keratoplasty ($p=.00$; SMD, -0.23 ; 95% CI, -0.37 to -0.10). Endothelial cell density was also significantly better preserved in the femtosecond laser-enabled keratoplasty group ($p=.03$, SMD: 0.63 ; 95% CI, 0.07 to 1.20). Results were comparable amongst both groups in spherical equivalent, graft rejection, graft failure, and complication.

Randomized Controlled Trials

Ivarsen et al. (2018) conducted an RCT of ultrathin Descemet stripping automated endothelial keratoplasty or femtosecond-prepared Descemet stripping automated endothelial keratoplasty using the Ziemer LDV Z8 femtosecond laser. (34) Outcome measures were planned after 1, 3, 6, 12 and 24 months with visual acuity, refraction, Scheimpflug tomography, whole eye scatter measurement, and anterior optical coherence tomography. However, graft dislocation occurred in all patients randomized to femtosecond-prepared Descemet stripping automated endothelial keratoplasty which was managed with rebubbling. No patients with ultrathin Descemet stripping automated endothelial keratoplasty experienced graft dislocation. Additionally, all patients treated with femtosecond-prepared Descemet stripping automated endothelial keratoplasty had significantly poorer clinical outcomes compared with ultrathin Descemet stripping automated endothelial keratoplasty patients. After 3 months, visual acuity was scored as approximately 2.5 times worse. The optical scatter index was also significantly greater in patients receiving femtosecond-prepared Descemet stripping automated endothelial keratoplasty compared to ultrathin Descemet stripping automated endothelial keratoplasty at 3 months (mean, 12; standard deviation [SD], 3; range, 8 to 16 vs. mean, 5; SD, 3; range, 2 to 9). While the planned enrollment was set at 80, after 1 month only 6 patients were treated with

femtosecond-prepared Descemet stripping automated endothelial keratoplasty and 5 patients received ultrathin Descemet stripping automated endothelial keratoplasty. Due to the large differences in observed clinical outcomes, no further patients were recruited, and the study was suspended.

Cheng et al. (2009) conducted a multicenter randomized trial in Europe that compared femtosecond laser-assisted endothelial keratoplasty with penetrating keratoplasty. (35) Eighty patients with Fuchs endothelial dystrophy, bullous keratopathy, or posterior polymorphous dystrophy, and a BSCVA less than 20/50 were included in the trial. In the femtosecond laser-assisted endothelial keratoplasty group, 4 of the 40 eyes did not receive treatment due to significant preoperative events and were excluded from the analysis. Eight (22%) of 36 eyes failed, and 2 patients were lost to follow-up due to death in the femtosecond laser-assisted endothelial keratoplasty group. One patient was lost to follow-up in the penetrating keratoplasty group due to health issues. At 12 months postoperatively, refractive astigmatism was lower in the femtosecond laser-assisted endothelial keratoplasty group (86%) than in the penetrating keratoplasty group (51%, with astigmatism of ≥ 3 D); however, there was a greater hyperopic shift in the femtosecond laser-assisted endothelial keratoplasty group than in the penetrating keratoplasty group. Mean BSCVA was better following penetrating keratoplasty than femtosecond laser-assisted endothelial keratoplasty at the 3-, 6-, and 12-month follow-ups. There was greater endothelial cell loss in the femtosecond laser-assisted endothelial keratoplasty group (65%) than in the penetrating keratoplasty group (23%). With the exception of dislocation and need to reposition the femtosecond laser-assisted endothelial keratoplasty grafts in 28% of eyes, the percentage of complications was similar between groups. Complications in the femtosecond laser-assisted endothelial keratoplasty group were due to pupillary block, graft failure, epithelial ingrowth, and elevated intraocular pressure, whereas complications in the penetrating keratoplasty group were related to the sutures and elevated intraocular pressure.

Nonrandomized Studies

Sorkin et al. (2019) reported 3-year outcomes of a retrospective, interventional study comparing femtosecond laser-assisted Descemet membrane endothelial keratoplasty with manual Descemet membrane endothelial keratoplasty in patients with Fuchs endothelial corneal dystrophy. (36) Sixteen eyes of 15 patients were evaluated in the femtosecond-prepared Descemet membrane endothelial keratoplasty group for an average follow-up up 33.0 \pm 9.0 months and 45 eyes of 40 patients were evaluated in the manual Descemet membrane endothelial keratoplasty group for an average follow-up of 32.0 \pm 7.0 months. Best spectacle-corrected visual acuity was not statistically different at 1-, 2-, and 3-years post-surgery ($p=.849$, $p=.465$, and $p=.936$, respectively). Rates of significant graft detachment were significantly higher in the manual Descemet membrane endothelial keratoplasty group than in the femtosecond prepared Descemet membrane endothelial keratoplasty group (35.6% vs. 6.25%; $p=.027$). Rebubbling rates were also significantly higher in the manual Descemet membrane endothelial keratoplasty group (33.3% vs. 6.25%; $p=.047$). Endothelial cell loss rates were significantly lower in the femtosecond prepared Descemet membrane endothelial keratoplasty group at 1 year (26.8% vs. 36.5%; $p=.042$) and 2 years (30.5% vs. 42.3%; $p=.008$), however, this

trend was lost at 3 years (37% vs. 47.5%; $p=.057$). (37) The primary graft failure rate was 0% in femtosecond prepared Descemet membrane endothelial keratoplasty compared to 8.9% in manual Descemet membrane endothelial keratoplasty ($p=.565$). While study authors speculate that the higher detachment and rebubbling rate in manual Descemet membrane endothelial keratoplasty may be related to retained Descemet tags and islands, this study is limited by its retrospective nature and nonrandomized design and cannot account for potential baseline differences in patient anatomy. Hosny et al. (2017) reported on results from a case series on 20 eyes (19 patients) that underwent a femtosecond prepared Descemet stripping automated endothelial keratoplasty. (38) After 3 months of follow-up, patients experienced significant improvements in corneal thickness, measured by anterior segment optical coherence tomography. Visual acuity significantly improved each month of the 3-month follow-up, with the largest improvement seen in the first month postprocedure. Complications specific to the femtosecond laser-assisted procedure were thickness disparities causing protrusion of the posterior disc ($n=6$) and air trapping in the interface ($n=2$). The former complication was corrected by modifying procedure parameters, and the latter was corrected by venting of the air bubble.

In a small retrospective cohort study, Vetter et al. (2013) found a reduction in visual acuity when the endothelial transplant was prepared with a laser (femtosecond laser-assisted endothelial keratoplasty=0.48 logMAR; $n=8$) compared with a microtome (Descemet stripping automated endothelial keratoplasty=0.33 logMAR; $n=14$). (39) There was also greater surface irregularity with femtosecond laser-assisted endothelial keratoplasty.

Section Summary: Femtosecond Laser-Assisted Endothelial Keratoplasty and Femtosecond and Excimer Laser-Assisted Endothelial Keratoplasty

Evidence for femtosecond laser-assisted endothelial keratoplasty consists of 3 small observational studies, 2 RCTs, and 1 systematic review. The systematic review reported that femtosecond laser-assisted endothelial keratoplasty may have advantages to achieving better outcomes in BSCVA and endothelial cell density preservation. One observational study showed improvements following the procedure, though there was no comparison group and the other showed worse outcomes with the laser compared with Descemet stripping automated endothelial keratoplasty. One RCT indicated that patients undergoing penetrating keratoplasty experienced better outcomes than patients in the femtosecond laser-assisted endothelial keratoplasty group after 1 year of follow-up. Complication rates were similar between groups. Another RCT reported better clinical outcomes and no instances of graft dislocation with microkeratome-prepared Descemet stripping automated endothelial keratoplasty compared to femtosecond prepared Descemet stripping automated endothelial keratoplasty.

Summary of Evidence

For individuals who have endothelial disease of the cornea who receive Descemet stripping endothelial keratoplasty or Descemet stripping automated endothelial keratoplasty, the evidence includes a number of cohort studies, randomized controlled trials (RCTs), and systematic reviews. Relevant outcomes are change in disease status, morbid events, and functional outcomes. The available literature has indicated that these procedures improve

visual outcomes and reduce serious complications associated with penetrating keratoplasty. Specifically, visual recovery occurs much earlier. Because endothelial keratoplasty maintains an intact globe without a sutured donor cornea, astigmatism or the risk of severe, sight-threatening complications such as expulsive suprachoroidal hemorrhage and postoperative catastrophic wound failure, are eliminated. The Descemet Endothelial Thickness Comparison Trial (DETECT) RCT reported improved visual acuity outcomes with Descemet membrane endothelial keratoplasty compared to ultra-thin Descemet stripping automated endothelial keratoplasty. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have endothelial disease of the cornea who receive Descemet membrane endothelial keratoplasty or Descemet membrane automated endothelial keratoplasty, the evidence includes a number of cohort studies and systematic reviews. Relevant outcomes are change in disease status, morbid events, and functional outcomes. Evidence from the cohort studies and meta-analyses has consistently shown that the use of Descemet membrane endothelial keratoplasty and Descemet membrane automated endothelial keratoplasty procedures improve visual acuity. When compared with Descemet stripping endothelial keratoplasty and Descemet stripping automated endothelial keratoplasty, Descemet membrane endothelial keratoplasty and Descemet membrane automated endothelial keratoplasty showed significantly greater improvements in visual acuity, both in the short term and through 1 year of follow-up. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have endothelial disease of the cornea who receive femtosecond laser-assisted endothelial keratoplasty and femtosecond and excimer laser-assisted endothelial keratoplasty, the evidence includes a multicenter RCT and a systematic review comparing femtosecond laser-assisted endothelial keratoplasty with penetrating keratoplasty, and an RCT comparing femtosecond-prepared Descemet stripping automated endothelial keratoplasty to microkeratome-prepared Descemet membrane automated endothelial keratoplasty. Relevant outcomes are change in disease status, morbid events, and functional outcomes. There were conflicting results in the evidence regarding mean best-corrected visual acuity and endothelial cell loss after femtosecond laser-assisted endothelial keratoplasty versus penetrating keratoplasty. Mean best-corrected visual acuity was worse after femtosecond laser-assisted endothelial keratoplasty than after penetrating keratoplasty, and endothelial cell loss was higher with femtosecond laser-assisted endothelial keratoplasty. With the exception of dislocation and need for repositioning of the femtosecond laser-assisted endothelial keratoplasty, the percentage of complications was similar between groups. Complications in the femtosecond laser-assisted endothelial keratoplasty group were due to pupillary block, graft failure, epithelial ingrowth, and elevated intraocular pressure, whereas complications in the penetrating keratoplasty group were related to sutures and elevated intraocular pressure. Worsened visual acuity and a 100% graft dislocation rate were reported for femtosecond-prepared Descemet stripping automated endothelial keratoplasty compared to 0% in manually prepared Descemet stripping automated endothelial keratoplasty. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Practice Guidelines and Position Statements

American Academy of Ophthalmology

In 2009, the American Academy of Ophthalmology (AAO) published a position paper on endothelial keratoplasty, stating that the optical advantages, speed of visual rehabilitation, and lower risk of catastrophic wound failure have driven the adoption of endothelial keratoplasty as the standard of care for patients with endothelial failure and otherwise healthy corneas. The 2009 AAO position paper was based in large part on an AAO comprehensive review of the literature on Descemet stripping automated endothelial keratoplasty. (3) The AAO concluded that "the evidence reviewed suggests Descemet stripping automated endothelial keratoplasty appears safe and efficacious for the treatment of endothelial diseases of the cornea. Evidence from retrospective and prospective Descemet stripping automated endothelial keratoplasty reports described a variety of complications from the procedure, but these complications do not appear to be permanently sight-threatening or detrimental to the ultimate vision recovery in the majority of cases. Long-term data on endothelial cell survival and the risk of late endothelial rejection cannot be determined with this review... Descemet stripping automated endothelial keratoplasty should not be used in lieu of penetrating keratoplasty for conditions with concurrent endothelial disease and anterior corneal disease. These situations would include concurrent anterior corneal dystrophies, anterior corneal scars from trauma or prior infection, and ectasia after previous laser vision correction surgery."

In 2018, the AAO published a Preferred Practice Pattern on corneal edema and opacification. (40) The guidance was updated in 2024. (41) Based on their findings, the following statement and recommendation was made by AAO in 2024: "Endothelial keratoplasty has supplanted penetrating keratoplasty as the procedure of choice in cases of endothelial failure in the absence of corneal scarring because patients achieve more rapid visual rehabilitation and reduced risk of immune-mediated rejection of the transplanted tissue and less induced astigmatism."

National Institute for Health and Care Excellence (NICE)

In 2009, NICE released guidance on corneal endothelial transplantation. (42) Additional data reviewed from the United Kingdom Transplant Register showed lower graft survival rates after endothelial keratoplasty than after penetrating keratoplasty; however, the difference in graft survival between the 2 procedures was noted to be narrowing with increased experience in endothelial keratoplasty use. The guidance concluded that "current evidence on the safety and efficacy of corneal endothelial transplantation (also known as endothelial keratoplasty) is adequate to support the use of this procedure." The guidance noted that techniques for this procedure continue to evolve, and thorough data collection should continue to allow future review of outcomes.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and/or unpublished trials that might influence this policy are listed in Table 7.

Table 7. Summary of Key Trials

NCT Number	Trial Name	Planned Enrollment	Completion Date
<i>Unpublished</i>			
NCT00543660	Descemet Stripping (Automated) Endothelial Keratoplasty (DSEK or DSAEK) (DSAEK)	20	Mar 2018
NCT00521898	Prospective Clinical Study on Descemet Membrane Endothelial Keratoplasty	1000	Feb 2020
NCT02793310	Corneal Transplantation by DMEK - is it Really Better Than DSAEK?	54	Feb 2019
NCT00800111	Open-enrollment, Prospective Study of Endothelial Keratoplasty Outcomes	2593	Feb 2018
NCT02470793	Technique and Results In Endothelial Keratoplasty (TREK)	62	Jan 2021
NCT03619434	Pilot Study of Femtolaser Assisted Keratoplasty Versus Conventional Keratoplasty	30	Dec 2021 (unknown)

NCT: national clinical trial; DMEK: Descemet membrane endothelial keratoplasty; DSAEK: Descemet stripping automated endothelial keratoplasty; DSEK: Descemet stripping endothelial keratoplasty.

Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member's benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

CPT Codes	65756, 65757
HCPCS Codes	None

*Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.

References

1. Eye Bank Association of America. 2019 Eye Banking Statistical Report (2019). Available at <<https://restoresight.org>> (accessed April 15, 2025).

2. Woo JH, Ang M, Htoo HM, et al. Descemet Membrane Endothelial Keratoplasty Versus Descemet Stripping Automated Endothelial Keratoplasty and Penetrating Keratoplasty. *Am J Ophthalmol.* Nov 2019; 207:288-303. PMID 31228467
3. Lee WB, Jacobs DS, Musch DC, et al. Descemet's stripping endothelial keratoplasty: safety and outcomes: a report by the American Academy of Ophthalmology. *Ophthalmology.* Sep 2009; 116(9):1818-1830. PMID 19643492
4. Stuart AJ, Romano V, Virgili G, et al. Descemet's membrane endothelial keratoplasty (DMEK) versus Descemet's stripping automated endothelial keratoplasty (DSAEK) for corneal endothelial failure. *Cochrane Database Syst Rev.* Jun 25 2018; 6(6):CD012097. PMID 29940078
5. Marques RE, Guerra PS, Sousa DC, et al. DMEK versus DSAEK for Fuchs' endothelial dystrophy: A meta-analysis. *Eur J Ophthalmol.* Jan 2019; 29(1):15-22. PMID 29661044
6. Chamberlain W, Lin CC, Austin A, et al. Descemet Endothelial Thickness Comparison Trial: A Randomized Trial Comparing Ultrathin Descemet Stripping Automated Endothelial Keratoplasty with Descemet Membrane Endothelial Keratoplasty. *Ophthalmology.* Jan 2019; 126(1):19-26. PMID 29945801
7. Duggan MJ, Rose-Nussbaumer J, Lin CC, et al. Corneal Higher-Order Aberrations in Descemet Membrane Endothelial Keratoplasty Versus Ultrathin DSAEK in the Descemet Endothelial Thickness Comparison Trial: A Randomized Clinical Trial. *Ophthalmology.* Jul 2019; 126(7):946-957. PMID 30776384
8. Lin CC, Chamberlain WD, Kakigi C, et al. Mediators of Visual Acuity in Descemet Membrane Endothelial Keratoplasty and Ultrathin Descemet Stripping Automated Endothelial Keratoplasty. *Cornea.* Jul 1 2024; 43(7):853-859. PMID 37986182
9. Hirabayashi KE, Chamberlain W, Rose-Nussbaumer J, et al. Corneal Light Scatter After Ultrathin Descemet Stripping Automated Endothelial Keratoplasty Versus Descemet Membrane Endothelial Keratoplasty in Descemet Endothelial Thickness Comparison Trial: A Randomized Controlled Trial. *Cornea.* Jun 2020; 39(6):691-696. PMID 31939923
10. Dunker SL, Dickman MM, Wisse RPL, et al. Descemet Membrane Endothelial Keratoplasty versus Ultrathin Descemet Stripping Automated Endothelial Keratoplasty: A Multicenter Randomized Controlled Clinical Trial. *Ophthalmology.* Sep 2020; 127(9):1152-1159. PMID 32386811
11. Wilhelm TI, Gauch L, Bhringer D, et al. Ten-year outcomes after DMEK, DSAEK, and PK: insights on graft survival, endothelial cell density loss, rejection and visual acuity. *Sci Rep.* Jan 8 2025; 15(1):1249. PMID 39774993
12. Fuest M, Ang M, Htoo HM, et al. Long-Term Visual Outcomes Comparing Descemet Stripping Automated Endothelial Keratoplasty And Penetrating Keratoplasty. *Am J Ophthalmol.* Oct 2017; 182:62-71. PMID 28739420
13. Heinzelmann S, Böhringer D, Eberwein P, et al. Outcomes of Descemet membrane endothelial keratoplasty, Descemet stripping automated endothelial keratoplasty and penetrating keratoplasty from a single centre study. *Graefes Arch Clin Exp Ophthalmol.* Mar 2016; 254(3):515-522. PMID 26743748
14. Wacker K, Baratz KH, Maguire LJ, et al. Descemet Stripping Endothelial Keratoplasty for Fuchs' Endothelial Corneal Dystrophy: Five-Year Results of a Prospective Study. *Ophthalmology.* Jan 2016; 123(1):154-160. PMID 26481820

15. Li JY, Terry MA, Goshe J, et al. Three-year visual acuity outcomes after Descemet's stripping automated endothelial keratoplasty. *Ophthalmology*. Jun 2012; 119(6):1126-1129. PMID 22364863
16. Dapena I, Ham L, Melles GR. Endothelial keratoplasty: DSEK/DSAEK or DMEK--the thinner the better? *Curr Opin Ophthalmol*. Jul 2009; 20(4):299-307. PMID 19417653
17. Rose L, Kelliher C, Jun AS. Endothelial keratoplasty: historical perspectives, current techniques, future directions. *Can J Ophthalmol*. Aug 2009; 44(4):401-405. PMID 19606160
18. Deng SX, Lee WB, Hammersmith KM, et al. Descemet Membrane Endothelial Keratoplasty: Safety And Outcomes: A Report by the American Academy of Ophthalmology. *Ophthalmology*. Feb 2018; 125(2):295-310. PMID 28923499
19. Singh A, Zarei-Ghanavati M, Avadhanam V, et al. Systematic Review And Meta-Analysis of Clinical Outcomes of Descemet Membrane Endothelial Keratoplasty Versus Descemet Stripping Endothelial Keratoplasty/Descemet Stripping Automated Endothelial Keratoplasty. *Cornea*. Nov 2017; 36(11):1437-1443. PMID 28834814
20. Pavlovic I, Shajari M, Herrmann E, et al. Meta-Analysis of Postoperative Outcome Parameters Comparing Descemet Membrane Endothelial Keratoplasty Versus Descemet Stripping Automated Endothelial Keratoplasty. *Cornea*. Dec 2017; 36(12):1445-1451. PMID 28957976
21. Li S, Liu L, Wang W, et al. Efficacy and safety of Descemet's membrane endothelial keratoplasty versus Descemet's stripping endothelial keratoplasty: A systematic review and meta-analysis. *PLoS One*. Dec 18 2017; 12(12):e0182275. PMID 29252983
22. Wu J, Wu T, Li J, et al. DSAEK or DMEK for failed penetrating keratoplasty: a systematic review and single-arm meta-analysis. *Int Ophthalmol*. Jul 2021; 41(7):2315-2328. PMID 34117964
23. Maier AB, Milek J, Joussen AM, et al. Systematic Review and Meta-analysis: Outcomes After Descemet Membrane Endothelial Keratoplasty Versus Ultrathin Descemet Stripping Automated Endothelial Keratoplasty. *Am J Ophthalmol*. Jan 2023; 245:222-232. PMID 36220351
24. Oellerich S, Baydoun L, Peraza-Nieves J, et al. Multicenter Study of 6-Month Clinical Outcomes After Descemet Membrane Endothelial Keratoplasty. *Cornea*. Dec 2017; 36(12):1467-1476. PMID 28957979
25. Tourtas T, Laaser K, Bachmann BO, et al. Descemet membrane endothelial keratoplasty versus descemet stripping automated endothelial keratoplasty. *Am J Ophthalmol*. Jun 2012; 153(6):1082-1090.e2. PMID 22397955
26. van Dijk K, Ham L, Tse WH, et al. Near complete visual recovery and refractive stability in modern corneal transplantation: Descemet membrane endothelial keratoplasty (DMEK). *Cont Lens Anterior Eye*. Feb 2013; 36(1):13-21. PMID 23108011
27. Ham L, Dapena I, van Luijk C, et al. Descemet membrane endothelial keratoplasty (DMEK) for Fuchs endothelial dystrophy: review of the first 50 consecutive cases. *Eye (Lond)*. Oct 2009; 23(10):1990-1998. PMID 19182768
28. Dapena I, Ham L, Droutras K, et al. Learning Curve in Descemet's Membrane Endothelial Keratoplasty: First Series of 135 Consecutive Cases. *Ophthalmology*. Nov 2011; 118(11):2147-2154. PMID 21777980

29. Price MO, Giebel AW, Fairchild KM, et al. Descemet's membrane endothelial keratoplasty: prospective multicenter study of visual and refractive outcomes and endothelial survival. *Ophthalmology*. Dec 2009; 116(12):2361-2368. PMID 19875170
30. Guerra FP, Anshu A, Price MO, et al. Descemet's membrane endothelial keratoplasty: prospective study of 1- year visual outcomes, graft survival, and endothelial cell loss. *Ophthalmology*. Dec 2011; 118(12):2368-2373. PMID 21872938
31. Anshu A, Price MO, Price FW, Jr. Risk of corneal transplant rejection significantly reduced with Descemet's membrane endothelial keratoplasty. *Ophthalmology*. Mar 2012; 119(3):536-540. PMID 22218143
32. McCauley MB, Price MO, Fairchild KM, et al. Prospective study of visual outcomes and endothelial survival with Descemet membrane automated endothelial keratoplasty. *Cornea*. Mar 2011; 30(3):315-319. PMID 21099412
33. Liu Y, Li X, Li W, et al. Systematic review and meta-analysis of femtosecond laser-enabled keratoplasty versus conventional penetrating keratoplasty. *Eur J Ophthalmol*. May 2021; 31(3):976-987. PMID 32223431
34. Ivarsen A, Hjortdal J. Clinical outcome of Descemet's stripping endothelial keratoplasty with femtosecond laser-prepared grafts. *Acta Ophthalmol*. Aug 2018; 96(5):e655-e656. PMID 29372934
35. Cheng YY, Schouten JS, Tahzib NG, et al. Efficacy and safety of femtosecond laser-assisted corneal endothelial keratoplasty: a randomized multicenter clinical trial. *Transplantation*. Dec 15 2009; 88(11):1294-1302. PMID 19996929
36. Sorkin N, Mednick Z, Einan-Lifshitz A, et al. Three-Year Outcome Comparison Between Femtosecond Laser-Assisted and Manual Descemet Membrane Endothelial Keratoplasty. *Cornea*. Jul 2019; 38(7):812-816. PMID 30973405
37. Singhal D, Maharana PK. RE: "Three-Year Outcome Comparison Between Femtosecond Laser-Assisted And Manual Descemet Membrane Endothelial Keratoplasty." *Cornea*. Nov 2019; 38(11):e51. PMID 31414998
38. Hosny MH, Marrie A, Sidky MK, et al. Results of Femtosecond Laser-Assisted Descemet Stripping Automated Endothelial Keratoplasty. *J Ophthalmol*. Jun 11 2017; 2017:8984367. PMID 28695004
39. Vetter JM, Butsch C, Faust M, et al. Irregularity of the posterior corneal surface after curved interface femtosecond laser-assisted versus microkeratome-assisted Descemet stripping automated endothelial keratoplasty. *Cornea*. Feb 2013; 32(2):118-124. PMID 23132446
40. Farid M, Rhee MK, Akpek EK, et al. Corneal Edema and Opacification Preferred Practice Pattern®. *Ophthalmology*. Jan 2019; 126(1):P216-P285. PMID 30366795
41. Mian SI, Viriya ET, Ahmad S, et al. Corneal Edema and Opacification Preferred Practice Pattern®. *Ophthalmology*. Apr 2024; 131(4):P247-P305. PMID 38349302
42. National Institute for Health and Care Excellence (NICE). Corneal endothelial transplantation [IPG304] (2009). Available at <<https://www.nice.org.uk>> (accessed January 14, 2025).

Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does not have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been developed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

Policy History/Revision	
Date	Description of Change
07/15/2025	Document updated with literature review. Coverage unchanged. References 8, 11, and 41 added.
12/15/2024	Document updated with literature review. Coverage unchanged. Added reference 38.
12/01/2023	Document updated with literature review. Coverage unchanged. References 20, 21, & 31 added, others updated or removed.
07/15/2022	Reviewed. No changes.
07/15/2021	Document updated with literature review. Coverage unchanged. References 2, 3, 5-9, 29, 31, & 32 added, others updated or removed.
06/15/2020	Reviewed. No changes.
07/01/2019	Document updated with literature review. Coverage unchanged. Added references: 1, 3, 9-13, 22, & 25.
06/15/2018	Reviewed. No changes.
07/15/2017	Document updated with literature review. Coverage unchanged.
08/01/2016	Reviewed. No changes.
01/01/2015	New Medical Document. Document updated with literature review. Medical document SUR713.001 was divided into (3) policies. The following changes were made to the coverage position: Added to coverage position to include: Endothelial keratoplasty (Descemet's stripping endothelial keratoplasty [DSEK], Descemet's stripping automated endothelial keratoplasty [DSAEK], Descemet's membrane endothelial keratoplasty [DMEK], or Descemet's membrane automated endothelial keratoplasty [DMAEK]) may be considered medically necessary for the treatment of endothelial dysfunction, including but not limited to 1) Ruptures in Descemet's membrane, Endothelial dystrophy 2.) Iridocorneal endothelial (ICE) syndrome 3.) Corneal edema attributed to endothelial failure. 4.) Femtosecond laser-assisted corneal endothelial keratoplasty (FLEK) or femtosecond and excimer lasers-assisted endothelial keratoplasty (FELEK) are considered experimental, investigational and/or unproven. 5.) Endothelial keratoplasty is considered not medically necessary when endothelial dysfunction is not the primary cause of decreased corneal clarity. This topic was previously addressed on

	SUR713.001, Refractive and Therapeutic Keratoplasty. CPT/HCPCS code(s) updated.
--	---