

Policy Number	RX501.148
Policy Effective Date	01/01/2025

Betibeglogene autotemcel

Table of Contents
Coverage
Policy Guidelines
Description
Rationale
Coding
References
Policy History

Related Policies (if applicable)
None

Disclaimer

Medical policies are a set of written guidelines that support current standards of practice. They are based on current peer-reviewed scientific literature. A requested therapy must be proven effective for the relevant diagnosis or procedure. For drug therapy, the proposed dose, frequency and duration of therapy must be consistent with recommendations in at least one authoritative source. This medical policy is supported by FDA-approved labeling and/or nationally recognized authoritative references to major drug compendia, peer reviewed scientific literature and acceptable standards of medical practice. These references include, but are not limited to: MCG care guidelines, DrugDex (IIa level of evidence or higher), NCCN Guidelines (IIb level of evidence or higher), NCCN Compendia (IIb level of evidence or higher), professional society guidelines, and CMS coverage policy.

Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

Legislative Mandates

EXCEPTION: For HCSC members residing in the state of Ohio, § 3923.60 requires any group or individual policy (Small, Mid-Market, Large Groups, Municipalities/Counties/Schools, State Employees, Fully-Insured, PPO, HMO, POS, EPO) that covers prescription drugs to provide for the coverage of any drug approved by the U. S. Food and Drug Administration (FDA) when it is prescribed for a use recognized as safe and effective for the treatment of a given indication in one or more of the standard medical reference compendia adopted by the United States Department of Health and Human Services or in medical literature even if the FDA has not approved the drug for that indication. Medical literature support is only satisfied when safety and efficacy has been confirmed in two articles from major peer-reviewed professional medical journals that present data supporting the proposed off-label use or uses as generally safe and effective. Examples of accepted journals include, but are not limited to, Journal of American Medical Association (JAMA), New England Journal of Medicine (NEJM), and Lancet. Accepted

study designs may include, but are not limited to, randomized, double blind, placebo controlled clinical trials. Evidence limited to case studies or case series is not sufficient to meet the standard of this criterion. Coverage is never required where the FDA has recognized a use to be contraindicated and coverage is not required for non-formulary drugs.

Coverage

Betibeglogene autotemcel (Zynteglo[®]) **may be considered medically necessary** if **ALL** the following criteria are met:

1. Adult and pediatric patients ≤50 years of age with transfusion dependent beta-thalassemia; AND
2. Documented history of transfusions evidenced by:
 - a. 100 mL/kg/per year or more of packed red blood cells (pRBCs); OR
 - b. 8 or more transfusions of packed red blood cells (pRBCs) per year in the preceding 1 year; AND
3. Negative serologic test for HIV infection; AND
4. Clinically stable and eligible to undergo hematopoietic stem cell (HSC) mobilization; AND
5. Individual does NOT have ANY of the following:
 - a. White blood cell count less than 3×10^9 /Liter and/or platelet count less than 100×10^9 not related to hypersplenism;
 - b. History of prior gene therapy or allogenic hematopoietic stem cell transplant;
 - c. Severe iron overload;
 - d. Advanced liver disease;
 - e. Prior or current malignancy, myeloproliferative disorder, or significant immunodeficiency.

Repeat treatment of Betibeglogene autotemcel (Zynteglo[®]) **is considered experimental, investigational, and/or unproven.**

Betibeglogene autotemcel (Zynteglo[®]) **is considered experimental, investigational, and/or unproven** for all other indications.

Policy Guidelines

None.

Description

Beta-thalassemia (β-thalassemia)

β-thalassemia is an inherited blood disorder that occurs as a result of a genetic variant in the *HBB* gene that codes for the production of β-globin chains. As a result, there is reduced synthesis or absence of β-globin chains leading to impaired production of hemoglobin. The clinical presentation is that of anemia which requires iron supplementation and multiple

downstream sequelae from the disease. These sequelae include growth retardation, skeletal changes (particularly in the face and long bones of the legs), osteoporosis, leg ulcers, and development of extramedullary masses. High output heart failure from anemia is also common without treatment. Without transfusion therapy, such patients die within the first few years of life, primarily from heart failure or infection. (1)

Life expectancy of individuals with transfusion-dependent β -thalassemia is much lower than population norms. From 2011 to 2021 the median age of death for a person in the US with transfusion-dependent β -thalassemia was 37. (2) Additionally, individuals with transfusion-dependent β -thalassemia report decreased quality of life due to the impact on physical and mental health. (3, 4)

All humans have 2 copies of the *HBB* gene and each copy produces the β -globin protein. Different types of β -thalassemia categorized by genotype are summarized in Table 1. When only 1 *HBB* gene is affected, the phenotype is less severe and individuals are generally asymptomatic due to compensation from the other normal gene. These individuals are called β -thalassemia minor or carrier. However, if both copies of *HBB* gene are affected there is a quantitative reduction or absence of β -globin protein. Phenotypes that manifest as a reduction in β -globin chains are referred to as “ β -thalassemia intermedia” and phenotypes that manifest as absence in β -globin chains are called “ β -thalassemia major.” (5)

More recently, patients have been classified according to their transfusion status (i.e., transfusion-dependent β -thalassemia or non-transfusion-dependent β -thalassemia). For this medical policy, focus is on transfusion-dependent β -thalassemia patients which generally includes “ β -thalassemia major” but occasionally may include patients with “ β -thalassemia intermedia”. Clinical studies reviewed define “transfusion dependence” as history of at least 100 mL/kg/year of peripheral red blood cells or ≥ 8 transfusions of peripheral red blood cells per year for the prior 2 years. “Transfusion independence” was defined as a weighted average hemoglobin (Hb) of at least 9 g/dL without any transfusions for a continuous period of at least 12 months at any time during the study after infusion of betibeglogene autotemcel.

Table 1. Different Types of β -Thalassemia (5-7)

Type	Genotype	Description
β -thalassemia major (generally transfusion dependent)	β^0/β^0 or β^0/β^+	<ul style="list-style-type: none"> • Presents within the first 2 years of life with severe microcytic anemia (typical hemoglobin 3 to 4 g/dL), mild jaundice, and hepatosplenomegaly. • Requires regular red blood cell transfusions and other medical treatments.
Thalassemia intermedia	β^+/β^+	<ul style="list-style-type: none"> • Presents at a later age with similar, but milder clinical signs and symptoms of thalassemia • Moderately severe anemia; some may need regular blood transfusions.

Thalassemia minor	β/β^0 or β/β^+	<ul style="list-style-type: none"> • Also called “β-thalassemia carrier” or “β-thalassemia trait.” • Usually clinically asymptomatic but may have a mild anemia. • Generally do not require any treatment.
-------------------	------------------------------------	---

β^0 refers to no beta globin production; β^+ refers to decreased beta globin production.

Epidemiology

β -thalassemia is 1 of the most common monogenic disorders, but its incidence varies geographically. Higher incidence and prevalence have been reported among individuals from Mediterranean, Africa, the Middle East, and Southeast Asia. While its occurrence is rare in the United States, the pattern shows an increasing trend with migration and is expected to increase in the future. According to Bluebird Bio, approximately 1300 people in the United States currently live with transfusion-dependent β -thalassemia. (8)

Diagnosis

The diagnostic pathway for symptomatic thalassemia syndromes (thalassemia major and thalassemia intermedia) in a neonate, infant, or child begins with either recognition of symptoms (anemia, evidence of hemolysis and extramedullary hematopoiesis such as jaundice, skeletal abnormalities, and/or splenomegaly) or may be suspected based on a known family history. Initial laboratory testing includes a complete blood count, review of the blood smear, and iron studies. DNA-based genotyping of globin gene can be done relatively inexpensively, is required for precise diagnosis, and is especially important in carrier detection, prenatal testing, and genetic counseling. (5)

Treatment

The current standard of care for transfusion-dependent β -thalassemia includes blood transfusion, iron chelation therapies, and allogeneic hematopoietic stem cell transplant.

As per the 2021 Thalassemia International Federation guidelines, transfusion is indicated when hemoglobin levels are less than 7 g/dL on 2 different occasions more than 2 weeks apart, or based on clinical criteria such as significant symptoms of anemia, poor growth or failure to thrive, complications from excessive intramedullary hematopoiesis (e.g., pathologic fractures, facial changes), or clinically significant extramedullary hematopoiesis, irrespective of hemoglobin level. (9) The goal of treatment is to maintain a hemoglobin level of 9 to 10.5 g/dL, which has been shown to promote normal growth, suppress bone marrow activity, and minimize iron accumulation. (10, 11) Transfusions are typically required every 2 to 5 weeks to reach this goal but can vary for patients such as those with heart failure who may require higher target hemoglobin levels. (12) Risks of repeated blood transfusions include transfusion reactions, allergic reactions, hemolytic anemia, transfusion-related acute lung injury, and transfusion-related graft versus host disease and alloimmunization. (13) In the event of alloimmunization, it becomes difficult to find a matched blood and also increases the likelihood of delayed transfusion reactions. However, the main complication from frequent blood transfusions is iron overload.

Iron overload as a result of frequent transfusion results in iron accumulation in the heart, liver, and pituitary gland and can lead to heart failure, cirrhosis, hepatocellular carcinoma, hypothyroidism, hypoparathyroidism, hypogonadism, diabetes, and growth failure. (14) Primary treatment for iron overload is chelation therapy (deferioxamine, deferasirox, deferiprone) and is typically initiated after 10 to 20 transfusions or when the serum ferritin level rises above 1000 mcg/L. (9) Chelation therapy is associated with side effects such as hearing problems, bone growth retardation and local reactions, gastrointestinal symptoms, arthralgia, and neutropenia. Another limitation of chelation therapy is lack of adherence when infused therapies are used as compared to higher adherence for patients taking oral therapy. (15)

Hematopoietic stem cell transplant is the only curative treatment with cure rates ranging from 80% to 90% in children who receive human leukocyte antigen-identical sibling transplant. (16) Cure rates in adults are lower with a reported range of 65% to 70%. (17) While the cure rates are high, the main limiting factor for hematopoietic stem cell transplant is lack of a compatible donor. Fewer than 25% of patients have compatible related or unrelated donors, and transplants with mismatched donors or unrelated umbilical cord blood have a lower success rate. (18) Complications from hematopoietic stem cell transplant include mucositis, infection, graft failure, and graft versus host disease. If available, hematopoietic stem cell transplant should be offered to patients early in the disease course, prior to the onset of iron overload. (9)

There are no randomized trials comparing hematopoietic stem cell transplant with medical therapy for transfusion-dependent thalassemia. (19) Only a 2017 retrospective case-control study has been published, showing no statistically different overall survival with transplantation versus conventional medical therapy (e.g., transfusions and iron chelation). (17) The Center for International Blood and Marrow Transplant Research reported the results of a retrospective cohort of 1110 individuals with β -thalassemia who received a hematopoietic stem cell transplant between 2000 and 2016. The median age at transplantation was 6 years (range: 1 to 25 years), 61% received transplants with grafts from HLA-matched related donors, 7% from HLA-mismatched related donors, 23% from HLA-matched unrelated donors, and 9% from HLA-mismatched unrelated donors. The results are summarized in Table 2.

Table 2. Outcomes of Retrospective Cohort of Individuals who Received Hematopoietic Stem Cell Transplant for β -Thalassemia

Outcome	Matched Sibling	Matched Unrelated	Mismatched Relative	Mismatched Unrelated
5-year survival	89% (n=677)	87% (n=252)	73% (n=78)	83% (n=103)
Graft failure	8.6% (n=677)	5.2% (n=252)	21.8% (n=78)	10.7% (n=103)
Grade 2-4 acute GVHD	11.9% (n=674)	21.5% (n=251)	35.1% (n=77)	19.8% (n=101)
Chronic GVHD	8.3% (n=627)	8.4% (n=249)	20% (n=70)	23.8% (n=101)

GVHD: graft versus host disease.

Betibeglogene autotemcel (Zynteglo®)

Betibeglogene autotemcel (Zynteglo®) is the first cell-based gene therapy for the treatment of adult and pediatric patients with beta-thalassemia who require regular red blood cell transfusions. A one-time product administered as a single dose, Zynteglo is customized using the patient's own bone marrow stem cells that are genetically modified to produce functional beta-globin, a hemoglobin component.

Regulatory Status

On August 17, 2022, Zynteglo (betibeglogene autotemcel) was approved by the U.S. Food and Drug Administration (FDA) for the treatment of adult and pediatric patients with β -thalassemia who require regular red blood cell transfusions. It was granted a rare pediatric disorder voucher, as well as receiving Priority Review, Fast Track, Breakthrough Therapy, and Orphan designations from the FDA.

Rationale

Medical policies assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

TRANSFUSION DEPENDENT B-THALASSEMIA

Betibeglogene autotemcel

Clinical Context and Therapy Purpose

The purpose of betibeglogene autotemcel is to provide a treatment option that is an improvement on existing therapies. Potential benefits of this one-time therapy may include the following:

- Obviates the need for repeated blood transfusion thereby eliminating its downstream consequences such as iron overload and alloimmunization.
- Reduced complexity of one-time treatment.
- Novel mechanism of action or approach may allow successful treatment of patients for whom other available treatments have failed.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with transfusion-dependent β -thalassemia.

Interventions

The therapy being considered is betibeglogene autotemcel. In this gene therapy protocol, hematopoietic stem cells are mobilized using granulocyte colony stimulating factor and plerixafor followed by apheresis to obtain a CD34+ cell-enriched population. These cells are then transduced *ex vivo* by BB305 lentiglobin viral vector which adds functional copies of β^{A-T87Q} -globin gene that encodes β -globin protein. Patients receive myeloablative conditioning with busulfan to deplete endogenous hematopoietic stem cells, enabling therapeutic repopulation of the individual bone marrow with hematopoietic stem cells containing the transgene. The treatment with betibeglogene autotemcel requires inpatient hospitalization. Betibeglogene autotemcel must be administered in a qualified treatment center (hospital setting) by a physician(s) with experience in hematopoietic stem cell transplantation and treatment of patients with β -thalassemia.

Comparators

The following strategies are currently being used to make decisions about management of transfusion-dependent β -thalassemia: blood transfusion, iron chelation therapies, activin A traps or activin A receptor IIA ligands such as luspatercept, and allogenic hematopoietic stem cell transplant.

Outcomes

The general outcomes of interest are change in disease status, quality of life, hospitalizations, medication use, treatment-related mortality, and treatment-related morbidity (Table 3). Follow up at 5 years is of interest to monitor outcomes.

Table 3. Health Outcome Measures Relevant to Transfusion-Dependent Thalassemia

Outcome	Measure (Units)	Thresholds for Improvement/Decline or Clinically Meaningful Difference
Change in disease status	<ul style="list-style-type: none"> • Change in iron levels (including serum ferritin, liver iron) 	Independence from transfusion defined in clinical trials as “weighted average Hb \geq 9 g/dL without RBC

	<p>concentration, and myocardial iron deposition)</p> <ul style="list-style-type: none"> • Change in Hb levels • Reduction in transfusion frequency • Independence from transfusion • Reduction in severity of clinical sequelae iron overload such as pulmonary hypertension, cardiovascular events (e.g., arrhythmia and congestive heart failure), liver disease, venous thromboembolism, bone pain, etc. 	transfusions for ≥12 months.”
Quality of life	<ul style="list-style-type: none"> • Quality of life (in trials, generic and age-appropriate measures for quality of life such as Pediatric Quality of Life Inventory for pediatric and adolescent patients and EuroQol-5D and Short Form-36 were used). 	-
Hospitalizations	<ul style="list-style-type: none"> • Reduction in frequency or length of hospital admission • Reduction in frequency of ER visit 	Not applicable
Medication use	<ul style="list-style-type: none"> • Reduction or avoidance of iron-chelating therapy 	Not applicable
Treatment-related mortality	<ul style="list-style-type: none"> • Mortality 	Not applicable
Treatment-related morbidity	<ul style="list-style-type: none"> • Serious adverse events • Adverse events leading to treatment discontinuation 	Not applicable

ER: emergency room; Hb: hemoglobin; RBC: red blood cells

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Nonrandomized Studies

In the early phase of clinical development, 2 proof of concept studies HGB-205 (NCT02151526) and HGB-204 (NCT01745120) were conducted. (20, 21) The clinical response in these studies was less than expected. Subsequently, improvements in manufacturing process were made to enhance transduction to increase vector copy number and bolster clinical response. As such, these proof of concept studies were not included in the medical policy. The clinical development program of betibeglogene autotemcel for individuals with transfusion dependent β -thalassemia consists of 2 open-label, phase III, single-arm studies (HGB-207 and -212) that included a total of 41 study participants who received a single intravenous infusion of betibeglogene autotemcel. Of the 2 phase III studies, 1 has been published. (22) Of the 41 participants, 36 participants in whom transfusion independence was evaluable were included in the efficacy analysis. Transfusion independence was achieved in 89% (95% confidence interval [CI], 74% to 97%) of study participants. The median duration of transfusion independence was not reached at the time of data cut-off. Study characteristics, baseline characteristics and results are summarized in Tables 4 to 6.

Table 4. Summary of Key Nonrandomized Trials Characteristics

Study	Study Type	Country	Dates	Participants	Treatment	Follow-Up
HGB-207 (Northstar-2) NCT02906202 (22)	Single-arm prospective	US, EU, UK, Thailand	2016- 2022	<ul style="list-style-type: none"> • Non β^0/β^0 genotype • Age ≤ 50 years • Transfusion dependent β-thalassemia^a • Clinically stable and eligible to undergo HSCT <p>Primary endpoint:</p> <ul style="list-style-type: none"> • Transfusion independence (weighted average Hb ≥ 9 g/dL without RBC transfusions for ≥ 12 months at any time during the study after gene therapy infusion) 	Betibeglogene autotemcel (N=23)	Target: 2 years As of March 9, 2021: Median 29.5 months (range, 13.0 to 48.2)
HGB-212 (Northstar-3) NCT03207009	Single-arm prospective	US, EU, UK	2017- 2022	<ul style="list-style-type: none"> • β^0/β^0 genotype • $\beta^0/IVS1-110$ genotype 	Betibeglogene autotemcel (N=18)	Target: 2 years As of

				<ul style="list-style-type: none"> • IVS1-110/IVS1-110 genotype • Age ≤50 years old • Transfusion dependent β thalassemia • Clinically stable and eligible to undergo HSCT <p>Primary endpoint:</p> <ul style="list-style-type: none"> • Transfusion independence (weighted average Hb \geq 9 g/dL without RBC transfusions for \geq 12 months at any time during the study after gene therapy infusion) 		March 9, 2021: Median 24.6 months (range, 4.1 to 35.5)
--	--	--	--	---	--	---

^a \geq 100 mL/kg/year of RBCs or \geq 8 peripheral RBC transfusions/year, for prior 2 years.

EU: European Union; Hb: hemoglobin; HSCT: hematopoietic stem cell transplant; RBC: red blood cell; UK: United Kingdom; US: United States.

Table 5. Summary of Baseline Demographics and Disease Characteristics in the Pivotal Trials

Characteristic (23)	HGB-207 (n=23)	HGB-212 (n=18)
Genotype	non- β^0/β^0	12 β^0/β^0 ; 6 non- β^0/β^0
Age, median (range), years	15 (4, 34)	13 (4, 33)
Male, n (%)	48%	56%
Race, n (%)		
Asian	57%	39%
White	35%	56%
Other/not reported	9%	6%
Baseline ^a transfusion volume (mL/kg/year), median (min, max)	208 (142, 274)	194 (75, 289)
Baseline ^a transfusion frequency (transfusions per year), median (min, max)	16 (12, 37)	17 (11, 40)
Lansky or Karnofsky Performance Score		

All patients, minimum score	≥80	≥90
Percentage of patients with score of 100	52%	56%
Cardiac T2* at baseline (msec), median (min, max)	37 (21, 57)	37 (15, 75)
Serum ferritin at baseline (pmol/L), median (min, max)	4439 (784, 22517)	3275 (1279, 8874)
Liver iron concentration at baseline (mg/g), median (min, max)	5.3 (1, 41)	3.6 (1.2, 13.2)

^a Baseline annualized based on data 2 years prior to enrollment.

Table 6. Summary of Key Nonrandomized Trial Results

Study	Transfusion Independence	Weighted Average Hb During Transfusion Independence^b (g/dL)	Observed Duration of Transfusion Independence in months (median)	Grade 3 or 4 Adverse Events in >10% of Participants, %
HGB-207 (Northstar-2) (23)	23	20	20	-
Betibeglogene autotemcel	91% (20/22); 95% CI: 77% to 99%	11.8 (range: 9.7 to 13.0)	Not reached (range: 15.7 to 39.4)	-
HGB-212 (Northstar-3) (23)	18 ^a	12	12	-
Betibeglogene autotemcel	86% (12/14); 95% CI: 57% to 98%	10.2 (range: 9.3 to 13.7)	Not reached (range: 12.5 to 32.8)	-
Combined (HGB-207 and HGB-212) (23)	36	32	32	41
Betibeglogene autotemcel	89% (32/36); 95% CI: 74% to 97%	11.5 (range 9.3 to 13.7)	Not reached (range: 12.5 to 39.4)	Neutropenia: 100% Thrombocytopenia: 100% Leukopenia: 100% Anemia: 95% Lymphopenia: 61% ALT Increased: 24% Hypophosphatemia: 20%

				Hyperglycemia: 14% Hypokalemia: 12% Hyperbilirubinemia: 10% Hyponatremia: 10%
--	--	--	--	--

ALT: alanine aminotransferase; CI: confidence interval; Hb: hemoglobin.

^a 4 study participants were not evaluable for transfusion independence at the data cutoff of March 9, 2021

^b The weighted average Hb is an average area under the curve during the period of transfusion independence, from the start of transfusion independence when the Hb is first ≥ 9 g/dL with no transfusions in the preceding 60 days to the last available Hb at which the transfusion independence criteria are still met.

The purpose of the study limitations table (see Tables 7) is to display notable limitations identified in each study. This information is synthesized as a summary of the body of evidence following each table and provides the conclusions on the sufficiency of evidence supporting the position statement. In addition to a limited sample, the length of follow-up is not long enough to remove uncertainty regarding the durability of effect over a longer time period. To date, no study participants that became transfusion independent have reverted to becoming transfusion dependent. Study participants in the phase III studies have a median duration of follow-up between 24 and 29 months. Long-term follow-up (>15 years) is required to establish precision around durability of the treatment effect as well as side effects. No deaths were reported in any of the studies, but both mild side effects and serious adverse events were observed in the studies. The limited sample sizes of the studies creates uncertainty around the estimates of some of the patient-important outcomes, particularly adverse events. Some serious harms are likely rare occurrences and as such may not be observed in trials. While most of the serious adverse events were attributable to known risks associated with myeloablative conditioning, uncertainty still remains about the degree of risk with betibeglogene autotemcel infusion in real-world practice. Insertional oncogenesis has been identified as a potential risk with transgene integration. There has been no evidence of insertional oncogenesis and no malignancies in the trials of betibeglogene autotemcel. However, cases of myelodysplastic syndrome and acute myeloid leukemia have been reported in gene therapy trials that use a lentiviral vector to treat other conditions.

Table 7. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Combined (HGB-207 and HGB-212) (23)	4. Enrolled populations do not reflect relevant diversity.				1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

The study limitations stated in this table are those notable in the current literature review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5: Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Section Summary: Betibeglogene autotemcel

In the 2 pivotal, open-label, phase III single-arm studies, a total of 41 study participants received a single intravenous infusion of betibeglogene autotemcel. Of the 41 participants, 36 participants in whom transfusion independence was evaluable were included in the efficacy analysis. Transfusion independence was achieved in 89% (95% CI, 74% to 97%) of study participants. There is uncertainty regarding the durability of effect over a longer time period. Long-term follow-up (>15 years) is required to establish precision around durability of the treatment effect. The limited sample size creates uncertainty around the estimates of some of the patient-important outcomes, particularly adverse events. Some serious harms are likely rare occurrences and as such may not be observed in trials. While most of the serious adverse events were attributable to known risks associated with myeloablative conditioning, uncertainty still remains about the degree of risk of betibeglogene autotemcel infusion in real-world practice. Insertional oncogenesis has been identified as a potential risk with transgene integration. There has been no evidence of insertional oncogenesis and no malignancies in the trials of betibeglogene autotemcel. However, cases of myelodysplastic syndrome and acute myeloid leukemia have been reported in gene therapy trials that use a lentiviral vector to treat other conditions.

Summary of Evidence

For individuals with transfusion-dependent β -thalassemia who receive betibeglogene autotemcel, the evidence includes 2 single-arm studies: HGB-207 (Northstar-2) and HGB-212 (Northstar-3). The Northstar-2 trial enrolled non- $\beta^0\beta^0$ genotype (less severe phenotype) while Northstar-3 trial enrolled β -thalassemia patients with either a β^0 or β^+ IVS1 110 (G>A) variant (severe phenotype) at both alleles of the *HBB* gene. Relevant outcomes are change in disease status, quality of life, hospitalizations, medication use, treatment-related morbidity and treatment-related mortality. The 2 open-label, phase III, single-arm studies included a total of 41 individuals who received a single intravenous infusion of betibeglogene autotemcel. Of the 41 participants, 36 participants in whom transfusion independence was evaluable were included in the efficacy analysis. Transfusion independence was achieved in 89% (95% confidence interval, 74% to 97%) of study participants. There is uncertainty regarding the durability of effect over a longer time period. Long-term follow-up (>15 years) is required to

establish precision around durability of the treatment effect. Limited sample size creates uncertainty around the estimates of some of the patient-important outcomes, particularly adverse events. Some serious harms are likely rare occurrences and, as such, may not be observed in trials. While most of the serious adverse events were attributable to known risks associated with myeloablative conditioning, uncertainty still remains about the degree of risk with betibeglogene autotemcel infusion in real-world practice. Insertional oncogenesis has been identified as a potential risk with transgene integration. There has been no evidence of insertional oncogenesis and no malignancies in the trials of betibeglogene autotemcel. However, cases of myelodysplastic syndrome and acute myeloid leukemia have been reported in gene therapy trials that use a lentiviral vector to treat other conditions. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Practice Guidelines and Position Statements

Institute for Clinical and Economic Review

The Institute for Clinical and Economic Review published a final report on comparative effectiveness and value of betibeglogene autotemcel for beta thalassemia on July 19, 2022. (24) The Report concluded that betibeglogene autotemcel to be incremental or better with moderate certainty of a small or substantial net health benefit (“B+”) versus standard of care.

Ongoing and Unpublished Clinical Trials

Ongoing and/or unpublished trials that might influence this policy are listed in Table 8.

Table 8. Summary of Ongoing and Unpublished Trials

NCT Number	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT02633943 ^a	Long-term Follow-up of Subjects With Transfusion-Dependent β-Thalassemia Treated With Ex Vivo Gene Therapy	66	Nov 2035

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member’s benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

CPT Codes	None
------------------	------

HCPCS Codes	J3393
--------------------	-------

*Current Procedural Terminology (CPT®) ©2023 American Medical Association: Chicago, IL.

References

1. Borgna-Pignatti C. The life of patients with thalassemia major. *Haematologica*. Mar 2010; 95(3):345-348. PMID 20207838
2. Chieco P and Butler C. 2021 CAF Information - The Cooleys Anemia Foundation. Published January 21, 2022; Available at <<https://www.thalassemia.org>> (accessed January 21, 2024).
3. Arian M, Mirmohammadkhani M, Ghorbani R, et al. Health-related quality of life (HRQoL) in beta-thalassemia major (β -TM) patients assessed by 36-item short form health survey (SF-36): a meta-analysis. *Qual Life Res*. Feb 2019; 28(2):321-334. PMID 30194626
4. Vitrano A, Calvaruso G, Lai E, et al. The era of comparable life expectancy between thalassaemia major and intermedia: Is it time to revisit the major-intermedia dichotomy? *Br J Haematol*. Jan 2017; 176(1):124-130. PMID 27748513
5. Langer AL. Beta-Thalassemia. 2000 Sep 28 [Updated 2023 Jul 20]. In: Adam MP, Mirzaa GM, Pagon RA, et al., editors. *GeneReviews* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: <<https://www.ncbi.nlm.nih.gov>>.
6. Galanello R, Origa R. Beta-thalassemia. *Orphanet J Rare Dis*. May 21 2010; 5:11. PMID 20492708
7. Chonat S, Quinn CT. Current Standards of Care and Long Term Outcomes for Thalassemia and Sickle Cell Disease. *Adv Exp Med Biol*. 2017; 1013:59-87. PMID 29127677
8. beta-thalassemia (beta-thal). *Bluebird Bio*. Available at <<https://www.bluebirdbio.com>> (accessed January 21, 2024).
9. Farmakis D, Porter J, Taher A, et al. 2021 Thalassaemia International Federation Guidelines for the Management of Transfusion-dependent Thalassemia. *Hemasphere*. Aug 2022; 6(8):e732. PMID 35928543
10. Cazzola M, Borgna-Pignatti C, Locatelli F, et al. A moderate transfusion regimen may reduce iron loading in beta-thalassemia major without producing excessive expansion of erythropoiesis. *Transfusion*. Feb 1997; 37(2):135-140. PMID 9051086
11. Cazzola M, Locatelli F, De Stefano P. Deferoxamine in thalassemia major. *N Engl J Med*. Jan 26 1995; 332(4):271-272; author reply 272-273. PMID 7808503
12. Kremastinos DT, Farmakis D, Aessopos A, et al. Beta-thalassemia cardiomyopathy: history, present considerations, and future perspectives. *Circ Heart Fail*. May 2010; 3(3):451-458. PMID 20484195
13. Thompson AA, Cunningham MJ, Singer ST, et al. Red cell alloimmunization in a diverse population of transfused patients with thalassaemia. *Br J Haematol*. Apr 2011; 153(1):121-128. PMID 21323889
14. Olivieri NF, Liu PP, Sher GD, et al. Brief report: combined liver and heart transplantation for end-stage iron-induced organ failure in an adult with homozygous beta-thalassemia. *N Engl J Med*. Apr 21 1994; 330(16):1125-1127. PMID 8133854
15. Trachtenberg F, Vichinsky E, Haines D, et al. Iron chelation adherence to deferoxamine and deferasirox in thalassemia. *Am J Hematol*. May 2011; 86(5):433-436. PMID 21523808

16. Baronciani D, Angelucci E, Potschger U, et al. Hemopoietic stem cell transplantation in thalassemia: a report from the European Society for Blood and Bone Marrow Transplantation Hemoglobinopathy Registry, 2000–2010. *Bone Marrow Transplant.* Apr 2016; 51(4):536-541. PMID 26752139
17. Caocci G, Orofino MG, Vacca A, et al. Long-term survival of beta thalassemia major patients treated with hematopoietic stem cell transplantation compared with survival with conventional treatment. *Am J Hematol.* Dec 2017; 92(12):1303-1310. PMID 28850704
18. Angelucci E, Matthes-Martin S, Baronciani D, et al. Hematopoietic stem cell transplantation in thalassemia major and sickle cell disease: indications and management recommendations from an international expert panel. *Haematologica.* May 2014; 99(5):811-820. PMID 24790059
19. Sharma A, Jagannath VA, Puri L. Hematopoietic stem cell transplantation for people with β -thalassaemia. *Cochrane Database Syst Rev.* Apr 21 2021; 4(4):CD008708. PMID 33880750
20. Thompson AA, Walters MC, Kwiatkowski J, et al. Gene Therapy in Patients with Transfusion-Dependent β -Thalassemia. *N Engl J Med.* Apr 19 2018; 378(16):1479-1493. PMID 29669226
21. Magrin E, Semeraro M, Hebert N, et al. Long-term outcomes of lentiviral gene therapy for the β -hemoglobinopathies: the HGB-205 trial. *Nat Med.* Jan 2022; 28(1):81-88. PMID 35075288
22. Locatelli F, Thompson AA, Kwiatkowski JL, et al. Betibeglogene Autotemcel Gene Therapy for Non- β 0 / β 0 Genotype β -Thalassemia. *N Engl J Med.* Feb 03 2022; 386(5):415-427. PMID 34891223
23. FDA – Zynteglo® (betibeglogene autotemcel) Highlights of Prescribing Information. U.S. Food and Drug Administration: Prescribing Information (8/2022). Available at <<https://www.fda.gov>> (accessed November 20, 2024).
24. Institute for Clinical and Evidence Review. Betibeglogene Autotemcel for Beta Thalassemia: Effectiveness and Value (Final Evidence Report July 19, 2022). Available at <<https://icer.org>> (accessed January 21, 2024).

Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does not have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been developed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

Policy History/Revision

Date	Description of Change

01/01/2025	Document updated with literature review. Coverage unchanged. Added references 5, 9, and 24.
09/15/2023	Reviewed. No changes.
03/01/2023	<p>New medical document. Betibeglogene autotemcel (Zynteglo®) may be considered medically necessary if ALL the following criteria are met: adult and pediatric patients (≤50 years of age) with transfusion dependent beta-thalassemia; AND documented history of transfusions evidenced by: 100 mL/kg/year or more of packed red blood cells (pRBCs) OR; 8 or more transfusions of packed red blood cells (pRBCs) per year in the preceding 1 year; AND negative serologic test for HIV infection; AND clinically stable and eligible to undergo hematopoietic stem cell (HSC) mobilization; AND individual does NOT have ANY of the following: white blood cell count less than 3 x 10⁹/Liter and/or platelet count less than 100 x 10⁹ not related to hypersplenism; history of prior gene therapy or allogeneic hematopoietic stem cell transplant; severe iron overload; advanced liver disease; prior or current malignancy, myeloproliferative disorder, or significant immunodeficiency. Repeat treatment of Betibeglogene autotemcel (Zynteglo®) is considered experimental, investigational, and/or unproven. Betibeglogene autotemcel (Zynteglo®) is considered experimental, investigational, and/or unproven for all other indications.</p>