

<b>Policy Number</b>	<b>SUR703.048</b>
<b>Policy Effective Date</b>	<b>01/01/2024</b>
<b>Policy End Date</b>	<b>12/31/2024</b>

## Stem Cell Therapy for Peripheral Arterial Disease (PAD)

Table of Contents
<a href="#">Coverage</a>
<a href="#">Policy Guidelines</a>
<a href="#">Description</a>
<a href="#">Rationale</a>
<a href="#">Coding</a>
<a href="#">References</a>
<a href="#">Policy History</a>

Related Policies (if applicable)
None

### Disclaimer

**Carefully check state regulations and/or the member contract.**

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

### Coverage

Treatment of peripheral arterial disease (PAD), including critical limb ischemia, with injection or infusion of stem cells from concentrated bone marrow, expanded in vitro, stimulated from peripheral blood, or from an allogeneic source, **is considered experimental, investigational and/or unproven.**

### Policy Guidelines

None.

### Description

Peripheral arterial disease (PAD) is a common atherosclerotic syndrome associated with significant morbidity and mortality. Critical limb ischemia (CLI) is the end stage of lower-extremity PAD in which severe obstruction of blood flow results in ischemic pain at rest, ulcers,

and a significant risk for limb loss. Use of autologous stem cells freshly harvested and allogeneic stem cells are reported to have a role in the treatment of PAD.

## **Background**

### Peripheral Arterial Disease

PAD is a common atherosclerotic syndrome associated with significant morbidity and mortality. A less common cause of PAD is Buerger disease (also called thromboangiitis obliterans), which is a nonatherosclerotic segmental inflammatory disease that occurs in younger patients and is associated with tobacco use. The development of PAD is characterized by narrowing and occlusion of arterial vessels and eventual reduction in distal perfusion. CLI is the end-stage of lower-extremity PAD in which severe obstruction of blood flow results in ischemic pain at rest, ulcers, and a significant risk for limb loss.

### *Physiology*

Two endogenous compensating mechanisms may occur with occlusion of arterial vessels: capillary growth (angiogenesis) and development of collateral arterial vessels (arteriogenesis). Capillary growth is mediated by the hypoxia-induced release of chemokines and cytokines such as vascular endothelial growth factor (VEGF) and occurs by sprouting of small endothelial tubes from preexisting capillary beds. The resulting capillaries are small and cannot sufficiently compensate for a large, occluded artery. Arteriogenesis with collateral growth is, in contrast, initiated by increasing shear forces against vessel walls when blood flow is redirected from the occluded transport artery to the small collateral branches, leading to an increase in the diameter of preexisting collateral arterioles.

The mechanism underlying arteriogenesis includes the migration of bone marrow-derived monocytes to the perivascular space. The bone marrow-derived monocytes adhere to and invade the collateral vessel wall. It is not known if the expansion of the collateral arteriole is due to the incorporation of stem cells into the wall of the vessel or to cytokines released by monocytic bone marrow cells that induce the proliferation of resident endothelial cells. It has been proposed that bone marrow-derived monocytic cells may be the putative circulating endothelial progenitor cells. Notably, the same risk factors for advanced ischemia (diabetes, smoking, hyperlipidemia, advanced age) are also risk factors for a lower number of circulating progenitor cells.

### *Treatment*

Use of autologous stem cells freshly harvested and allogeneic stem cells are reported to have a role in the treatment of PAD. Stem cells can be administered in a variety of routes, derived from different progenitors, and be grouped with different co-factors, many of which are being studied in order to determine the best clinical option for patients. The primary outcome in stem cell therapy trials regulated by the U.S. Food and Drug Administration (FDA) is amputation-free survival, defined as time to major amputation and/or death from any cause. Other outcomes for CLI include the Rutherford criteria for limb status, healing of ulcers, the Ankle-Brachial Index, transcutaneous oxygen pressure, and pain-free walking. The Ankle-Brachial Index

measures arterial segmental pressures on the ankle and brachium and indexes ankle systolic pressure against brachial systolic pressure (normative range, 0.95 to 1.2 mm Hg).

### Regulatory Status

Several point-of-care concentrations of bone marrow aspirate have been cleared by the FDA through the 510(k) process and are summarized in Table 1.

**Table 1. The U.S. Food and Drug Administration Approved Point-of-Care Concentration of Bone Marrow Aspirate Devices**

Device	Manufacturer	Location	Date Cleared	510(k) Number
The SmartPReP Bone Marrow Aspirate Concentrate System, SmartPReP Platelet Concentration System	Harvest Technologies (now MD Biologix)	Lakewood, CO	12/06/2010	K103340
MarrowStim Concentration System (MSC system)	Biomet Biologics, Inc. (now Zimmer Biomet)	Warsaw, IN	12/18/2009	BK090008
PureBMC SupraPhysiologic Concentrating System	EmCyte Corporation®	Fort Myers, Florida	5/30/2019	K183205
Arthrex Angel® System Kit	Arthrex, Inc.	Naples, Florida	5/23/2018	BK180180
Magellan® Autologous Platelet Separator System	Arteriocyte Medical Systems (Medtronic)	Memphis, TN	11/09/2004	BK040068
BioCUE Platelet Concentration Kit (now BioCUE® Blood and Bone Marrow Aspiration (bBMA) Concentration Kit)	Biomet Biologics, Inc. (now Zimmer Biomet)	Warsaw, IN	5/26/2010	BK100027
ART BMC System/ART BMC PLUS System	SpineSmith Holdings, LLC (now Ceiling Biosciences)	Austin, TX	Not available	Not available
PXP® System (now PXP®-1000)	ThermoGenesis Corp.	Rancho Cordova, CA	07/10/2008	K081345

FDA product code: JQC.

## Rationale

This policy was created in 2011 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through December 2, 2022.

Medical policies assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life (QOL), and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

## **Stem Cell Therapy in Individuals with Peripheral Arterial Disease (PAD)**

### Clinical Context and Therapy Purpose

The purpose of stem cell therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with PAD.

The following PICO was used to select literature to inform this policy.

### *Populations*

The relevant population of interest is individuals with PAD.

### *Interventions*

The therapy being considered is stem cell therapy. The rationale for hematopoietic cell or bone marrow-cell therapy in PAD is to induce arteriogenesis by boosting the physiologic repair processes. This requires large numbers of functionally active autologous precursor cells and, subsequently, a large quantity of bone marrow (e.g., 240 to 500 mL) or another source of stem cells.

### *Comparators*

Comparators of interest include conservative management or surgical intervention. The standard therapy for severe, limb-threatening ischemia is revascularization aiming to improve blood flow to the affected extremity. If revascularization fails or is not possible, amputation is often necessary.

## *Outcomes*

The general outcomes of interest are overall survival, symptoms, change in disease status, morbid events, functional outcomes, QOL, and treatment-related morbidity including amputation rates, improved amputation-free survival, improved wound healing, ulcer healing, and pain-free walking distance. Follow-up at 3, 6, and 12 months is of interest for stem cell therapy to monitor relevant outcomes. Longer-term follow-up is also of interest.

## Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

At this time, the literature on stem cell therapy consists primarily of small RCTs, systematic reviews, and meta-analyses. (1, 2)

## Systematic Reviews

Several systematic reviews have been published (Table 2). Rigato et al. (2017) published a systematic review of autologous cell therapy for PAD. (3) The authors identified 19 RCTs (837 patients), 7 nonrandomized controlled studies (338 patients), and 41 noncontrolled studies (1177 patients). There was heterogeneity across studies in setting, underlying diseases, types and doses of cells, routes of administration, and follow-up durations. Many studies were pilot or phase 2 trials and were rated as low quality. There was an indication of publication bias. A meta-analysis of all RCTs showed a significant reduction in amputation rates, improved amputation-free survival, and improved wound healing. However, when only the placebo-controlled trials (n=19) were analyzed, the effects were no longer statistically significant, and analysis of only RCTs with a low risk of bias (n=3) found no benefit of cell therapy.

In a meta-analysis of RCTs, Xie et al. (2018) reviewed published evidence evaluating the safety and efficacy of autologous stem cell therapy in critical limb ischemia (CLI). (4) Cell therapy increased the probability of angiogenesis (relative risk=5.91, confidence interval [CI]: 2.49 to 14.02,  $p<0.0001$ ), ulcer healing (relative risk=1.73, CI: 1.45 to 2.06,  $p<0.00001$ ), and a reduction in amputation rates (relative risk=0.59, CI: 0.46-0.76,  $p<0.0001$ ). Compared with the control group, significant improvement in the cell therapy group was also seen in Ankle-Brachial Index (ABI) (mean difference=0.13, CI=0.11 to 0.15,  $p<0.00001$ ), transcutaneous oxygen tension (mean difference=12.22, CI=5.03 to 19.41,  $p=0.0009$ ), and pain-free walking distance (mean difference=144.84, CI=53.03 to 236.66,  $p=0.002$ ).

Gao et al. (2019) reviewed 27 RCTs and found that 1186 patients and 1280 extremities were included. (5) A majority of studies showed a high-risk of bias. Meta-analysis indicated that

autologous stem cell therapy was more effective than conventional therapy on the healing rate of ulcers. There was also a significant improvement in ABI, total carbon dioxide, and pain-free walking distance while a significant reduction was shown in amputation rate and rest pain scores. However, the result presented no significant improvement in major limb salvage.

Pu et al. (2022) included 12 RCTs (N=630) in a meta-analysis of patients with atherosclerosis obliterans (the most common type of PAD). (6) Autologous cell implantation was compared with placebo or standard care in all studies. A single injection of cell products was administered in all but 1 study in which injections were repeatedly administered. Follow-up periods ranged from 1 to 12 months. The analysis found improvements in total amputation (RR, 0.64; 95% CI, 0.47 to 0.87;  $p=.004$ ;  $I^2$ , 12%), major amputation (RR, 0.69; 95% CI, 0.50 to 0.94;  $p=.02$ ;  $I^2$ , 12%), and ABI (mean difference, 0.08; 95% CI, 0.02 to 0.13;  $p=.004$ ;  $I^2$ , 84%). Death and ulcer size were not improved with cell therapy. Findings of this analysis are applicable only to patients with no other therapy options. The analysis is limited by the small sample size in each trial (range, 10 to 160 patients) and heterogeneity in cell therapy methods (e.g., dosage, cell type, route of administration).

Moazzami et al. (2022) published a Cochrane review of 4 RCTs (N=176) in patients with CLI who were treated with autologous bone marrow mononuclear cells (BM-MNCs). (7) It was uncertain if amputations were lower (4 studies; RR, 0.52; 95% CI, 0.27 to 0.99), and mortality was not reduced with BM-MNCs (3 studies; RR, 1.0; 95% CI, 0.15 to 6.63). Data were limited by risk of bias, imprecision, and inconsistency.

**Table 2. Systematic Reviews of Trials Assessing Autologous Cell Therapy for Peripheral Arterial Disease**

Study (Year)	Literature Search	Studies	Participants	N	Design	Results
Moazzami et al. (2022) (7)	Nov 2021	4	Patients with CLI who were treated with local intramuscular transplantation of autologous adult BM-MNCs	176	RCTs	Pooled analysis of 4 RCTs found very low-to low-certainty evidence and no conclusion regarding BM-MNC for improving clinical outcomes can be drawn.
Pu et al. (2022) (6)	Mar 2021	12	Patients with atherosclerosis obliterans and "no available treatment" who received autologous cell therapy	630	RCTs	Pooled analysis of 12 RCTs showed a significant improvement in total amputation, major amputation, and ABI but not all-cause death or ulcer size.

Gao et al. (2019) (5)	May 2019	27	Patients with PAD or CLI who received autologous stem cell therapy	1186	RCTs	Pooled analysis of 27 RCTs showed a significant improvement in ABI, total carbon dioxide, and pain-free walking distance while significant reduction was showed in amputation rate and rest pain scores.
Rigato et al. (2017) (3)	Jul 2016	67	Patients with severe intractable PAD or CLI who received autologous cell therapy	2352	RCTs, cohort	Pooled analysis of 19 RCTs showed a reduction in amputation rates, improved amputation-free survival, and improved wound healing.
Xie et al. (2018) (4)	Jan 2018	23	Patients with PAD or CLI who received autologous stem cell therapy	1118	RCTs	Pooled analysis of 18 studies showed a reduction in amputation rate, ulcer healing, and pain-free walking distance (n=512).

ABI: Ankle-Brachial Index; CLI: critical limb ischemia; PAD: peripheral arterial disease, RCT: randomized controlled trial; BM-MNC: bone marrow mononuclear cells.

### Randomized and Nonrandomized Trials

#### *Concentrated Bone Marrow Aspirate (Monocytes and Mesenchymal Stem Cells [MSCs]) – Intramuscular Injection*

Prochazka et al. (2010) reported on a randomized study of 96 patients with CLI, and foot ulcers. (8) Patient inclusion criteria were CLI as defined by an ABI score of 0.4 or less, ankle systolic pressure of 50 mm Hg or less or toe systolic pressure of 30 mm Hg or less, and failure of basic conservative and revascularization treatment (surgical or endovascular). Patients were randomized to treatment with bone marrow concentrate (n=42) or standard medical care (n=54). The primary end points were major limb amputation during the 120 days posttreatment and degree of pain and function at 90- and 120-day follow-ups. At baseline, the control group compared with the treatment group had a higher proportion of patients with diabetes (98.2% versus 88.1%), hyperlipidemia (80.0% versus 54.8%), and ischemic heart disease (76.4% versus 57.1%), respectively. Additionally, the control group had a higher proportion of patients (72% versus 40%) with the University of Texas Wound Classification stage DIII (deep ulcers with osteitis). For the 42 patients in the treatment group, there was a history of 50 revascularization

procedures; 46 of 54 patients in the control group had a history of revascularization procedures. All 42 patients in the bone marrow group finished 90 days of follow-up, and 37 of 54 patients in the control group finished 120 days of follow-up. Differences in lengths of follow-up for the primary outcome measure were unexplained. Five patients in the bone marrow group and 8 in the control group died of causes unrelated to the therapy during follow-up. At follow-up, the frequency of major limb amputation was 21% in patients treated with bone marrow concentrate and 44% in controls. Secondary end points were assessed only in those treated with bone marrow concentrate. In the treatment group with salvaged limbs, toe pressure and Toe-Brachial Index score increased from 22.66 to 25.63 mm Hg and from 0.14 to 0.17, respectively. Interpretation of results is limited by unequal baseline measures, lack of blinding, differences in lengths of follow-up, differences in losses to follow-up, and differences in follow-up measures for the 2 groups.

Benoit et al. (2011) reported on a U.S. Food and Drug Administration (FDA)-regulated, double-blind pilot RCT of 48 patients with CLI who were randomized 2:1 to bone marrow concentrate using the SmartPreP system or to iliac crest puncture with an intramuscular injection of diluted peripheral blood. (9) At a 6-month follow-up, the differences in the percentages of amputations between the bone marrow concentrate group (29.4%) and diluted peripheral blood group (35.7%) were not statistically significant. In a subgroup analysis of patients with tissue loss at baseline (Rutherford 5), intramuscular injection of bone marrow concentrate resulted in a lower amputation rate (39.1%) than placebo (71.4%).

Intramuscular injection with a combination of bone marrow mononuclear cells (BM-MNCs) and gene therapy with a vascular endothelial growth factor (VEGF) plasmid was tested in a 2015 European RCT assessing 32 patients. (10) Controls in this trial were treated pharmacologically, and therefore the groups were not blinded to treatment. Several objective measures were improved in the BM-MNC group, but not in the control group. These measures included ABI scores, development of collateral vessels measured with angiography, and healing rates of ischemic ulcers. Amputations were performed in 25% of patients in the BM-MNC group and in 50% of patients in the control group.

Gupta et al. (2017) evaluated the efficacy and safety of intramuscular adult human bone marrow derived, cultured, pooled, allogeneic mesenchymal stromal cells (Stempeutics Research, Bangalore, India) in a phase II prospective, open-label dose-ranging study. (11) Ninety patients were nonrandomly allocated to 3 groups: 1 million cells/kg body weight (n=36), 2 million cells/kg body weight (n=36), and standard of care (SOC; n=18). Compared with the SOC group, greater reduction in rest pain and healing of ulcers were seen in the 2 million cells/kg body weight group (0.3 units per month [standard error (SE): 0.13], CI: -0.55 to -0.05, p=0.0193 and 11.0% decrease in size per month [SE: 0.05%], CI: 0.80 to 0.99, p=0.0253, respectively) and in the 1 million cells/kg body weight group (0.23 per month [SE: 0.13], CI: -0.49 to 0.03, p=0.081 and 2.0% decrease in size per month [SE: 0.06%], CI: 0.87 to 1.10, p=0.6967, respectively). Limitations of this study included the geographically and ethnically homogenous cohort and a lack of clearly defined methods for cohort selection. Additionally, patients in the cell



administration groups had lower ABI values and larger ulcers indicating potential investigator bias to allocate more severe patients to the treatment groups.

Dubsky et al. (2022) compared standard therapy with BM-MNC in patients with CLI and diabetic foot. (12) Forty patients with no-option chronic limb-threatening ischemia and no available treatment options were randomized to no treatment or BM-MNC for 12 weeks. Transcutaneous oxygen pressure (a marker of wound healing) had greater improvement in the BM-MNC group compared with no treatment (difference, 21.8 mm Hg;  $p=.034$ ). There were more healed ulcers at 12 weeks in the BM-MNC group (31.3% vs. 0%;  $p=.48$ ). The amputation rate and amputation-free survival was not different between groups. Although short-term improvements in outcomes were seen in this trial, the trial is limited by its small sample size, lack of placebo comparator, and single-center design.

#### *Concentrated Bone Marrow Aspirate (Monocytes and MSCs) – Intra-Arterial Injection*

The Rejuvenating Endothelial Progenitor Cells via Transcutaneous Intra-arterial Supplementation trial was a randomized, double-blind, placebo-controlled study (2015) from Europe. (13) This foundation-supported trial evaluated the clinical effects of repeated intra-arterial infusion of BM-MNCs in 160 patients with nonrevascularizable CLI. Patients received a repeated intra-arterial infusion of BM-MNCs or placebo (autologous peripheral blood erythrocytes) into the common femoral artery. The primary outcome measure (rate of major amputation after 6 months) did not differ significantly between groups (19% for BM-MNCs versus 13% controls). Secondary outcomes of QOL, rest pain, ABI score, and transcutaneous oxygen pressure improved to a similar extent in both groups, reinforcing the need for a placebo control in this type of trial. Results from a long-term follow-up analysis of 109 of the participants found improvements in self-reported QOL persisted for a median of 35 months in both groups, who remained blinded to treatment assignment. (14) The percentages of patients undergoing amputation also remained similar in the 2 groups (25.9% for the BM-MNC group versus 25.3% for the control group).

Results from the multicenter Intra-arterial Progenitor Cell Transplantation of Bone Marrow Mononuclear Cells for Induction of Neovascularization in Patients with Peripheral Arterial Occlusive Disease trial (2011) were reported. (15) In this double-blind, phase 2 trial, 40 patients with CLI who were not candidates or had failed to respond to interventional or surgical procedures were randomized to intra-arterial administration of BM-MNC or placebo. The cell suspension included hematopoietic, mesenchymal and other progenitor cells. After 3 months, both groups were treated with BM-MNC in an open-label phase. Twelve patients received additional treatment with BM-MNC between 6 months and 18 months. The primary outcome measure (a significant increase in the ABI score at 3 months) was not achieved (from 0.66 at baseline to 0.75 at 3 months). Limb salvage and amputation-free survival rates differed between groups. There was a significant improvement in ulcer healing (ulcer area, 1.89 cm<sup>2</sup> versus 2.89 cm<sup>2</sup>) and reduced pain at rest (an improvement on a 10-point visual analog scale score of  $\approx 3$  versus 0.05) following intra-arterial BM-MNC administration.

#### *Subsection Summary: Concentrated Bone Marrow Aspirate (Monocytes and MSCs)*

There is preliminary evidence of benefit to the use of intramuscular concentrated bone aspirate injection outcomes in CLI patients. Randomized controlled trials and a non-randomized comparative study have been published. (8, 9, 10, 11) Two RCTs have been published with intra-arterial injection of concentrated bone marrow aspirate. (13, 15) The RCTs did not find support for their respective primary outcome measures; the rate of major amputation after 6 months or a significant increase in the ABI score at 3 months.

#### *Expanded Monocytes and MSCs*

Interim and final results from the industry-sponsored phase 2, randomized, double-blind, placebo-controlled RESTORE-CLI trial, which used cultured and expanded monocytes and MSCs derived from bone marrow aspirate (ixmyelocel-T), were reported by Powell et al. (2011, 2012). (16, 17) Seventy-two patients with CLI received ixmyelocel-T (n=48) or placebo with sham bone marrow aspiration (n=24) and were followed for 12 months. There was a 40% reduction in any treatment failure (due primarily to differences in doubling of total wound surface area and de novo gangrene), but no significant differences in amputation rates at 12 months.

#### *Granulocyte-Macrophage Colony-Stimulating Factor Mobilization*

Poole et al. (2013) reported on the results of a phase 2, double-blind, placebo-controlled trial of granulocyte-macrophage colony-stimulating factor (GM-CSF) in 159 patients with intermittent claudication due to PAD. (18) Patients were treated with subcutaneous injections of GM-CSF or placebo 3 times weekly for 4 weeks. The primary outcome (peak treadmill walking time at 3 months) increased by 109 seconds (296 to 405 seconds) in the GM-CSF group and by 68 seconds (308 to 376 seconds) in the placebo group (p=0.08). Changes in the physical functioning subscale score of the 36-Item Short-Form Health Survey (SF-36) and distance score of the Walking Impairment Questionnaire were significantly better in patients treated with GM-CSF. However, there were no significant differences between the groups in ABI score, Walking Impairment Questionnaire distance or speed scores, claudication onset time, or SF-36 Mental Component or Physical Component Summary scores. The post hoc exploratory analysis found that patients with more than a 100% increase in progenitor cells (CD34-positive/CD133-positive) had a significantly greater increase in peak walking times (131 seconds) than patients who had less than 100% increase in progenitor cells (60 seconds).

McDermott et al. (2017) reported results from an RCT of 210 patients with PAD that evaluated whether GM-CSF combined with supervised treadmill exercise improves 6-minute walk distance (6MWT) compared with exercise alone and compared with GM-CSF alone and to determine whether GM-CSF alone improves 6MWT more than placebo and whether exercise improves 6MWT more than an attention control intervention. (19) Supervised exercise consisted of treadmill exercise 3 times weekly for 6 months. Participants were randomized to 1 of 4 groups: supervised exercise + GM-CSF (exercise + GM-CSF) (n = 53), supervised exercise + placebo (exercise alone) (n = 53), attention control + GM-CSF (GM-CSF alone) (n = 53), attention control + placebo (n = 51). The attention control consisted of weekly educational lectures by clinicians for 6 months. The primary outcome was change in 6MWT distance at a 12-week follow-up, with a minimum clinically important difference of 20 meters. Ninety-three percent of patients completed a 12-week follow-up. At follow-up, exercise + GM-CSF did not significantly improve

6MWT distance more than exercise alone ( $p = 0.61$ ) or more than GM-CSF alone (Hochberg-adjusted  $p = .052$ ). GM-CSF alone did not improve a 6MWT more than attention control + placebo ( $p = 0.91$ ). Exercise alone improved a 6MWT compared with attention control + placebo (Hochberg-adjusted  $p = 0.02$ ).

Horie et al. (2018) reported results from an RCT (IMPACT: Improvement of Peripheral Arterial Disease by Granulocyte Colony-Stimulating Factor-Mobilized Autologous Peripheral-Blood-Mononuclear Cell Transplantation) of 107 patients with PAD characterized as Buerger disease that evaluated the efficacy and safety of GM-CSF-mobilized peripheral blood mononuclear cell (PBMNC) transplantation compared with SOC (Tables 3 and 4). (20) Participants were randomized to guideline-based SOC or SOC plus intramuscular weight-based PBMNC administration. After disease progression or completion of a 1-year follow-up, 17 patients in the control group underwent cell therapy. Furthermore, 21 patients underwent revascularization after completion of the protocol treatment period or after discontinuation of the study (12 in the cell therapy group, 9 in the control group; 18 patients underwent percutaneous transluminal angioplasty, 2 had bypass surgery, and 1 had thrombectomy). Serious adverse events occurred in 20% of the cell therapy group compared with 11.3% of the control group ( $p = 0.28$ ). Leukopenia, alkaline phosphatase elevation, and hyperuricemia were determined to be adverse events related to GM-CSF administration. This study was limited by a small number of advanced cases (Fontaine stage IV cases [20.4%]), a high-risk group of hemodialysis patients and a high number of patients who did not complete treatment (cell therapy group: 38.5%; control group: 50.9%).

**Table 3. Key Characteristics of Randomized Controlled Trial with Intramuscular Granulocyte-Macrophage Colony-Stimulating Factor-Mobilized Peripheral Blood Mononuclear Cells for PAD**

Study (Year)	Countries	Sites	Dates	Participants	Treatment	
					Active	Comparator
Horie et al. (2018) (20) IMPACT	Japan	17	2009 to 2013	Patients with PAD, Fontaine classification II-IV (n=107)	Intramuscular GM-CSF, single dose of 200 $\mu\text{g}/\text{m}^2$ per day for 4 days (n=52)	Guideline based standard of care <sup>1</sup> (n=55)

GM-CSF: granulocyte-macrophage colony-stimulating factor; PAD: peripheral arterial disease.

<sup>1</sup> Includes the use of lipid, antihypertensive, antidiabetic, antithrombotic drugs, exercise, and prostanoids.

**Table 4. Results of Randomized Controlled Trial with Intramuscular Granulocyte-Macrophage Colony-Stimulating Factor-Mobilized Peripheral Blood Mononuclear Cells for PAD**

Study (Year)	PFS (95% CI)	Frequency of major limb amputation	New ulcer or gangrene	Serious AE
--------------	--------------	------------------------------------	-----------------------	------------

<b>Horie et al. (2018) (20) IMPACT</b>				
Cell Therapy group	0.42 (0.13-1.36)	6.0%	18%	20.0
Control group	0.62 (0.28-1.36)	5.7%	15.1%	11.3
p-value	0.07	1.00	0.80	0.28

AE: adverse events; CI: confidence intervals; PFS: progression-free survival.

The purpose of the limitations tables (see Tables 5 and 6) is to display notable limitations identified in the study. This information is synthesized as a summary of the body of evidence following each table and provides the conclusions on the sufficiency of evidence supporting the position statement.

**Table 5. Study Relevance Limitations**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Duration of Follow-up <sup>e</sup>
Horie et al. (2018) (20) IMPACT		3 - fixed dosing was used (not weight-based)			

The study limitations stated in this table are those notable in the current literature review; this is not a comprehensive gaps assessment.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

<sup>b</sup> Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

<sup>c</sup> Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

<sup>d</sup> Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

<sup>e</sup> Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

**Table 6. Study Design and Conduct Limitations**

Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness <sup>d</sup>	Power <sup>e</sup>	Statistical <sup>f</sup>
Horie et al. (2018) (20) IMPACT		1, 2, 3 - open-label trial				

The study limitations stated in this table are those notable in the current literature review; this is not a comprehensive gaps assessment.

<sup>a</sup> Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup> Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

<sup>c</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup> Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

<sup>e</sup> Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

<sup>f</sup> Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

### Subsection Summary: Granulocyte-Macrophage Colony-Stimulating Factor Mobilization

Three RCTs have been published. (18, 19, 20) The route of administration of cell therapy and the primary outcomes differed between studies. In the trial that added cell therapy to guideline-based care, there were no significant differences in PFS and frequency of limb amputation at 1 year of follow-up. (20) There was a substantial rate of subsequent surgical intervention in both arms.

### **Summary of Evidence**

For individuals who have peripheral arterial disease (PAD) who receive stem cell therapy, the evidence includes small, randomized trials and systematic reviews. Relevant outcomes are overall survival, symptoms, change in disease status, morbid events, functional outcomes, quality of life (QOL), and treatment-related morbidity. The current literature on stem cells as a treatment for critical limb ischemia (CLI) due to PAD consists primarily of phase 2 studies using various cell preparation methods and methods of administration. A meta-analysis of the trials with the lowest risk of bias has shown no significant benefit of stem cell therapy for overall survival, amputation-free survival, or amputation rates. Three randomized controlled trials (RCTs) have been published that used granulocyte colony-stimulating factor mobilized peripheral blood mononuclear cells. The route of administration of cell therapy and the primary outcomes differed between studies. In the trial that added cell therapy to guideline-based care, there were no significant differences in progression-free survival and frequency of limb amputation at 1 year of follow-up. There was a substantial rate of subsequent surgical intervention in both arms. Well-designed RCTs with a larger number of subjects and low risk of bias are needed to evaluate the health outcomes of these various procedures. Several are in progress, including multicenter randomized, double-blind, placebo-controlled trials. More data on the safety and durability of these treatments are also needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### **Practice Guidelines and Position Statements**

#### American Heart Association and the American College of Cardiology

In 2016, the guidelines from the American Heart Association and the American College of Cardiology provided recommendations on the management of patients with lower-extremity

peripheral arterial disease (PAD), including surgical and endovascular revascularization for CLI. (21, 22) Stem cell therapy for PAD was not addressed.

### Global Vascular Guideline

In 2019, a Global Vascular Guideline on management of chronic limb-threatening ischemia summarized the available literature on therapeutic angiogenesis for various etiologies. (23) The guideline was a joint venture of the Society for Vascular Surgery, the European Society for Vascular Surgery, and the World Federation of Vascular Societies. Based on a moderate level of evidence, the guideline recommended that therapeutic angiogenesis in patients with chronic limb-threatening ischemia should be limited to the context of a clinical trial (strong recommendation). The authors noted that Phase 3 clinical trials are planned or underway so additional data may be forthcoming in the future.

### European Society of Cardiology

In 2011, the European Society of Cardiology guidelines on the diagnosis and treatment of PAD did not recommend for or against stem cell therapy for PAD. (24) However, in 2017, updated guidelines, published in collaboration with the European Society of Vascular Surgery, stated: “Angiogenic gene and stem cell therapy are still being investigated with insufficient evidence in favor of these treatments.” The current recommendation is that stem cell/gene therapy is not indicated in patients with chronic limb-threatening ischemia (class of recommendation: III; level of evidence: B). (25)

### **Ongoing and Unpublished Clinical Trials**

Some currently ongoing and unpublished trials that might influence this policy are listed in Table 7.

**Table 7. Summary of Key Trials**

<b>NCT Number</b>	<b>Trial Name</b>	<b>Planned Enrollment</b>	<b>Completion Date</b>
<b><i>Ongoing</i></b>			
NCT03968198	Autologous Transplantation of Adipose Tissue Derived Mesenchymal Stroma/Stem Cells (ASC) in Patients With Critical Limb Ischemia	43	Jul 2023
NCT03304821	Granulocyte-Macrophage Stimulating Factor (GM-CSF) in Peripheral Artery Disease: the GPAD-3 Study	176	Sep 2022
NCT02685098	A Clinical and Histological Analysis of Mesenchymal Stem Cells in Amputation (CHAMP)	16	May 2026
NCT02805023 <sup>a</sup>	Phase 1/2, Double Blind Randomized Placebo Controlled Study to Assess the	50	Dec 2023

	Safety and Efficacy of BGC101 (EnEPC) in the Treatment of PAD & CLI		
<b>Unpublished</b>			
NCT04466007	Multicenter, Randomized, Dose-search, Parallel, Double-blind, and Placebo-controlled Clinical Trial to Evaluate the Safety and Efficacy of Intramuscular Administration of Allogeneic Adipose Tissue Adult Mesenchymal Stem Cells in Diabetic Patients With Critical Limb Ischemia Without Possibility of Revascularization	90	Sep 2022
NCT02551679 <sup>a</sup>	A Randomized Double Blind Placebo Controlled Clinical Study to Assess Blood-Derived Autologous Angiogenic Cell Precursor Therapy in Patients With Critical Limb Ischemia (ACP-CLI)	95	Dec 2020 (status unknown; last update Jan 2020)
NCT01745744	Clinical Trial Phase I / II, Multicentre, Open, Randomized Study of the Use of Mesenchymal Stem Cells From Adipose Tissue (CeTMAd) as Cell Regeneration Therapy in Critical Chronic Ischemic Syndrome of Lower Limb in Nondiabetic Patients	33	July 2018
NCT00498069 <sup>a</sup>	Feasibility Study of the Safety and Activity of Autologous Bone Marrow Aspirate Concentrate (BMAC) for the Treatment of Critical Limb Ischemia Due to Peripheral Arterial Occlusive Disease	48	Mar 2015
NCT02538978 <sup>a</sup>	Safety and Effectiveness of the SurgWerks <sup>TM</sup> -CLI Kit and VXPTM System for the Rapid Intra-operative Aspiration, Preparation and Intramuscular Injection of Concentrated Autologous Bone Marrow Cells Into the Ischemic Index Limb of Rutherford Category 5 Non-Reconstructable Critical Limb Ischemia Patients.	224	Mar 2019 (last update posted 2016 not yet recruiting)
NCT01679990 <sup>a</sup>	A Phase II, Randomized, Double-Blind, Multicenter, Multinational, Placebo-Controlled, Parallel-Groups Study to Evaluate the Safety and Efficacy of Intramuscular Injections of Allogeneic PLX-	180	Feb 2019 (last update Feb 2019)

	PAD Cells for the Treatment of Subjects With Intermittent Claudication (IC)		
NCT03042572	Allogeneic Mesenchymal Stromal Cells for Angiogenesis and Neovascularization in No-option Ischemic Limbs; A Double-blind, Randomized, Placebo-controlled Trial	60	Jul 2021 (unknown status; last update May 2018)
NCT01049919 <sup>a</sup>	MarrowStim PAD Kit for the Treatment of Critical Limb Ischemia (CLI) in Subjects With Severe Peripheral Arterial Disease (PAD) (MOBILE)	153	Feb 2020 (terminated; primary endpoint not met)
NCT03809494 <sup>a</sup>	Clinical Study of the Use of an Autologous Blood Filtration Device in the Treatment of Critical Limb Ischemia	2	Nov 2020 (Terminated)

NCT: national clinical trial.

<sup>a</sup> Denotes industry-sponsored or cosponsored trial.

## Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member's benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

<b>CPT Codes</b>	0263T, 0264T, 0265T
<b>HCPCS Codes</b>	None

\*Current Procedural Terminology (CPT®) ©2022 American Medical Association: Chicago, IL.

## References

1. Lawall H, Bramlage P, Amann B. Treatment of peripheral arterial disease using stem and progenitor cell therapy. J Vasc Surg. Feb 2011; 53(2):445-453. PMID 21030198
2. Fadini GP, Agostini C, Avogaro A. Autologous stem cell therapy for peripheral arterial disease meta-analysis and systematic review of the literature. Atherosclerosis. Mar 2010; 209(1):10-17. PMID 19740466
3. Rigato M, Monami M, Fadini GP. Autologous cell therapy for peripheral arterial disease: systematic review and meta-analysis of randomized, nonrandomized, and noncontrolled studies. Circ Res. Apr 14 2017; 120(8):1326-1340. PMID 28096194



4. Xie B, Luo H, Zhang Y, et al. Autologous Stem Cell Therapy in Critical Limb Ischemia: A Meta-Analysis of Randomized Controlled Trials. *Stem Cells Int.* 2018; 2018:7528464. PMID 29977308
5. Gao W, Chen D, Liu G, et al. Autologous stem cell therapy for peripheral arterial disease: a systematic review and meta-analysis of randomized controlled trials. *Stem Cell Res Ther.* May 21 2019; 10(1):140. PMID 31113463
6. Pu H, Huang Q, Zhang X, et al. A meta-analysis of randomized controlled trials on therapeutic efficacy and safety of autologous cell therapy for atherosclerosis obliterans. *J Vasc Surg. Apr 2022; 75(4):1440-1449.e5.* PMID 34788653
7. Moazzami B, Mohammadpour Z, Zabala ZE, et al. Local intramuscular transplantation of autologous bone marrow mononuclear cells for critical lower limb ischemia. *Cochrane Database Syst Rev.* Jul 08 2022; 7(7):CD008347. PMID 35802393
8. Prochazka V, Gumulec J, Jaluvka F, et al. Cell therapy, a new standard in management of chronic critical limb ischemia and foot ulcer. *Cell Transplant.* Jun 2010; 19(11):1413-1424. PMID 20529449
9. Benoit E, O'Donnell TF Jr, Iafrati MD, et al. The role of amputation as an outcome measure in cellular therapy for critical limb ischemia: implications for clinical trial design. *J Transl Med.* Sep 27 2011; 9:165. PMID 21951607
10. Skora J, Pupka A, Janczak D, et al. Combined autologous bone marrow mononuclear cell and gene therapy as the last resort for patients with critical limb ischemia. *Arch Med Sci.* Apr 25 2015; 11(2):325-331. PMID 25995748
11. Gupta PK, Krishna M, Chullikana A, et al. Administration of Adult Human Bone Marrow-Derived, Cultured, Pooled, Allogeneic Mesenchymal Stromal Cells in Critical Limb Ischemia Due to Buerger's Disease: Phase II Study Report Suggests Clinical Efficacy. *Stem Cells Transl Med.* Mar 2017; 6(3):689-699. PMID 28297569
12. Dubský M, Husáková J, Bem R, et al. Comparison of the impact of autologous cell therapy and conservative standard treatment on tissue oxygen supply and course of the diabetic foot in patients with chronic limb-threatening ischemia: A randomized controlled trial. *Front Endocrinol (Lausanne).* 2022; 13:888809. PMID 36105404
13. Teraa M, Sprengers RW, Schutgens RE, et al. Effect of repetitive intra-arterial infusion of bone marrow mononuclear cells in patients with no-option limb ischemia: The randomized, double-blind, placebo-controlled Rejuvenating Endothelial Progenitor Cells via Transcutaneous Intra-arterial Supplementation (JUVENTAS) Trial. *Circulation.* Mar 10 2015; 131(10):851-860. PMID 25567765
14. Peeters Weem SM, Teraa M, den Ruijter HM, et al. Quality of life after treatment with autologous bone marrow derived cells in no option severe limb ischemia. *Eur J Vasc Endovasc Surg.* Jan 2016; 51(1):83-89. PMID 26511056
15. Walter DH, Krankenberg H, Balzer JO, et al. Intraarterial administration of bone marrow mononuclear cells in patients with critical limb ischemia: a randomized-start, placebo-controlled pilot trial (PROVASA). *Circ Cardiovasc Interv.* Feb 1 2011; 4(1):26-37. PMID 21205939
16. Powell RJ, Comerota AJ, Berceli SA, et al. Interim analysis results from the RESTORE-CLI, a randomized, double-blind multicenter phase II trial comparing expanded autologous bone

- marrow-derived tissue repair cells and placebo in patients with critical limb ischemia. *J Vasc Surg.* Oct 2011; 54(4):1032-1041. PMID 21684715
17. Powell RJ, Marston WA, Berceci SA, et al. Cellular therapy with Ixmyelocel-T to treat critical limb ischemia: the randomized, double-blind, placebo-controlled RESTORE-CLI trial. *Mol Ther.* Jun 2012; 20(6):1280-1286. PMID 22453769
  18. Poole J, Mavromatis K, Binongo JN, et al. Effect of progenitor cell mobilization with granulocyte-macrophage colony-stimulating factor in patients with peripheral artery disease: a randomized clinical trial. *JAMA.* Dec 25 2013; 310(24):2631-2639. PMID 24247554
  19. McDermott MM, Ferrucci L, Tian L, et al. Effect of granulocyte-macrophage colony-stimulating factor with or without supervised exercise on walking performance in patients with peripheral artery disease: The PROPEL randomized clinical trial. *JAMA.* Dec 5 2017; 318(21):2089-2098. PMID 29141087
  20. Horie T, Yamazaki S, Hanada S, et al. Outcome from a randomized controlled clinical trial - improvement of peripheral arterial disease by granulocyte colony-stimulating factor-mobilized autologous peripheral-blood-mononuclear cell transplantation (IMPACT). *Circ J.* Jul 25 2018; 82(8):2165-2174. PMID 29877199
  21. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the management of patients with lower extremity peripheral artery disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* Mar 21 2017; 69(11):e71-e126. PMID 27851992
  22. Valentine EA, Ochroch EA. 2016 American College of Cardiology/American Heart Association guideline on the management of patients with lower extremity peripheral artery disease: perioperative implications. *J Cardiothorac Vasc Anesth.* Oct 2017; 31(5):1543-1553. PMID 28826846
  23. Conte MS, Bradbury AW, Kolh P, et al. Global vascular guidelines on the management of chronic limb-threatening ischemia. *J Vasc Surg.* Jun 2019; 69(6S):3S-125S.e40. PMID 31159978
  24. Tendera M, Aboyans V, Bartelink ML, et al. ESC Guidelines on the diagnosis and treatment of peripheral artery diseases: Document covering atherosclerotic disease of extracranial carotid and vertebral, mesenteric, renal, upper, and lower extremity arteries: the Task Force on the Diagnosis and Treatment of Peripheral Artery Diseases of the European Society of Cardiology (ESC). *Eur Heart J.* Nov 2011; 32(22):2851-906. PMID 21873417
  25. Aboyans V, Ricco JB, Bartelink MEL, et al. 2017 ESC Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with the European Society for Vascular Surgery (ESVS): Document covering atherosclerotic disease of extracranial carotid and vertebral, mesenteric, renal, upper and lower extremity arteries Endorsed by: the European Stroke Organization (ESO) The Task Force for the Diagnosis and Treatment of Peripheral Arterial Diseases of the European Society of Cardiology (ESC) and of the European Society for Vascular Surgery (ESVS). *Eur Heart J.* Mar 01 2018; 39(9):763-816. PMID 28886620

## Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does not have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been developed since this medical policy document was written. See Medicare's National Coverage at <<http://www.cms.hhs.gov>>.

<b>Policy History/Revision</b>	
<b>Date</b>	<b>Description of Change</b>
01/01/2024	Document updated with literature review. Coverage unchanged. References 6, 7, and 12 added; others removed.
05/01/2022	Document updated with literature review. Coverage unchanged. Reference 20 added; some updated and others removed.
08/01/2021	Reviewed. No changes.
07/15/2020	Document updated with literature review. Coverage unchanged. References 4, 6, 10, 18, and 19 were added and some removed.
05/01/2019	Reviewed. No changes.
08/01/2018	Document updated with literature review. Coverage updated to describe specific sources of stem-cells. References 3, 14-15, and 17 added.
01/15/2018	Reviewed. No changes.
07/15/2016	Document updated with literature search. Coverage unchanged.
03/01/2015	Document updated with literature search. Coverage unchanged.
08/15/2013	Document updated with literature search. Title changed from Autologous Cell Therapy for the Treatment of Peripheral Arterial Disease. Coverage statement changed from “autologous cell therapy” to “stem-cell therapy”; otherwise, coverage unchanged. Rationale reorganized.
07/01/2011	New medical document. Autologous cell therapy, including injection or infusion of cells concentrated from hematopoietic bone marrow stem-cells aspirate, is considered experimental, investigational and unproven as a treatment of peripheral arterial disease, including critical limb ischemia.