

Policy Number	SUR704.012
Policy Effective Date	07/01/2024

Bioengineered Skin and Soft Tissue Substitutes

Table of Contents
Coverage
Policy Guidelines
Description
Rationale
Coding
References
Policy History

Related Policies (if applicable)
SUR704.011 Amniotic Membrane and Amniotic Fluid

Disclaimer

Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

Legislative Mandates

EXCEPTION: For Illinois only: Illinois Public Act 103-0123 (IL HB 1384) Coverage for Reconstructive Services requires the following policies amended, delivered, issued, or renewed on or after January 1, 2025 (Individual and family PPO/HMO/POS; Student; Group [Small Group; Mid-Market; Large Group Fully Insured PPO/HMO/POS] or Medicaid), to provide coverage for medically necessary services that are intended to restore physical appearance on structures of the body damaged by trauma.

EXCEPTION: For HCSC members residing in the state of Arkansas, § 23-99-405 related to coverage of mastectomy and reconstruction services, should an enrollee elect reconstruction after a mastectomy, requires coverage for surgery and reconstruction of the breast on which the mastectomy has been performed, surgery and reconstruction of the other breast to produce a symmetrical appearance, and prostheses and coverage for physical complications at all stages of a mastectomy, including lymphedema. This applies to the following: Fully Insured Group, Student, Small Group, Mid-Market, Large Group, HMO, EPO, PPO, POS. Unless indicated by the group, this mandate or coverage will not apply to ASO groups.

Coverage

NOTE 1: This policy does not address amniotic membrane and amniotic fluid products. See SUR704.011 Amniotic Membrane and Amniotic Fluid for coverage position on those products.

Breast Reconstructive Surgery

Breast reconstructive surgery using allogeneic acellular dermal matrix products^a including each of the following: AlloDerm[®], AlloMend[®], Cortiva[®] [AlloMax[™]], DermACELL[™], DermaMatrix[™], FlexHD[®], FlexHD[®] Pliable[™], GraftJacket[®] **may be considered medically necessary:**

- When there is insufficient tissue expander or implant coverage by the pectoralis major muscle and additional coverage is required;
- When there is viable but compromised or thin postmastectomy skin flaps that are at risk of dehiscence or necrosis; or
- The inframammary fold and lateral mammary folds have been undermined during mastectomy and reestablishment of these landmarks is needed.

Diabetic Lower-Extremity Ulcers

Treatment of chronic, noninfected, full-thickness diabetic lower-extremity ulcers using the following tissue-engineered skin substitutes **may be considered medically necessary:**

- AlloPatch^{®a};
- Apligraf^{®b};
- Dermagraft^{®b};
- Integra[®] Omnigraft[™] Dermal Regeneration Matrix (also known as Omnigraft[™]); and Integra Flowable Wound Matrix;
- mVASC[®];
- TheraSkin[®].

Lower-Extremity Skin Ulcers due to Venous Insufficiency

Treatment of chronic, noninfected, partial- or full-thickness lower-extremity skin ulcers due to venous insufficiency, which have not adequately responded following a 1-month period of conventional ulcer therapy, using the following tissue-engineered skin substitutes **may be considered medically necessary:**

- Apligraf^{®b}; or
- Oasis[™] Wound Matrix^c.

Dystrophic Epidermolysis Bullosa

Treatment of dystrophic epidermolysis bullosa using the following tissue-engineered skin substitutes **may be considered medically necessary:**

- OrCel[™] (for the treatment of mitten-hand deformity when standard wound therapy has failed and when provided in accordance with the humanitarian device exemption [HDE] specifications of the U.S. Food and Drug Administration [FDA])^d.

Burns

Treatment of second- and third-degree burns using the following tissue-engineered skin substitutes **may be considered medically necessary:**

- Epicel® (for the treatment of deep dermal or full-thickness burns comprising a total body surface area $\geq 30\%$ when provided in accordance with the HDE specifications of the FDA)^d;
or
- Integra® Dermal Regeneration Template^b.

^aBanked human tissue.

^bFDA premarket approval.

^cFDA 510(k) clearance.

^dFDA-approved under an HDE.

All other uses of the bioengineered skin and soft tissue substitutes listed above are **considered experimental, investigational and/or unproven.**

All other skin and soft tissue substitutes not listed above are considered **experimental, investigational and/or unproven**, including, but not limited to:

- AC5® Advanced Wound System;
- ACell® UBM Hydrated/Lyophilized Wound Dressing;
- AlloSkin™;
- AlloSkin™ RT;
- Aongen™ Collagen Matrix;
- Apis®;
- Architect® ECM, PX, FX;
- Artacent® Wound
- ArthroFlex™ (Flex Graft);
- AxoGuard® Nerve Protector (AxoGen);
- BellaCell HD or Surederm;
- Biobrane®/Biobrane-L;
- Bio-ConneKt® Wound Matrix;
- CollaCare®;
- CollaCare® Dental;
- Collagen Wound Dressing (Oasis Research);
- CollaGUARD®;
- CollaMend™;
- CollaWound™;
- Coll-e-derm;
- Collexa®;
- Collieva®;
- Conexa™;
- Coreleader Colla-Pad;
- CorMatrix®;
- Cymetra™ (Micronized AlloDerm™);
- Cytal™ (previously MatriStem®);

- DeNovoSkin™
- Dermadapt™ Wound Dressing;
- Derma-gide;
- DermaPure™;
- DermaSpan™;
- DressSkin;
- Duragen® XS, Duragen™ Plus;
- Durepair Regeneration Matrix®;
- Endoform Dermal Template™;
- ENDURAGen™;
- Excellagen;
- ExpressGraft™;
- E-Z Derm™;
- FlowerDerm™;
- GammaGraft;
- Geistlich Derma-Gine™;
- GraftJacket® Xpress, injectable;
- Helicoll™;
- hMatrix®;
- Hyalomatrix®;
- Hyalomatrix® PA;
- Innovaburn™;
- Innovamatrix® XL;
- Innovamatrix® PD;
- Innovamatrix FS;
- Integra™ Bilayer Wound Matrix;
- Integra® Matrix Wound Dressing (previously Avagen);
- InteguPly®;
- Keramatrix®;
- Kerecis™ Omega3;
- Keroxx™;
- MatriDerm®;
- MatriStem;
- Matrix HD™;
- Mediskin®;
- MemoDerm™;
- Microderm®;
- Microderm® biologic wound matrix;
- Microlyte® Matrix;
- MicroMatrix®;
- Miro3d® Wound Matrix;
- MiroTract® Wound Matrix;

- Mirragen® Advanced Wound Matrix;
- MyOwn skin;
- NeoMatriX® Wound Matrix;
- NovoSorb SynPath;
- Oasis® Burn Matrix;
- Oasis® Ultra;
- Ologen™ Collagen Matrix;
- Omega3 Wound (originally Merigen wound dressing);
- Omeza Collagen Matrix®
- OvaTex
- Permacol™;
- PermeaDermB, PermeaDerm Glove, PermeaDerm C
- Phoenix Wound Matrix;
- PriMatrix™;
- PriMatrix™ Dermal Repair Scaffold;
- Progenamatrix;
- Puracol® and Puracol® Plus Collagen Wound Dressings;
- PuraPly™ Wound Matrix (previously FortaDerm™);
- PuraPly™ AM (Antimicrobial Wound Matrix);
- Puros® Dermis;
- RegenePro™;
- Repliform®;
- ReCell®;
- Repriza™;
- Resolve Matrix™;
- Restrata;
- SimpiDerm;
- Skin TE™;
- StrataGraft®;
- Strattice™ (xenograft);
- Suprathel®;
- Supra SDRM;
- SUPRATHE;
- SurgiMend®;
- Symphony™;
- Talymed®;
- TenoGlide™;
- TenSIX™ Acellular Dermal Matrix;
- TissueMend;
- TheraForm™ Standard/Sheet;
- TheraGenesis;
- TransCyte™;

- TruSkin™;
- Tutomesh™ Fenestrated Bovine Pericardium;
- Veritas® Collagen Matrix;
- Xcellistem;
- XCM Biologic® Tissue Matrix; or,
- XenMatrix™ AB.

Policy Guidelines

None.

Description

Skin and Soft Tissue Substitutes

Bioengineered skin and soft tissue substitutes may be either acellular or cellular. Acellular products (e.g., dermis with cellular material removed) contain a matrix or scaffold composed of materials such as collagen, hyaluronic acid, and fibronectin. Acellular dermal matrix (ADM) products can differ in a number of ways, including as species source (human, bovine, porcine), tissue source (e.g., dermis, pericardium, intestinal mucosa), additives (e.g., antibiotics, surfactants), hydration (wet, freeze-dried), and required preparation (multiple rinses, rehydration).

Cellular products contain living cells such as fibroblasts and keratinocytes within a matrix. The cells contained within the matrix may be autologous, allogeneic, or derived from other species (e.g., bovine, porcine). Skin substitutes may also be composed of dermal cells, epidermal cells, or a combination of dermal and epidermal cells, and may provide growth factors to stimulate healing. Bioengineered skin substitutes can be used as either temporary or permanent wound coverings.

Applications

There are a large number of potential applications for artificial skin and soft tissue products. One large category is nonhealing wounds, which potentially encompasses diabetic neuropathic ulcers, vascular insufficiency ulcers, and pressure ulcers. A substantial minority of such wounds do not heal adequately with standard wound care, leading to prolonged morbidity and increased risk of mortality. For example, nonhealing lower-extremity wounds represent an ongoing risk for infection, sepsis, limb amputation, and death. Bioengineered skin and soft tissue substitutes have the potential to improve rates of healing and reduce secondary complications.

Other situations in which bioengineered skin products might substitute for living skin grafts include certain postsurgical states (e.g., breast reconstruction) in which skin coverage is inadequate for the procedure performed or for surgical wounds in patients with compromised ability to heal. Second- and third-degree burns are another indication in which artificial skin

products may substitute for auto- or allografts. Certain primary dermatologic conditions that involve large areas of skin breakdown (e.g., bullous diseases) may also be conditions in which artificial skin products can be considered as substitutes for skin grafts. ADM products are also being evaluated in the repair of other soft tissues including rotator cuff repair, following oral and facial surgery, hernias, and other conditions.

Regulatory Status

The U.S. Food and Drug Administration (FDA) does not refer to any single product or class of products as “skin substitutes”. Products in this review cover products that do not require FDA approval or clearance as well as a number of products cleared through the 510(k) pathway with a variety of FDA product codes. A large number of artificial skin products are commercially available or in development. Commercial availability is not a reflection of a product's regulatory status. The following section summarizes a subset of commercially available skin and soft-tissue substitutes. This is not a complete list of all commercially available products. Information on additional products is available in a 2020 Technical Brief on skin substitutes for treating chronic wounds that was commissioned by the Agency for Healthcare Research and Quality. (1)

Acellular Dermal Matrix (ADM) Products

Allograft ADM products derived from cadaveric human skin tissue are supplied by tissue banks compliant with standards of the American Association of Tissue Banks and FDA guidelines. The processing removes the cellular components (i.e., epidermis, all viable dermal cells) that can lead to rejection and infection. ADM products from human skin tissue are regarded as minimally processed and not significantly changed in structure from the natural material; FDA classifies ADM products as banked human tissue and therefore, not requiring FDA approval for homologous use.

In 2017, the FDA published clarification of what is considered minimal manipulation and homologous use for human cells, tissues, and cellular and tissue-based products (HCT/Ps). (2)

HCT/Ps are defined as human cells or tissues that are intended for implantation, transplantation, infusion, or transfer into a human recipient. If an HCT/P does not meet the criteria below and does not qualify for any of the stated exceptions, the HCT/P will be regulated as a drug, device, and/or biological product and applicable regulations and premarket review will be required.

An HCT/P is regulated solely under section 361 of the PHS Act and 21 CFR Part 1271 if it meets all of the following criteria:

- 1) “The HCT/P is minimally manipulated;
- 2) The HCT/P is intended for homologous use only, as reflected by the labeling, advertising, or other indications of the manufacturer’s objective intent;
- 3) The manufacture of the HCT/P does not involve the combination of the cells or tissues with another article, except for water, crystalloids, or a sterilizing, preserving, or storage agent, provided that the addition of water, crystalloids, or the sterilizing, preserving, or storage agent does not raise new clinical safety concerns with respect to the HCT/P; and

4) Either:

- i) The HCT/P does not have a systemic effect and is not dependent upon the metabolic activity of living cells for its primary function; or
- ii) The HCT/P has a systemic effect or is dependent upon the metabolic activity of living cells for its primary function, and: a) Is for autologous use; b) Is for allogeneic use in a first-degree or second-degree blood relative; or c) Is for reproductive use."

- AlloDerm® (LifeCell Corp.) is an ADM (allograft) tissue-replacement product created from native human skin and processed so that the basement membrane and cellular matrix remain intact. Originally, AlloDerm® required refrigeration and rehydration before use. It is currently available in a ready-to-use product stored at room temperature. An injectable micronized form of AlloDerm® (Cymetra) is available.
- AlloPatch® (Musculoskeletal Transplant Foundation) is an acellular human dermis allograft derived from the reticular layer of the dermis and marketed for wound care. This product is also marketed as FlexHD® for postmastectomy breast reconstruction.
- Cortiva® (previously marketed as AlloMax™ Surgical Graft and before that NeoForm™) is an acellular non-cross-linked human dermis allograft.
- FlexHD® and the newer formulation FlexHD® Pliable™ (Musculoskeletal Transplant Foundation) are acellular hydrated reticular dermis allograft derived from donated human skin.
- DermACELL™ (LifeNet Health) is an allogeneic ADM processed with proprietary technologies MATRACELL® and PRESERVON®.
- DermaMatrix™ (Synthes) is a freeze-dried ADM derived from donated human skin tissue. DermaMatrix Acellular Dermis is processed by the Musculoskeletal Transplant Foundation.
- DermaPure™ (Tissue Regenix Wound Care) is a single-layer decellularized human dermal allograft for the treatment of acute and chronic wounds.
- GraftJacket® Regenerative Tissue Matrix (also called GraftJacket Skin Substitute; KCI) is an acellular regenerative tissue matrix that has been processed from human skin supplied from U.S. tissue banks. The allograft is minimally processed to remove the epidermal and dermal cells while preserving dermal structure. GraftJacket Xpress® is an injectable product.
- mVASC® (MicroVascular Tissues, Inc.) is a microvascular tissue structural allograft made of small blood vessels and extracellular matrix, inherent non-viable cells, and associated biological signaling factors harvested from subcutaneous tissue of cadaveric human donors.
- TheraSkin® (LifeNet Health) is a cryopreserved split-thickness human skin allograft composed of living fibroblasts and keratinocytes and an extracellular matrix in epidermal and dermal layers. TheraSkin® is derived from human skin allograft supplied by tissue banks compliant with the American Association of Tissue Banks and FDA guidelines. It is considered a minimally processed human cell, tissue, and cellular- and tissue-based product by the FDA.

Although frequently used by surgeons for breast reconstruction, the FDA does not consider this homologous use and has not cleared or approved any surgical mesh device (synthetic, animal collagen-derived, or human collagen-derived) for use in breast surgery. The indication of

surgical mesh for general use in “Plastic and reconstructive surgery” was cleared by the FDA before surgical mesh was described for breast reconstruction in 2005. The FDA states that the specific use of surgical mesh in breast procedures represents a new intended use and that a substantial equivalence evaluation via 510(k) review is not appropriate and a pre-market approval evaluation is required. (3)

In March 2019, the FDA held an Advisory Committee meeting on breast implants, at which time the panel noted that while there is data about ADM for breast reconstruction, the FDA has not yet determined the safety and effectiveness of ADM use for breast reconstruction. The panel recommended that patients are informed and also recommended studies to assess the benefit and risk of ADM use in breast reconstruction. (3)

In March 2021, FDA issued a Safety Communication to inform patients, caregivers, and health care providers that certain ADM products used in implant-based breast reconstruction may have a higher chance for complications or problems. An FDA analysis of patient-level data from real-world use of ADMs for implant-based breast reconstruction suggested that 2 ADMs—FlexHD and Allomax—may have a higher risk profile than others. (4)

In October 2021, an FDA advisory panel on general and plastic surgery voted against recommending FDA approval of the SurgiMend mesh for the specific indication of breast reconstruction. The advisory panel concluded that the benefits of using the device did not outweigh the risks. (4)

FDA product codes: FTM, OXF.

Xenogenic Products

Cytal™ (previously called MatriStem®) Wound Matrix, Multilayer Wound Matrix, Pelvic Floor Matrix, MicroMatrix, and Burn Matrix (all manufactured by ACell) are composed of porcine-derived urinary bladder matrix.

Helicoll (Encol) is an acellular collagen matrix derived from bovine dermis. In 2004, it was cleared for marketing by the FDA through the 510(k) process for topical wound management that includes partial and full-thickness wounds, pressure ulcers, venous ulcers, chronic vascular ulcers, diabetic ulcers, trauma wounds (e.g., abrasions, lacerations, second-degree burns, skin tears), and surgical wounds including donor sites/grafts.

Keramatrix® (Keraplast Research) is an open-cell foam comprised of freeze-dried keratin that is derived from acellular animal protein. In 2009, it was cleared for marketing by the FDA through the 510(k) process under the name of Keratec. The wound dressings are indicated in the management of the following types of dry, light, and moderately exudating partial and full-thickness wounds: pressure (stage I-IV) and venous stasis ulcers, ulcers caused by mixed vascular etiologies, diabetic ulcers, donor sites, and grafts.

Kerecis™ Omega3 Wound (Kerecis) is an ADM derived from fish skin. It has a high content of omega 3 fatty acids and is intended for use in burn wounds, chronic wounds, and other applications.

Oasis™ Wound Matrix (Cook Biotech) is a collagen scaffold (extracellular matrix) derived from porcine small intestinal submucosa. In 2000, it was cleared for marketing by the FDA through the 510(k) process for the management of partial- and full-thickness wounds, including pressure ulcers, venous ulcers, diabetic ulcers, chronic vascular ulcers, tunneled undermined wounds, surgical wounds, trauma wounds, and draining wounds.

Permacol™ (Covidien) is xenogeneic and composed of cross-linked porcine dermal collagen. Cross-linking improves the tensile strength and long-term durability but decreases pliability.

PriMatrix™ (TEI Biosciences; a subsidiary of Integra Life Sciences) is a xenogeneic ADM processed from fetal bovine dermis. It was cleared for marketing by the FDA through the 510(k) process for partial- and full-thickness wounds; diabetic, pressure, and venous stasis ulcers; surgical wounds; and tunneling, draining, and traumatic wounds.

SurgiMend® PRS (TEI Biosciences; a subsidiary of Integra Life Sciences) is a xenogeneic ADM processed from fetal and neonatal bovine dermis.

Strattice™ Reconstructive Tissue Matrix (LifeCell Corp.) is a xenogenic non-cross-linked porcine-derived ADM. There are pliable and firm versions, which are stored at room temperature and come fully hydrated.

FDA Product codes: KGN, FTL, FTM.

Living Cell Therapy

Apligraf® (Organogenesis) is a bilayered living cell therapy composed of an epidermal layer of living human keratinocytes and a dermal layer of living human fibroblasts. Apligraf® is supplied as needed, in 1 size, with a shelf-life of 10 days. In 1998, it was approved by the FDA for use in conjunction with compression therapy for the treatment of noninfected, partial- and full-thickness skin ulcers due to venous insufficiency and in 2001 for full-thickness neuropathic diabetic lower-extremity ulcers nonresponsive to standard wound therapy.

Dermagraft® (Organogenesis) is composed of cryopreserved human-derived fibroblasts and collagen derived from newborn human foreskin and cultured on a bioabsorbable polyglactin mesh scaffold. Dermagraft has been approved by the FDA for repair of diabetic foot ulcers.

Epicel® (Genzyme Biosurgery) is an epithelial autograft composed of a patient's own keratinocytes cultured ex vivo and is FDA-approved under a humanitarian device exemption (HDE) for the treatment of deep dermal or full-thickness burns comprising a total body surface area of 30% or more. It may be used in conjunction with split-thickness autografts or alone in patients for whom split-thickness autografts may not be an option due to the severity and extent of their burns.

OrCel™ (Forticell Bioscience; formerly Composite Cultured Skin) is an absorbable allogeneic bilayered cellular matrix, made of bovine collagen, in which human dermal cells have been cultured. It was approved by the FDA premarket approval for healing donor site wounds in burn victims and under a HDE for use in patients with recessive dystrophic epidermolysis bullosa undergoing hand reconstruction surgery to close and heal wounds created by the surgery, including those at donor sites.

FDA product codes: FTM, PFC, OCE, ODS.

Autologous Cell Harvesting Device

Recell® (Avita Medical) was initially approved by the FDA in September 2018 under the PMA process (PMA BP170122). It is an autologous cell harvesting device indicated for the treatment of acute partial-thickness thermal burn wound when used by an appropriately-licensed healthcare professional at the patient's point of care to prepare autologous RES Regenerative Epidermal Suspension. The initial indication was for use in patients 18 years of age and older in combination with meshed autografting. Subsequently, indications were expanded to include direct application to acute partial-thickness thermal burn wounds in patients 18 years of age and older or application in combination with meshed autografting for acute full-thickness thermal burn wounds in pediatric as well as adult patients and full-thickness skin defects after traumatic avulsion (e.g., degloving) or surgical excision (e.g., necrotizing tissue infection) or resection (e.g., skin cancer) in patients 15 years of age and older.

FDA product code: QCZ.

Biosynthetic Products

Biobrane®/Biobrane-L (Smith & Nephew) is a biosynthetic wound dressing constructed of a silicon film with a nylon fabric partially imbedded into the film. The fabric creates a complex 3-dimensional structure of tri-filament thread, which chemically binds collagen. Blood/sera clot in the nylon matrix, adhering the dressing to the wound until epithelialization occurs.

Integra® Dermal Regeneration Template (also marketed as Omnigraft Dermal Regeneration Matrix; Integra LifeSciences) is a bovine, collagen/glycosaminoglycan dermal replacement covered by a silicone temporary epidermal substitute. It was approved by the FDA for use in the post-excisional treatment of life-threatening full-thickness or deep partial-thickness thermal injury where sufficient autograft is not available at the time of excision or not desirable because of the physiologic condition of the patient and for certain diabetic foot ulcers. Integra® Matrix Wound Dressing and Integra® Meshed Bilayer Wound Matrix are substantially equivalent skin substitutes and were cleared for marketing by the FDA through the 510(k) process for other indications. Integra® Bilayer Matrix Wound Dressing (Integra LifeSciences) is designed to be used in conjunction with negative pressure wound therapy. The meshed bilayer provides a flexible wound covering and allows drainage of wound exudate.

TransCyte™ (Advanced Tissue Sciences) consists of human dermal fibroblasts grown on nylon mesh, combined with a synthetic epidermal layer and was approved by the FDA in 1997. TransCyte is intended as a temporary covering over burns until autografting is possible. It can also be used as a temporary covering for some burn wounds that heal without autografting.

FDA product codes: FRO, MDD, MGR.

Synthetic Products

Suprathel® (PolyMedics Innovations) is a synthetic copolymer membrane fabricated from a tripolymer of polylactide, trimethylene carbonate, and *s*-caprolactone. It is used to provide temporary coverage of superficial dermal burns and wounds. Suprathel® is covered with gauze and a dressing that is left in place until the wound has healed.

Rationale

This medical policy was created in 2019 and has been updated regularly with searches of the PubMed database. The most recent literature update was performed through November 13, 2023.

Medical policies assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

There is no standard definition of "skin substitute". Products reviewed in the following sections include products that do not require U.S. Food and Drug Administration (FDA) approval or clearance as well as a number of products cleared through the 510(k) pathway with a variety of FDA product codes. The FDA product codes that include these products are not limited to skin

substitute products and may include other indications not related to wound healing or wound care.

Breast Reconstruction

Clinical Context and Therapy Purpose

A variety of breast reconstruction techniques are used postmastectomy, including implant-based (immediate or delayed following use of a tissue expander) and those using autologous tissue flaps. Some of these techniques have been used with acellular dermal matrix (ADM) to provide additional support or tissue coverage. The purpose of bioengineered soft tissue substitutes in individuals who are undergoing breast reconstruction is to provide a treatment option that is an alternative to or an improvement on breast reconstruction without use of a biological or biosynthetic matrix.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals who are undergoing breast reconstruction, typically following mastectomy.

Interventions

The therapy being considered is bioengineered soft tissue substitutes as a biological matrix that is used to facilitate one-stage tissue expander reconstruction. As noted in the regulatory status section, the U.S. Food and Drug Administration (FDA) has not cleared or approved any surgical mesh device (synthetic, animal collagen-derived, or human collagen-derived) for use in breast surgery. In October 2021, an FDA advisory panel on general and plastic surgery voted against recommending FDA approval of the SurgiMend mesh for the specific indication of breast reconstruction. The advisory panel concluded that the benefits of using the device did not outweigh the risks. (4)

Comparators

The following therapies are currently being used to make decisions about soft tissue substitutes or biological matrices: 2-stage tissue expander reconstruction without a biological matrix.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity. Specific outcomes are the time to permanent implant, pain during and after the procedure, and adverse events including seroma, infection, and necrosis rates, rates of capsular contracture, and malposition of implants. Short-term outcomes would be measured within 3 months with longer-term outcomes apparent by 2 years.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials, with preference for RCTs* were sought.

- In the absence of such trials, comparative observational studies, with preference for prospective studies were sought.
- To assess longer-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Within each category of study design, larger sample size studies and longer duration studies were preferred.
- Studies with duplicative or overlapping populations were excluded.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

The literature on ADM for breast reconstruction consists primarily of retrospective, uncontrolled series and systematic reviews of these studies.

A 2013 study used data from the American College of Surgeon’s National Surgical Quality Improvement Program to compare ADM-assisted tissue expander breast reconstruction (n=1717) to submuscular tissue expander breast reconstruction (n=7442) after mastectomy. (5) Complication rates did not differ significantly between the ADM-assisted (5.5%) and the submuscular tissue expander groups (5.3%; p=0.68). Rates of reconstruction-related complications, major complications, and 30-day reoperation did not differ significantly between cohorts.

Systematic Reviews

A meta-analysis by Lee and Mun (2016) included 23 studies (total N=6199 cases) on implant-based breast reconstruction that were published between February 2011 and December 2014. (6) The analysis included an RCT and 3 prospective comparative cohort studies; the remainder was retrospective comparative cohort studies. Use of ADM did not affect the total complication rate (see Table 1). ADM significantly increased the risk of major infection, seroma, and flap necrosis, but reduced risks of capsular contracture and implant malposition. Use of ADM allowed for significantly greater intraoperative expansion (mean difference [MD], 79.63; 95% confidence interval [CI], 41.99 to 117.26; p<0.001) and percentage of intraoperative filling (MD=13.30; 95% CI, 9.95 to 16.65; p<0.001), and reduced the frequency of injections to complete expansion (MD = -1.56; 95% CI, -2.77 to -0.35; p=0.01).

Table 1. Meta-Analysis of Breast Reconstruction Outcomes With and Without ADM

Outcome Measure	Relative Risk	95% Confidence Interval	p
Infection	1.42	1.02 to 1.99	0.04
Seroma	1.41	1.12 to 1.78	0.004
Mastectomy flap necrosis	1.44	1.11 to 1.87	0.006
Unplanned return to the operating room	1.09	0.63 to 1.90	NS
Implant loss	1.00	0.68 to 1.48	NS

Total complications	1.08	0.87 to 1.34	NS
Capsular contracture	0.26	0.15 to 0.47	<0.001
Implant malposition	0.21	0.07 to 0.59	0.003

Adapted from Lee and Mun (2016). (6)

ADM: acellular dermal matrix; NS: not significant.

AlloDerm

Randomized Controlled Trials

McCarthy et al. (2012) reported on a multicenter, blinded RCT of AlloDerm in 2-stage expander/implant reconstruction. (7) Seventy patients were randomized to AlloDerm ADM-assisted tissue expander/implant reconstruction or to submuscular tissue expander/implant placement. The trial was adequately powered to detect clinically significant differences in immediate postoperative pain but underpowered to detect the secondary end point of pain during tissue expansion. There were no significant differences between the groups in the primary outcomes of immediate postoperative pain (54.6 AlloDerm vs 42.8 controls on a 100-point visual analog scale) or pain during the expansion phase (17.0 AlloDerm vs 4.6 controls), or in the secondary outcome of rate of tissue expansion (91 days AlloDerm vs 108 days controls) and patient-reported physical well-being. There was no significant difference in adverse events, although the total number of adverse events was small.

Comparison Between Products

AlloDerm vs AlloMax

Hinchcliff et al. (2017) conducted an RCT that compared AlloDerm with AlloMax (n=15 each) for implant-based breast reconstruction. (8) Complications were assessed 7, 14, and 30 days postoperatively and biopsies of the ADMs were taken during implant exchange. Vessel density in the AlloMax biopsies was higher than in the AlloDerm biopsies. Complications were reported in 26.1% of AlloMax cases and 8.0% of AlloDerm cases; these complication rates did not differ statistically with the 30 patients in this trial.

AlloDerm vs DermaMatrix

Mendenhall et al. (2017) conducted an RCT that compared AlloDerm with DermaMatrix in 111 patients (173 breasts). (9) There were no significant differences in overall rates of complications (AlloDerm, 15.4%; DermaMatrix, 18.3%; p=0.8) or implant loss (AlloDerm, 2.2%; DermaMatrix, 3.7%; p=0.5) between the 2 ADMs at 3 months postoperative. (9) There were no statistically significant differences in the overall complication rates (6% vs. 13%; p=.3), severity of complications, or patient satisfaction between the AlloDerm and DermaMatrix groups at 2 years after definitive reconstruction. (10)

Strattice

Dikmans et al. (2017) reported on early safety outcomes from an open-label multicenter RCT that compared porcine ADM-assisted 1-stage expansion with 2-stage implant-based breast reconstruction (see Table 2). (11) One-stage breast reconstruction with porcine ADM was associated with a higher risk of surgical complications, reoperation, and with removal of implant, ADM, or both (see Table 3). The trial was stopped early due to safety concerns, but it

cannot be determined from this study design whether the increase in complications was due to the use of the xenogenic ADM or to the comparison between 1-stage and 2-stage reconstruction.

Table 2. Summary of Key RCT Characteristics

Author	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Dikmans et al. (2017) (11)	EU	8	2013-2015	Women intending to undergo skin-sparing mastectomy and immediate IBBR	59 patients (91 breasts) undergoing 1-stage IBBR with ADM	62 women (92 breasts) undergoing 20-stage IBBR

ADM: Acellular dermal matrix; EU: European Union; IBBR: implant-based breast reconstruction; RCT: randomized controlled trial.

Table 3. Summary of Key RCT Outcomes

Study	Surgical Outcomes	Severe Adverse Effects	Reoperation	Removal of Implant, ADM, or Both
Dikmans et al. (2017) (11)				
1-stage with ADM, n (%)	27 (46)	26 (29)	22 (37)	24 (26)
2-stage with ADM, n (%)	11 (18)	5 (5)	9 (15)	4 (5)
OR (95% CI)	3.81 (2.67 to 5.43)		3.38 (2.10 to 5.45)	8.80 (8.24 to 9.40)
p	<0.001		<0.001	<0.001

ADM: acellular dermal matrix; CI: confidence interval; OR: odds ratio; RCT: randomized controlled trial.

Section Summary: Breast Reconstruction

Results of a systematic review found no difference in overall complication rates between ADM allograft and standard procedures for breast reconstruction. Although reconstructions with ADM have been reported to have higher seroma, infection, and necrosis rates than reconstructions without ADM, rates of capsular contracture and malposition of implants may be reduced. Thus, in cases where there is limited tissue coverage, the available studies may be considered sufficient to permit informed decision-making about risks and benefits of using allogeneic ADM for breast reconstruction.

Tendon Repair

Clinical Context and Therapy Purpose

The purpose of bio-engineered soft tissue substitutes in individuals who are undergoing tendon repair is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals undergoing tendon repair.

Interventions

The therapy being considered is bioengineered soft-tissue substitutes.

Comparators

The following therapies are currently being used to make decisions about tendon repair: tendon repair without bioengineered soft-tissue substitutes.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity. Short-term outcomes would be measured within 3 months with longer-term outcomes apparent by 2 years.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials, with preference for RCTs* were sought.
- In the absence of such trials, comparative observational studies, with preference for prospective studies were sought.
- To assess longer-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Within each category of study design, larger sample size studies and longer duration studies were preferred.
- Studies with duplicative or overlapping populations were excluded.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

GraftJacket

Barber et al. (2012) reported an industry-sponsored multicenter RCT of augmentation with GraftJacket human ADM for arthroscopic repair of large (>3 cm) rotator cuff tears involving 2 tendons. (12) Twenty-two patients were randomized to GraftJacket augmentation and 20 patients to no augmentation. At a mean follow-up of 24 months (range, 12-38 months), the American Shoulder and Elbow Surgeons score improved from 48.5 to 98.9 in the GraftJacket group and from 46.0 to 94.8 in the control group (p=0.035). The Constant score improved from 41 to 91.9 in the GraftJacket group and from 45.8 to 85.3 in the control group (p=0.008). The University of California, Los Angeles score did not differ significantly between groups. Gadolinium-enhanced magnetic resonance imaging (MRI) scans showed intact cuffs in 85% of repairs in the GraftJacket group and 40% of repairs in the control group. However, no correlation was found between MRI findings and clinical outcomes. Rotator cuff re-tears occurred in 3 (14%) patients in the GraftJacket group and 9 (45%) patients in the control group.

Rashid et al. (2020) reported disruption of the native extracellular matrix with either GraftJacket or Permacol (porcine acellular dermis) as a patch overlay for rotator cuff repair in a small controlled study with 13 patients. (13) The disruption was greater in the Permacol group and there was an immune response in 1 of 3 patients following use of the xenograft.

Section Summary: Tendon Repair

One small RCT was identified that found improved outcomes with GraftJacket ADM allograft for rotator cuff repair. Although results of this trial were promising, additional study with a larger number of patients is needed to evaluate consistency of findings and determine the effects of this technology with greater certainty.

Surgical Repair of Hernias or Parastomal Reinforcement

Clinical Context and Therapy Purpose

The purpose of bio-engineered soft tissue substitutes in individuals who are undergoing surgical repair of hernias or require parastomal reinforcement is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals undergoing surgical repair of hernias or require parastomal reinforcement.

Interventions

The therapy being considered is bioengineered matrix support.

Comparators

The following therapies are currently being used for surgical repair of hernias or parastomal reinforcement: synthetic mesh.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity. Specific outcomes are surgical site occurrence of postoperative infection, seroma/hematoma, pain, bulging, dehiscence, fistula, or mechanical failure. Short-term outcomes would be measured within 3 months with longer-term outcomes apparent by 2 years.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials, with preference for RCTs* were sought.
- In the absence of such trials, comparative observational studies, with preference for prospective studies were sought.
- To assess longer-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.

- Within each category of study design, larger sample size studies and longer duration studies were preferred.
- Studies with duplicative or overlapping populations were excluded.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials.

A 2013 systematic review evaluated the clinical effectiveness of acellular collagen-based scaffolds for the repair of incisional hernias. (14) The bioprosthetic materials could be harvested from bovine pericardium, human cadaveric dermis, porcine small intestine mucosa, porcine dermal collagen, or bovine dermal collagen. Products included in the search were Surgisis, Tutomesh, Veritas, AlloDerm, FlexHD, AlloMax, CollaMend, Permacol, Strattice, FortaGen, ACell, DermaMatrix, XenMatrix, and SurgiMend. Sixty publications with 1,212 repairs were identified and included in the review, although meta-analysis could not be performed. There were 4 level III studies (2 AlloDerm, 2 Permacol); the remainder was level IV or V. The largest number of publications were on AlloDerm (n=27) and Permacol (n=18). No publications on incisional hernia repair were identified for AlloMax, FortaGen, DermaMatrix, or ACell. The overall incidence of a surgical site occurrence (e.g., postoperative infection, seroma/hematoma, pain, bulging, dehiscence, fistula, mechanical failure) was 82.6% for porcine small intestine mucosa, 50.7% for xenogenic dermis, 48.3% for human dermis, and 6.3% for xenogenic pericardium. No comparative data were identified that could establish superiority to permanent synthetic meshes.

AlloDerm as an Overlay

Espinosa-de-los-Monteros et al. (2007) retrospectively reviewed 39 abdominal wall reconstructions with AlloDerm performed in 37 patients and compared them with 39 randomly selected cases. (15) They reported a significant decrease in recurrence rates when human cadaveric acellular dermis was added as an overlay to primary closure plus rectus muscle advancement and imbrication in patients with medium-sized hernias. However, no differences were observed when adding human cadaveric acellular dermis as an overlay to patients with large-size hernias treated with underlay mesh.

Comparisons Between Products

AlloDerm vs Surgisis Gold

Gupta et al. (2006) compared the efficacy and complications associated with use of AlloDerm and Surgisis bioactive mesh in 74 patients who underwent ventral hernia repair. (16) The first 41 procedures were performed using Surgisis Gold 8-ply mesh formed from porcine small intestine submucosa, and the remaining 33 patients had ventral hernia repair with AlloDerm. Patients were seen 7 to 10 days after discharge from the hospital and at 6 weeks. Any signs of wound infection, diastasis, hernia recurrence, changes in bowel habits, and seroma formation were evaluated. The use of the AlloDerm mesh resulted in 8 (24%) hernia recurrences. Fifteen (45%) of the AlloDerm patients developed a diastasis or bulging at the repair site. Seroma formation was only a problem in 2 patients.

AlloDerm vs FlexHD

A 2013 study compared AlloDerm with FlexHD for complicated hernia surgery. (17) From 2005 to 2007, AlloDerm was used to repair large (>200 cm²) symptomatic complicated ventral hernias that resulted from trauma or emergency surgery (n=55). From 2008 to 2010, FlexHD was used to repair large, complicated ventral hernias in patients meeting the same criteria (n=40). The 2 groups were comparable at baseline. At 1-year follow-up, all AlloDerm patients were diagnosed with hernia recurrence (abdominal laxity, functional recurrence, true recurrence) requiring a second repair. Eleven (31%) patients in the FlexHD group required a second repair. This comparative study is limited by the use of nonconcurrent comparisons, which is prone to selection bias and does not control for temporal trends in outcomes.

FlexHD vs Strattice

Roth et al. (2017) reported on a prospective study assessing clinical and quality of life outcomes following complex hernia repair with a human (FlexHD) or porcine (Strattice) ADM. (18) The study was funded by the Musculoskeletal Transplant Foundation, which prepares and supplies FlexHD. Patients were enrolled if they had a hernia at least 6 cm in the transverse dimension, active or prior infection of the abdominal wall, and/or enterocutaneous fistula requiring mesh removal. Eighteen (51%) of the 35 patients had undergone a previous hernia repair. After abdominal wall repair with the ADM, 20 (57%) patients had a surgical site occurrence, and nearly one-third had hospital readmission. The type of biologic material did not impact hernia outcomes. There was no comparison with synthetic mesh in this study, limiting interpretation.

Strattice vs Synthetic Mesh

Bellows et al. (2014) reported early results of an industry-sponsored multicenter RCT that compared Strattice (non-cross-linked porcine ADM, n=84) with a standard synthetic mesh (n=88) for the repair of inguinal hernias. (19) The trial was designed by the surgeons and was patient-and assessor-blinded to reduce risk of bias. Blinding continued through 2 years of follow-up. The primary outcome was resumption of activities of daily living at 1 year. Secondary outcomes included complications, recurrences, or chronic pain (i.e., pain that did not disappear by 3 months postsurgery). At 3-month follow-up, there were no significant differences in either the occurrence or type of wound events (relative risk, 0.98; 95% CI, 0.52 to 1.86). Pain was reduced from 1 to 3 days postoperative in the group treated with Strattice, but at 3-month follow-up pain scores did not differ significantly between groups.

Strattice vs No Reinforcement

Also, in 2014, the Parastomal Reinforcement with Strattice (PRISM) Study Group reported a multicenter, double-blinded, randomized trial of Strattice for parastomal reinforcement in patients undergoing surgery for permanent abdominal wall ostomies. (20) Patients were randomized to standard stoma construction with no reinforcement (n=58) or stoma construction with Strattice as parastomal reinforcement (n=55). At 24-month follow-up (n=75), the incidence of parastomal hernias was similar for the 2 groups (13.2% of controls, 12.2% of study group).

Adverse Events

Permacol (porcine acellular dermal matrix) was reported in a case series of 13 patients to result in recurrent intestinal fistulation and intestinal failure when used for abdominal reconstructive surgery. (21)

Section Summary: Surgical Repair of Hernias or Parastomal Reinforcement

Current evidence does not support a benefit of ADMs in hernia repair or prevention of parastomal hernia. Additional RCTs are needed to compare biologic mesh with synthetic mesh and to determine if there is a patient population that would benefit from these products.

Diabetic Lower-Extremity Ulcers

Clinical Context and Therapy Purpose

The purpose of bio-engineered soft tissue substitutes in individuals who have diabetic lower extremity ulcers is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with diabetic lower extremity ulcers.

Interventions

The therapy being considered is bioengineered skin substitutes.

Comparators

The following therapies are currently being used: standard wound care which involves regular debridement and moist wound covering.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, and QOL.

The primary endpoints of interest for trials of wound closure are as follows, consistent with guidance from the FDA for industry in developing products for treatment of chronic cutaneous ulcer and burn wounds:

- Incidence of complete wound closure.
- Time to complete wound closure (reflecting accelerated wound closure).
- Incidence of complete wound closure following surgical wound closure.
- Pain control.

Time to wound closure can be measured at 6 months with longer-term outcomes apparent by 1 year. More complex wounds may require more than 6 months to heal.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials, with preference for RCTs* were sought.
- In the absence of such trials, comparative observational studies, with preference for prospective studies were sought.
- To assess longer-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Within each category of study design, larger sample size studies and longer duration studies were preferred.
- Studies with duplicative or overlapping populations were excluded.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials

Systematic Reviews

A 2016 Cochrane review evaluated skin substitutes for the treatment of diabetic foot ulcers. (22) Seventeen trials (total N=1655 participants) were included in the meta-analysis. Most trials identified were industry-sponsored, and an asymmetric funnel plot indicated publication bias. Pooled results of published trials found that skin substitutes increased the likelihood of achieving complete ulcer closure compared with standard of care (SOC) alone (relative risk, 1.55; 95% CI, 1.30 to 1.85). Use of skin substitutes also led to a statistically significant reduction in amputations (relative risk, 0.43; 95% CI, 0.23 to 0.81), although the absolute risk difference was small. Analysis by individual products found a statistically significant benefit on ulcer closure for Apligraf, EpiFix, and Hyalograft-3D. The products that did not show a statistically significant benefit for ulcer closure were Dermagraft, GraftJacket, Kaloderm, and OrCel.

Apligraf, Dermagraft, AlloPatch, Integra Dermal Regeneration Template, or Integra Flowable Wound Matrix, mVASC, or TheraSkin

Apligraf

Veves et al. (2001) reported on a randomized prospective trial on the effectiveness of Apligraf (previously called Graftskin), a living skin equivalent, in treating noninfected nonischemic chronic plantar diabetic foot ulcers. (23) The trial involved 24 centers in the United States; 208 patients were randomized to ulcer treatment with Apligraf (112 patients) or saline-moistened gauze (96 patients, control group). Standard state-of-the-art adjunctive therapy, including extensive surgical débridement and adequate foot off-loading, was provided in both groups. Apligraf was applied at the beginning of the study and weekly thereafter for a maximum of 4 weeks (maximum of 5 applications) or earlier if complete healing occurred. At the 12-week follow-up visit, 63 (56%) Apligraf-treated patients achieved complete wound healing compared with 36 (38%) in the control group ($p=0.004$). The Kaplan-Meier method median time to complete closure was 65 days for Apligraf, significantly lower than the 90 days observed in the control group ($p=0.003$). The rates of adverse reactions were similar between groups, except osteomyelitis and lower-limb amputations, both of which were less frequent in the Apligraf group. Trialists concluded that application of Apligraf for a maximum of 4 weeks resulted in higher healing rates than state-of-the-art treatment and was not associated with any significant adverse events.

Dermagraft

A 2003 pivotal multicenter FDA-regulated trial randomized 314 patients with chronic diabetic ulcers to Dermagraft (human-derived fibroblasts cultured on mesh) or control. (24) Over the 12-week study, patients received up to 8 applications of Dermagraft. All patients received pressure-reducing footwear and were encouraged to stay off their study foot as much as possible. At 12 weeks, the median percent wound closure for the Dermagraft group was 91% compared with 78% for the control group. Ulcers treated with Dermagraft closed significantly faster than ulcers treated with conventional therapy. No serious adverse events were attributed to Dermagraft. Ulcer infections developed in 10.4% of the Dermagraft patients compared with 17.9% of the control patients. Together, there was a lower rate of infection, cellulitis, and osteomyelitis in the Dermagraft-treated group (19% vs 32.5%). A 2015 retrospective analysis of the trial data found a significant reduction in amputation/bone resection rates with Dermagraft (5.5% vs 12.6%, $p=0.031$). (25) Of the 28 cases of amputation/bone resection, 27 were preceded by ulcer-related infection.

AlloPatch

AlloPatch Pliable human reticular acellular dermis was compared with SOC in an industry-sponsored multicenter trial by Zelen et al. (2017, 2018). (26, 27) The initial trial with 20 patients per group was extended to determine the percent healing at 6 weeks with 40 patients per group. Healing was evaluated by the site investigator and confirmed by an independent panel. At 6 weeks, 68% (27/40) of wounds treated using AlloPatch had healed compared with 15% (6/40) in the SOC-alone group ($p<0.001$). At 12 weeks, 80% (32/40) of patients in the AlloPatch group had healed compared to 30% (12/40) in the control group. Mean time to heal within 12 weeks was 38 days (95% CI: 29-47 days) for the HR-ADM group and 72 days (95% CI: 66-78 days) for the SOC group ($p < 0.001$).

Integra Omnigraft Dermal Regeneration Template or Integra Flowable Wound Matrix

Integra Dermal Regeneration Template is a biosynthetic skin substitute that is FDA-approved for life-threatening thermal injury. The FOUNDER (Foot Ulcer New Dermal Replacement) multicenter study (32 sites) assessed Integra Dermal Regeneration Template (marketed as Omnigraft) for chronic nonhealing diabetic foot ulcers under an FDA-regulated investigational device exemption. (28) A total of 307 patients with at least 1 chronic diabetic foot ulcer were randomized to treatment with the Integra Template or a control condition (sodium chloride gel 0.9%). Treatment was given for 16 weeks or until wound closure. There was a modest increase in wound closure with the Integra Template (51% vs 32%, $p=0.001$) and a shorter median time to closure (43 days vs 78 days, $p=0.001$). There was a strong correlation between investigator-assessed and computerized planimetry assessment of wound healing ($r=0.97$). Kaplan-Meier analysis showed the greatest difference between groups in wound closure up to 10 weeks, with diminishing differences after 10 weeks. Trial strengths included adequate power to detect an increase in wound healing of 18%, which was considered to be clinically significant, secondary outcomes of wound closure and time to wound closure by computerized planimetry, and ITT analysis.

Integra Flowable Wound Matrix is composed of a porous matrix of cross-linked bovine tendon collagen and glycosaminoglycan. It is supplied as a granular product that is mixed with saline. Campitiello et al. (2017) published an RCT that compared the flowable matrix with wet dressing in 46 patients who had Wagner grade 3 diabetic foot ulcers. (29) The ulcers had developed over 39 weeks. Complete healing at 6 weeks was achieved in significantly more patients in the Integra Flowable Wound Matrix group than in the control group, while the risk of rehospitalization and major amputation was reduced with Integra Flowable Wound Matrix (see Table 4).

Table 4. Probability of Wound Healing with IFWM vs SOC

Study	Complete Wound Healing	Rehospitalization	Major Amputation
Campitiello et al. (2017) (29)			
IFWM, n (%)	20 (86.95)	2 (6.69)	1 (4.34)
SOC, n (%)	12 (52.17)	10 (43.47)	7 (30.43)
RR (95% CI)	1.67 (1.09 to 2.54)	0.10 (0.01 to 0.72)	0.16 (0.02 to 1.17)
p	0.010	0.001	0.028

CI: confidence interval; IFWM Integra Flowable Wound Matrix; RR: relative risk; SOC: standard of care.

mVACS

Tables 5 and 6 summarize the trial characteristics and results for RCTs of mVASC. Tables 7 and 8 evaluate study limitations.

Gould et al. (2023) reported results of the HIFLO (Healing in Diabetic Foot Ulcers with Microvascular Tissue) Trial, a multicenter (6 US sites) RCT comparing weekly application of the processed microvascular tissue (PMVT) allograft, mVASC in addition to a standardized diabetic foot ulcer protocol versus standard wound care with a collagen alginate dressing control in 100 adults with Wagner Grade 1 and 2 diabetic foot ulcers of ≥ 4 weeks and < 52 weeks duration. (30) Wound and local peripheral neuropathy assessment were performed weekly. The primary outcome of the study was complete wound closure at 12 weeks. The investigator and a blinded physician made the initial determination of wound closure, followed by adjudication and confirmation by an independent, blinded panel of plastic surgeons. All participants who attended at least 1 treatment visit were included in the analysis. There was missing data for 15 participants at week 12 (3 in mVASC vs. 12 in control) and 14 of these were missing due to adverse events related to the wound. These were included in the primary analysis and counted as wound healing failures. The mean age of participants was 60 years, 90% of participants were White and 10% were Black, and 66% of participants were men. At randomization, the mean size of the wound area was 3.3 cm and the mean duration of the wound was 15 weeks. The proportion of participants with complete wound closure at week 12 was 74% (37/50) for mVASC versus 38% (19/50) for control ($p < .001$). Of the wounds that healed, the mean time to healing was also statistically significantly faster for the mVASC group (54 days; 95% CI, 46 to 61 vs 64 days; 95% CI, 57 to 72; $p = .009$). The 10-point Semmes-Weinstein monofilament (SWM) test of peripheral neuropathy also favored mVASC (118% vs. 11%; $p = .028$). No adverse events or serious adverse events related to the study treatment or the procedure were reported.

There were 11 adverse events (3, mVASC vs. 8, control) reported that were related to the wound.

Table 5. Randomized Controlled Trial of mVASC for Diabetic Foot Ulcers-Characteristics

Study	Countries	Sites	Dates	Participants	Inerventions	
					Active	Comparator
Gould 2023; HIFLO (30)	US	6	2017-2020	Adults with chronic Wagner Grade 1 or 2 DFU Mean age, 60 y 90% White 10% Black 66% Male Mean wound size 3.3 cm	mVASC + SOC (n=50)	SOC (n=50)

DFU: Diabetic Foot Ulcers; HIFLO: Healing in Diabetic Foot Ulcers with Microvascular Tissue; SOC: Standard of Care

Table 6. Randomized Controlled Trial of mVASC for Diabetic Foot Ulcers-Results

Study	Wounds Healed	Time to Heal	% Area Reduction	Adverse Events
Gould 2023; HIFLO (30)	at 12 weeks	by 12 weeks	at 12 weeks	--
N analyzed	100	56	100	100
mVACS	74% (37/50)	Mean, 54 d	76%	3
SOC	38% (19/50)	Mean, 64 d	24%	8
p-value	<.001	.009	.009	--

DFU: Diabetic Foot Ulcers; HIFLO: Healing in Diabetic Foot Ulcers with Microvascular Tissue; SOC: Standard of Care

Table 7. Randomized Controlled Trial of mVASC for Diabetic Foot Ulcers-Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Gould 2023; HIFLO (30)	4. Lack of racial and ethnic diversity				1. Follow-up not sufficient to determine ulcer recurrence

HIFLO: Healing in Diabetic Foot Ulcers with Microvascular Tissue

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^aPopulation key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^bIntervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5: Other.

^cComparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^dOutcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^eFollow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 8. Randomized Controlled Trial of mVASC for Diabetic Foot Ulcers-Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Gould 2023; HIFLO (30)			1. Registered retrospectively in European registry			3. Confidence intervals not reported

HIFLO: Healing in Diabetic Foot Ulcers with Microvascular Tissue

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^aAllocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^bBlinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^cSelective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^dData Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^ePower key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^fStatistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other

TheraSkin Versus Standard of Care

Tables 9 and 10 summarize the trial characteristics and results for RCTs of TheraSkin compared to SOC. Tables 11 and 12 evaluate study limitations.

Armstrong et al. (2022) reported results of an RCT including 100 adults with non-healing Wagner 1 diabetic foot ulcers comparing Theraskin (n=50) to SOC (n=50). (31) The index ulcer had to have been present for greater than 4 weeks and less than 1 year with a minimum size of

1.0 cm² and a maximum size of 25 cm². Standard of care included glucose monitoring, weekly debridement as appropriate, and an offloading device. The dressing in the SOC group was calcium alginate (Fibracol Plus). The primary outcome was the proportion of full-thickness wounds healed at 12 weeks. Wound healing was assessed initially by the investigator and confirmed by blinded adjudication panel. Wounds were closed when there was 100% re-epithelization and no drainage. The mean age of participants was 60 years; 53% of participants were male, 70% were White, and 15% were Black. The mean wound area at baseline was 4.1 cm². Participants who did not have healing of at least 50% by 6 weeks were allowed to seek alternative rescue wound care (TheraSkin, n=1; SOC, n=11). In addition, 3 participants in the TheraSkin group and 8 in the SOC group had worsening of the wound or an adverse event before week 12. All enrolled participants were included in analysis and missing data were imputed using last observation carried forward. The percent of participants with complete wound healing at week 12 was 76% (38/50) in the intervention group compared with 36% (18/50) in the SOC group (p<.01). The mean percent area reduction at 12 weeks was 77.8% in the TheraSkin group compared with 49.6% in the SOC group (p<.01). There were no statistically significant differences between groups in QOL or pain score measures.

Table 9. Randomized Controlled Trial of TheraSkin vs. SOC for Diabetic Foot Ulcers-Characteristics

Study	Countries	Sites	Dates	Participants	Inerventions	
					Active	Comparator
Armstrong (2022); NCT04040426 (31)	US	5	2019-2021	Adults with non-healing Wagner Grade 1 DFUs Mean wound size 4.1 cm ² Mean age, 60 y 53% Male 70% White 15% Black	TheraSkin (n=50)	SOC with calcium alginate dressing (n=50)

DFU: Diabetic Foot Ulcers; SOC: standard of care

Table 10. Randomized Controlled Trial of TheraSkin vs. SOC for Diabetic Foot Ulcers-Results

Study	Wounds Healed	Time to Heal	% Area Reduction	Adverse Events
Armstrong (2022); NCT04040426 (31)	at 12 weeks	by 12 weeks	at 12 weeks	--
N analyzed	100	100	100	100

TheraSkin	76% (38/50)	Mean, 47 days (95% CI, 39 to 55)	78% (SD=63)	2
SOC	36% (18/50)	Mean, 65 days (95% CI, 58 to 73)	50% (SD=98)	4
p-value	<.01	<.01	<.01	NR

CI: confidence interval; NR: not reported; SD: standard deviation; SOC: standard of care

Table 11. Randomized Controlled Trial of TheraSkin vs. SOC for Diabetic Foot Ulcers-Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Armstrong (2022); NCT04040426 (31)	4. Lack of racial and ethnic diversity				1. Follow-up not sufficient to determine ulcer recurrence

SOC: standard of care.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^aPopulation key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^bIntervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5: Other.

^cComparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^dOutcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^eFollow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 12. Randomized Controlled Trial of TheraSkin vs. SOC for Diabetic Foot Ulcers-Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reportin ^g _c	Data Completeness ^d	Power ^e	Statistical ^f
Armstrong (2022); NCT 04040426 (31)		1. Investigators not blinded		2. Missing data imputed by last observation carried forward; no sensitivity		

				analyses provided		
--	--	--	--	-------------------	--	--

SOC: standard of care.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^aAllocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^bBlinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^cSelective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^dData Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^ePower key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^fStatistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other

TheraSkin vs Dermagraft

Sanders et al. (2014) reported on a small (N=23) industry-funded randomized comparison of TheraSkin (cryopreserved human skin allograft with living fibroblasts and keratinocytes) and Dermagraft for diabetic foot ulcers. (32) Wound size at baseline ranged from 0.5 to 18.02 cm²; the average wound size was about 5 cm² and was similar for the 2 groups (p=0.51). Grafts were applied according to manufacturers' instructions over the first 12 weeks of the study until healing, with an average of 4.4 TheraSkin grafts (every 2 weeks) compared with 8.9 Dermagraft applications (every week). At week 12, complete wound healing was observed in 63.6% of ulcers treated with TheraSkin and 33.3% of ulcers treated with Dermagraft (p<0.049). At 20 weeks, complete wound healing was observed in 90.9% of the TheraSkin-treated ulcers compared with 66.7% of the Dermagraft group (p=0.428).

TheraSkin vs Apligraf

DiDomenico et al. (2011) compared TheraSkin with Apligraf for the treatment of diabetic foot ulcers in a small (N=29) RCT. (33) The risk of bias in this study is uncertain, because reporting did not include a description of power analysis, statistical analysis, method of randomization, or blinding. The percentage of wounds closed at 12 weeks was 41.3% in the Apligraf group and 66.7% in the TheraSkin group. Results at 20 weeks were not substantially changed from those at 12 weeks, with 47.1% of wounds closed in the Apligraf group and 66.7% closed in the TheraSkin group. The percentage healed in the Apligraf group was lower than expected based on prior studies. The average number of grafts applied was similar for both groups (1.53 for Apligraf, 1.38 for TheraSkin). The low number of dressing changes may have influenced results, with little change in the percentage of wounds closed between 12 and 20 weeks. An adequately powered trial with blinded evaluation of wound healing and a standard treatment regimen would permit greater certainty on the efficacy of this product.

Subsection Summary: Apligraf, Dermagraft, AlloPatch, Integra, mVASC or TheraSkin for Diabetic Lower-Extremity Ulcers

RCTs reporting complete wound healing outcomes with at least 12 weeks of follow-up have demonstrated the efficacy of Apligraf, Dermagraft, AlloPatch, Integra Dermal Regeneration Template, Integra Flowable Wound Matrix, mVASC and TheraSkin over SOC for the treatment of diabetic lower-extremity ulcers.

Bioengineered Skin Substitutes Other Than Apligraf, Dermagraft, AlloPatch, Integra, mVASC, or Theraskin

GraftJacket Regenerative Tissue Matrix

Brigido et al. (2004) reported a small (N=40) randomized pilot study comparing GraftJacket with conventional treatment for chronic nonhealing diabetic foot ulcers. (34) Control patients received conventional therapy with débridement, wound gel with gauze dressing, and off-loading. GraftJacket patients received surgical application of the scaffold using skin staples or sutures and moistened compressive dressing. A second graft application was necessary after the initial application for all patients in the GraftJacket group. Preliminary one-month results showed that, after a single treatment, ulcers treated with GraftJacket healed at a faster rate than conventional treatment. There were significantly greater decreases in wound length (51% vs 15%), width (50% vs 23%), area (73% vs 34%), and depth (89% vs 25%), respectively. With follow-up to four weeks, no data were reported on the proportion with complete closure or the mean time to heal. All grafts were incorporated into the host tissue.

Reyzelman et al. (2009) reported an industry-sponsored multicenter randomized study that compared a single application of GraftJacket with SOC in 86 patients with diabetic foot ulcers. (35) Eight patients, six in the study group and two in the control group, did not complete the trial. At 12 weeks, complete healing was observed in 69.6% of the GraftJacket group and 46.2% of controls. After adjusting for ulcer size at presentation, a statistically significant difference in nonhealing rate was calculated, with odds of healing 2.0 times higher in the study group. Mean healing time was 5.7 weeks for the GraftJacket group vs 6.8 weeks for the control group. The authors did not report whether this difference was statistically significant. Median time to healing was 4.5 weeks for GraftJacket (range, 1-12 weeks) and 7.0 weeks for control (range, 2-12 weeks). Kaplan-Meier method survivorship analysis for time to complete healing at 12 weeks showed a significantly lower nonhealing rate for the study group (30.4%) than for the control group (53.9%). The authors commented that a single application of GraftJacket, as used in this study, was often sufficient for complete healing.

Reyzelman and Bazarov (2015) (36) reported an industry-sponsored meta-analysis of GraftJacket for diabetic foot ulcers that included the 2 studies described above and a third RCT by Brigido (2006) (37) with 28 patients (total N=154 patients). The time to heal was estimated for the Brigido (2004) study, based on the average wound reduction per week. The estimated difference in time to heal was considerably larger for Brigido's 2004 study (-4.30 weeks) than for the other 2 studies that measured the difference in time to heal (-1.58 weeks and -1.10 weeks). Analysis of the proportion of wounds that healed included Brigido (2006) and

Reyzelman et al. (2009). The odds ratio in the smaller study by Brigido (2006) was considerably larger, with a lack of precision in the estimate (odds ratio, 15.0; 95% CI, 2.26 to 99.64), and the combined odds (3.75; 95% CI, 1.72 to 8.19) was not significant when analyzed using a random-effects model. Potential sources of bias, noted by Reyzelman and Bazarov (2015), included publication and reporting biases, study selection biases, incomplete data selection, post hoc manipulation of data, and subjective choice of analytic methods. Overall, results of these studies do not provide convincing evidence that GraftJacket is more effective than SOC for healing diabetic foot ulcers.

DermACELL vs GraftJacket Regenerative Tissue Matrix or SOC

DermACELL and GraftJacket are both composed of human ADM. Walters et al. (2016) reported on a multicenter randomized comparison of DermACELL, GraftJacket, or SOC (2:1:2 ratio) in 168 patients with diabetic foot ulcers. (38) The study was sponsored by LifeNet Health, a nonprofit organ procurement association and processor for DermACELL. At 16 weeks, the proportion of completely healed ulcers was 67.9% for DermACELL, 47.8% for GraftJacket, and 48.1% for SOC. The 20% difference in completely healed ulcers was statistically significant for DermACELL vs SOC (p=0.039). The mean time to complete wound closure did not differ significantly for DermACELL (8.6 weeks), GraftJacket (8.6 weeks), and SOC (8.7 weeks).

A second report from this study was published in 2017. (39) This analysis compared DermACELL with SOC and did not include the GraftJacket arm. The authors reported that either 1 or 2 applications DermACELL led to a greater proportion of wounds healed compared with SOC in per protocol analysis (see Table 13), but there was no significant difference between DermACELL (1 or 2 applications) and SOC when analyzed by intention-to-treat (ITT). For the group of patients who received only a single application, the percentage of patients who achieved complete wound healing was significantly higher than SOC at 16 and 24 weeks, but not at 12 weeks. Although reported as ITT analysis, results were analyzed only for the group who received a single application of DermACELL. This would not typically be considered ITT.

Table 13. Probability of Wound Healing in Per Protocol Analysis of DermACELL vs SOC

Study	Single Application			1 or 2 Applications		
	% With Wound Healing at 12 Wk	% With Wound Healing at 16 Wk	% With Wound Healing at 24 Wk	% With Wound Healing at 12 Wk	% With Wound Healing at 16 Wk	% With Wound Healing at 24 Wk
Cazzell et al. (2017) (39)						
DermACELL, %	65.0%	82.5%	89.7%	NR	67.9%	83.7%
SOC, %	41.1%	48.1%	67.3%	NR	48.1%	67.3%
HR (95% CI)	1.97 (1.1 to 3.5)	2.40 (1.4 to 4.1)	2.11 (1.3 to 3.5)	--	1.72 (1.04 to 2.83)	1.55 (0.98 to 2.44)
p	0.012	<0.001	<0.001	NS	0.028	0.049

CI: confidence interval; HR: hazard ratio; NR: not reported; NS: not significant; SOC: standard of care; Wk: week.

Cytal (MatriStem) vs Dermagraft

Frykberg et al. (2017) reported a prespecified interim analysis of an industry-funded multicenter noninferiority trial of Cytal (a porcine urinary bladder-derived extracellular matrix) vs Dermagraft in 56 patients with diabetic foot ulcers. (40) The mean duration of ulcers before treatment was 263 days (range, 30-1095 days). The primary outcome was the percent wound closure with up to 8 weeks of treatment using blinded evaluation of photographs. Intention-to-treat (ITT) analysis found complete wound closure in 5 (18.5%) wounds treated with Cytal compared with 2 (6.9%) wounds treated with Dermagraft (p=NS [not significant]). Quality of life, measured by the Diabetic Foot Ulcer Scale, improved from 181.56 to 151.11 in the Cytal group and from 184.46 to 195.73 in the Dermagraft group (p=0.074). It should be noted that this scale is a subjective measure and patients were not blinded to treatment. Power analysis indicated that 92 patients would be required; further recruitment is ongoing for completion of the study.

PriMatrix

Lantis et al. (2021) reported on a multicenter RCT comparing PriMatrix plus standard of care to PriMatrix alone in 226 patients with diabetic foot ulcers (Tables 14 and 15). (41)

Study subjects underwent a 2-week run-in period of SOC treatment and were excluded if they had a wound reduction of 30% or more. Patients randomized to the SOC group received weekly treatment at the study site identical to the SOC treatment applied during the screening period. In addition, control group patients performed daily dressing changes, which consisted of wound cleaning, application of saline gel and secondary dressings. The primary endpoint was the percentage of subjects with complete wound closure, defined as 100% re-epithelialization without drainage during the 12-week treatment phase.

Significantly more patients in the PriMatrix group experienced complete wound closure at 12 weeks (45.6% vs 27.9%; p=.008). It is unclear if this difference (17.7%) is clinically significant; the study was powered to detect a 20% difference between groups. The time to complete healing did not differ between groups for the wounds that healed. Major study limitations include lack of blinding, limited generalizability, and insufficient duration of follow-up to assess wound recurrence (Tables 16 and 17).

Table 14. Randomized Controlled Trial of PriMatrix for Diabetic Foot Ulcers-Characteristics

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Lantis et al. (2021) (41) NCT03010319	US	21	2019-2020	Diabetic foot ulcer for a minimum of 2 weeks, adequate vascular perfusion to the affected extremity	PriMatrix plus standard of care n = 103	Standard of care n = 104

US: United States.

Table 15. Randomized Control Trial of PriMatrix for Diabetic Foot Ulcers-Results

Study	Wound Healed at 12 weeks	Median Time to Heal, days (range)	AEs
Lantis et al. (2021) (41) NCT03010319			
Number analyzed	207	76	226
Primatrix	47/103 (45.6%)	43 (22 to 93)	Any AE: 44.8%
Standard Care	29/104 (27.9%)	57 (16 to 88)	Any AE: 46.4%
Treatment Effect	HR 2.02 (95% CI 1.3 to 3.2)		
p	.008	.362	

AE: adverse events; CI: confidence interval; HR: hazard ratio

Table 16. Randomized Controlled Trial of PriMatrix for Diabetic Foot Ulcers-Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Lantis et al. (2021) (41) NCT03010319	4. Race and ethnicity of the study population was not reported and is not included in the demographics table.		3. Standard of care patients received additional dressing changes at home, which could have potentially exposed the wound to unknown factors.		1. 4-week follow-up not sufficient to determine ulcer recurrence.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5: Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 17. Randomized Controlled Trial of PriMatrix for Diabetic Foot Ulcers-Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Lantis et al. (2021) (41) NCT03010319	3. Allocation concealment not described.	1. Patients and investigator not blinded		1. 24 subjects from the treatment group and 22 from the control group discontinued from each arm prior to meeting the protocol-defined primary endpoint and were counted as treatment failures. 207 of 226 randomized were included in primary analysis (91.6%)		3. Confidence intervals not reported

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other

Oasis Wound Matrix vs Regranex Gel

Niezgoda et al. (2005) compared healing rates at 12 weeks for full-thickness diabetic foot ulcers treated with OASIS Wound Matrix (a porcine acellular wound care product) to Regranex Gel.

(42) This industry-sponsored, multicenter RCT was conducted at 9 outpatient wound care clinics and involved 73 patients with at least 1 diabetic foot ulcer. Patients were randomized to receive either Oasis Wound Matrix (n=37) or Regranex Gel (n=36) and a secondary dressing. Wounds were cleansed and débrided, if needed, at a weekly visit. The maximum treatment period for each patient was 12 weeks. After 12 weeks, 18 (49%) Oasis-treated patients had complete wound closure compared with 10 (28%) Regranex-treated patients. Oasis treatment met the noninferiority margin but did not demonstrate that healing in the Oasis group was statistically superior ($p=0.055$). Post hoc subgroup analysis showed no significant difference in incidence of healing in patients with type 1 diabetes (33% vs 25%) but showed a significant improvement in patients with type 2 diabetes (63% vs 29%). There was also an increased healing of plantar ulcers in the Oasis group (52% vs 14%). These post hoc findings are considered hypothesis-generating. Additional study with a larger number of subjects is needed to compare the effect of Oasis treatment to current SOC.

Autologous Grafting on HYAFF Scaffolds

Uccioli et al. (2011) reported a multicenter RCT of cultured expanded fibroblasts and keratinocytes grown on a HYAFF scaffold (benzyl ester of hyaluronic acid) compared with paraffin gauze for difficult diabetic foot ulcers. (43) A total of 180 patients were randomized. At 12 weeks, complete ulcer healing was similar for the two groups (24% treated vs 21% controls). At 20 weeks, complete ulcer healing was achieved in a similar proportion of the treatment group (50%) and the control group (43%, log-rank test = 0.344). Subgroup analysis, adjusted for baseline factors and possibly post-hoc, found a statistically significant benefit of treatment on dorsal ulcers but not plantar ulcers.

Kerecis Omega3 Wound

Lullove et al. (2021, 2022) reported interim results and Lantis et al. (2023) reported the final results of a RCT of Omega3 Wound (Kerecis) plus standard wound care compared to standard care alone in individuals with diabetic lower extremity skin ulcers (Table 18). (44, 45, 46) The primary outcome of the trial was healing at 12 weeks. Complete ulcer healing was based on the site investigator's assessment, as evidenced by complete (100%) re-epithelialization without drainage and need of dressing. An independent panel of wound care experts who were blinded to the patient allocation process and the principal investigator's assessment reviewed all study-related decisions made by the site investigators and confirmed healing status. Secondary outcomes were time to heal and wound area reduction by percentage at 12 weeks. Patients underwent a 2-week run-in period prior to randomization. If the ulcer reduced in area by 20% or more after 14 days of standard care, the patient was excluded as a screening failure. If the wound area was reduced by less than 20%, the patient was randomized and enrolled in the study.

Study results are summarized in Table 19. At 12 weeks, the complete healing rate was significantly higher in the intervention arm (57% vs 31%), but time to healing did not differ between groups for wounds that healed completely. Among the subset of wounds that did not heal completely by 12 weeks (n = 65), there was a larger percent wound reduction in the intervention group (86% vs 64%; $p = .03$). Of the 45 participants whose wound healed during the

12 weeks of the trial, 42 were available for follow-up 6 to 12 months following healing. 3 (11%) ulcer recurrences were reported in the intervention arm compared to 1 (7%) in the control arm.

Study limitations are detailed in Tables 20 and 21. Notably, 2 larger RCTs are registered and reported as completed but have not been published.

Table 18. Randomized Controlled Trial of Omega3 Wound for Diabetic Foot Ulcers- Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Lantis et al. (2023) (46) Lullove et al. (2021) (44, 45) NCT04133493	US	16	2019-2022	Diabetic foot ulcer for a minimum of 4 weeks, adequate renal function and perfusion to the affected extremity Mean age, 60 years 69% Men 80% White 7% Black Mean wound size, 4.4 cm	Omega3 Wound plus standard of care (n=51)	Standard of care (n=51)

US: United States.

Table 19. Randomized Controlled Trial of Omega3 Wound for Diabetic Foot Ulcers-Results

Study	Wound Healed at 12 weeks	Time to Heal	Percent Wound Reduction at 12 Weeks for Wounds that did not heal	Adverse Events
Lantis et al. (2023) (46) Lullove et al. (2021) (44, 45) NCT04133493				
N analyzed	102		65	
Omega3 Wound	57% (29/51)	Mean 7 weeks in both groups	86%	3
Standard of Care	31% (16/51)		64%	5
p	.02		.03	

Table 20. Randomized Controlled Trial of Omega3 Wound for Diabetic Foot Ulcers-Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e

Lantis et al. (2023) (46) Lullove et al. (2021) (44, 45) NCT04133493	4. Lack of racial and ethnic diversity		3. Standard of care patients received additional dressing changes at home, which could have potentially exposed the wound to unknown factors.		
--	--	--	---	--	--

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5: Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 21. Randomized Controlled Trial of Omega3 Wound for Diabetic Foot Ulcers-Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Lantis et al. (2023) (46) Lullove et al. (2021) (44, 45) NCT 04133493			3. Two larger RCTs are reported as completed on clinicaltrials.gov but have not been published (NCT04257370 and NCT04537520)	1, 2. 25% of participants did not complete week 12. Although they were included in the primary ITT analysis, the method of imputation was unclear.		3. Confidence intervals not reported

ITT: intention-to-treat; RCT: randomized controlled trial.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other

Subsection Summary: Bioengineered Skin Substitutes Other Than Apligraf, Dermagraft, AlloPatch, Integra, mVASC or TheraSkin for Diabetic Lower-Extremity Ulcers, Integra

Results from a multicenter RCT showed some benefit of DermACELL that was primarily for the subgroup of patients who only required a single application of the ADM. Studies are needed to further define the population who might benefit from this treatment. Additional study with a larger number of subjects is needed to evaluate the effect of GraftJacket, DermACELL, Cytal, PriMatrix, and Oasis Wound Matrix, compared with current SOC or other advanced wound therapies. Keresis has RCTs that are reported as completed on clinicaltrials.gov but which have not been published (NCT04257370 and NCT04537520).

Lower-Extremity Ulcers due to Venous Insufficiency

Clinical Context and Therapy Purpose

The purpose of bio-engineered soft tissue substitutes in individuals who have lower extremity ulcers due to venous insufficiency is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals who have lower extremity ulcers due to venous insufficiency.

Interventions

The therapy being considered is bioengineered skin substitutes.

Comparators

The following therapies are currently being used: SOC which includes debridement of necrotic tissue and compression.

A Cochrane review by O'Meara et al. (2012) that evaluated compression for venous leg ulcers included 48 RCTs with 59 different comparisons. (47) Most RCTs were small. Measures of healing were the time to complete healing, the proportion of ulcers healed within the trial

period (typically 12 weeks), the change in ulcer size, and the rate of change in ulcer size. Evidence from 8 trials indicated that venous ulcers healed more rapidly with compression than without. Findings suggested that multicomponent systems (bandages or stockings) were more effective than single-component compression. Also, multicomponent systems containing an elastic bandage appeared more effective than those composed mainly of inelastic constituents. Although these meta-analyses did not include time to healing, studies included in the review reported the mean time to ulcer healing was approximately 2 months, while the median time to healing in other reports was 3 to 5 months.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, and QOL.

The primary endpoints of interest for trials of wound closure are as follows, consistent with guidance from the FDA for industry in developing products for treatment of chronic cutaneous ulcer and burn wounds:

- Incidence of complete wound closure.
- Time to complete wound closure (reflecting accelerated wound closure).
- Incidence of complete wound closure following surgical wound closure.
- Pain control.

Time to wound closure can be measured at 6 months with longer-term outcomes apparent by 1 year. Complex wounds may require more than 6 months to heal.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials, with preference for RCTs* were sought.
- In the absence of such trials, comparative observational studies, with preference for prospective studies were sought.
- To assess longer-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Within each category of study design, larger sample size studies and longer duration studies were preferred.
- Studies with duplicative or overlapping populations were excluded.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials

Apligraf

Falanga et al. (1998) reported on a multicenter randomized trial of Apligraf living cell therapy. (48) A total of 293 patients with venous insufficiency and clinical signs of venous ulceration were randomized to compression therapy alone or to compression therapy and treatment with Apligraf. Apligraf was applied up to a maximum of 5 (mean, 3.3) times per patient during the initial 3 weeks. The primary end points were the percentage of patients with complete healing

by 6 months after initiation of treatment and the time required for complete healing. At 6-month follow-up, the percentage of patients healed was higher with Apligraf (63% vs 49%), and the median time to complete wound closure was shorter (61 days vs 181 days). Treatment with Apligraf was superior to compression therapy in healing larger (>1000 mm²) and deeper ulcers and ulcers of more than 6 months in duration. There were no symptoms or signs of rejection, and the occurrence of adverse events was similar in both groups.

Oasis Wound Matrix

Mostow et al. (2005) reported on an industry-sponsored multicenter (12 sites) randomized trial that compared weekly treatment using Oasis Wound Matrix (xenogenic collagen scaffold from porcine small intestinal mucosa) with SOC in 120 patients who had chronic ulcers due to venous insufficiency that had not adequately responding to conventional therapy. (49) Healing was assessed weekly for up to 12 weeks, with follow-up performed after 6 months to assess recurrence. After 12 weeks of treatment, there was a significant improvement in the percentage of wounds healed in the Oasis group (55% vs 34%). After adjusting for baseline ulcer size, patients in the Oasis group were 3 times more likely to heal than those in the group receiving SOC. Patients in the SOC group whose wounds did not heal by week 12 were allowed to cross over to Oasis treatment. None of the healed patients treated with Oasis wound matrix seen for the 6-month follow-up experienced ulcer recurrence.

A research group in Europe has described 2 comparative studies of the Oasis matrix for mixed arteriovenous ulcers. In a quasi-randomized study, Romanelli et al. (2007) compared the efficacy of 2 extracellular matrix-based products, Oasis and Hyaloskin (extracellular matrix with hyaluronic acid). (50) Fifty-four patients with mixed arteriovenous leg ulcers were assigned to the 2 arms based on order of entry into the study; 50 patients completed the study. Patients were followed twice weekly, and dressings changed more than once a week, only when necessary. After 16 weeks of treatment, complete wound closure was achieved in 82.6% of Oasis-treated ulcers compared with 46.2% of Hyaloskin-treated ulcers. Oasis treatment significantly increased the time to dressing change (mean, 6.4 days vs 2.4 days), reduced pain on a 10-point scale (3.7 vs 6.2), and improved patient comfort (2.5 vs 6.7).

Romanelli et al. (2010) compared Oasis with a moist wound dressing (SOC) in 23 patients with mixed arteriovenous ulcers and 27 patients with venous ulcers. (51) The trial was described as randomized, but the method of randomization was not described. After the 8-week study period, patients were followed monthly for 6 months to assess wound closure. Complete wound closure was achieved in 80% of the Oasis-treated ulcers at 8 weeks compared with 65% of the SOC group. On average, Oasis-treated ulcers achieved complete healing in 5.4 weeks compared with 8.3 weeks for the SOC group. Treatment with Oasis also increased the time to dressing change (5.2 days vs 2.1 days) and the percentage of granulation tissue formed (65% vs 38%).

Subsection Summary: Apligraf or Oasis Wound Matrix for Lower-Extremity Ulcers due to Venous Insufficiency

RCTs have demonstrated the efficacy of Apligraf or Oasis Wound Matrix over SOC for lower-extremity ulcers due to venous insufficiency.

Bioengineered Skin Substitutes Other Than Apligraf or Oasis Wound Matrix for Lower-Extremity Ulcers due to Venous Insufficiency

Dermagraft

Dermagraft living cell therapy has been approved by the FDA for repair of diabetic foot ulcers. Use of Dermagraft for venous ulcers is an off-label indication. Harding et al. (2013) reported an open-label multicenter RCT that compared Dermagraft plus compression therapy (n=186) with compression therapy alone (n=180). (52) The trial had numerous inclusion and exclusion criteria that restricted the population to patients who had nonhealing ulcers with compression therapy but had the capacity to heal. ITT analysis revealed no significant difference between the 2 groups in the primary outcome measure, the proportion of patients with completely healed ulcers by 12 weeks (34% Dermagraft vs 31% control). Prespecified subgroup analysis revealed a significant improvement in the percentage of wounds healed for ulcers of 12 months or less in duration (52% vs 37%) and for ulcers of 10 cm or less in diameter (47% vs 39%). There were no significant differences in the secondary outcomes of time to healing, complete healing by week 24, and percent reduction in ulcer area.

DermACELL

Cazzell (2019) published an RCT on DermACELL ADM for venous leg ulcers in 18 patients (see Table 22). (53) This was part of a larger study of the acellular dermal matrix for chronic wounds of the lower extremity in 202 patients; the component on diabetic lower extremity ulcers was previously reported by Cazzell et al. (2017) and is described above. (39) When including patients who required more than 1 application of the ADM, the percent of wounds closed at 24 weeks was 29.4% with DermACELL and 33.3% with SOC, suggesting no benefit DermACELL for the treatment of venous ulcers in this small substudy.

Table 22. Summary of Key RCT Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Cazzell (2019) (53) NCT01970163	U.S.	7	2013- 2016	Venous leg ulcer present for at least 60 days (n=18)	1 or 2 applications of DermACELL plus SOC (n=18)	SOC (debridement and compression, n=10)

RCT: randomized controlled trial; SOC: standard of care.

Subsection Summary: Bioengineered Skin Substitutes Other Than Apligraf or Oasis Wound Matrix for Lower-Extremity Ulcers due to Venous Insufficiency

In a moderately large RCT, Dermagraft was not shown to be more effective than controls in the primary or secondary end points for the entire population and was slightly more effective than controls (an 8%-15% increase in healing) only in subgroups of patients with ulcer duration of 12

months or less or wound diameter of 10 cm or less. An initial study with 18 patients found that DermACELL (ADM) was not more effective than SOC. An initial study with 18 patients found that and DermACELL (ADM) was not more effective than SOC.

Deep Dermal Burns

Clinical Context and Therapy Purpose

The purpose of bio-engineered soft tissue substitutes in individuals who have deep dermal burns is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with deep dermal burns.

Interventions

The therapy being considered is bioengineered skin substitutes.

Comparators

The following therapies are currently being used; standard therapy for burns.

Outcomes

The general outcomes of interest are disease-specific survival, symptoms, change in disease status, morbid events, functional outcomes, QOL, and treatment-related morbidity.

The primary endpoints of interest for trials of wound closure are as follows, consistent with guidance from the FDA for industry in developing products for treatment of chronic cutaneous ulcer and burn wounds:

- Incidence of complete wound closure.
- Time to complete wound closure (reflecting accelerated wound closure).
- Incidence of complete wound closure following surgical wound closure.
- Pain control.

Time to wound closure can be measured at 6 months with longer-term outcomes apparent by 1 year.

Study Selection Criteria

- To assess efficacy outcomes, comparative controlled prospective trials, with preference for RCTs* were sought.
- In the absence of such trials, comparative observational studies, with preference for prospective studies were sought.
- To assess longer-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.

- Within each category of study design, larger sample size studies and longer duration studies were preferred.
- Studies with duplicative or overlapping populations were excluded.

* Includes various RCT designs such as adaptive trials, pragmatic trials, and cluster trials

Epicel

One case series from 2000 has described the treatment of 30 severely burned patients with Epicel. (54) The cultured epithelial autografts were applied to a mean of 37% of total body surface area (TBSA). Epicel achieved permanent coverage of a mean of 26% of TBSA, an area similar to that covered by conventional autografts (mean, 25%). Survival was 90% in these severely burned patients.

Integra Dermal Regeneration Template

A 2013 study compared Integra with split-thickness skin graft and with viscose cellulose sponge (Cellonex), using three 10x5 cm test sites on each of 10 burn patients. (55) The surrounding burn area was covered with meshed autograft. Biopsies were taken from each site on days 3, 7, 14, and 21, and at months 3 and 12. The tissue samples were stained and examined for markers of inflammation and proliferation. The Vancouver Scar Scale was used to assess scars. At 12-month follow-up, the 3 methods resulted in similar clinical appearance, along with similar histologic and immunohistochemical findings.

Branski et al. (2007) reported on a randomized trial that compared Integra with a standard autograft-allograft technique in 20 children with an average burn size of 73% TBSA (71% full-thickness burns). (56) Once vascularized (about 14-21 days), the Silastic epidermis was stripped and replaced with thin (0.05-0.13 mm) epidermal autograft. There were no significant differences between the Integra group and controls in burn size (70% vs 74% TBSA), mortality (40% vs 30%), and hospital length of stay (41 vs 39 days), all respectively. Long-term follow-up revealed a significant increase in bone mineral content and density (24 months) and improved scarring in terms of height, thickness, vascularity, and pigmentation (at 12 months and 18-24 months) in the Integra group. No differences were observed between groups in the time to first reconstructive procedure, cumulative reconstructive procedures required during 2 years, and cumulative operating room time required for these procedures. The authors concluded that Integra can be used for immediate wound coverage in children with severe burns without the associated risks of cadaver skin.

Heimbach et al. (2003) reported on a multicenter (13 U.S. burn care facilities) postapproval study involving 222 burn injury patients (36.5% TBSA; range, 1%-95%) who were treated with Integra Dermal Regeneration Template. (57) Within 2 to 3 weeks, the dermal layer regenerated, and a thin epidermal autograft was placed over the wound. The incidence of infection was 16.3%. Mean take rate (absence of graft failure) of Integra was 76.2%; the median take rate was 98%. The mean take rate of epidermal autograft placed over Integra was 87.7%; the median take rate was 95%.

Hicks et al. (2019) conducted a systematic review of Integra dermal regeneration template for the treatment of acute full thickness burns and burn reconstruction. (58) A total of 72 studies with 1084 patients (4 RCTs, 4 comparative studies, 5 cohort studies, 2 case control studies, 24 case series, and 33 case reports) were included in the review. The majority of patients (74%) were treated with Integra for acute burns, and the remainder (26%) for burn reconstruction. The take of the skin substitute was 86% (range 0–100%) for acute burn injuries and 95% (range 0–100%) for reconstruction. The take of the split-thickness skin graft over the template was 90% for acute burn injuries and 93% for reconstruction. There was high variability in reporting of outcomes, but studies generally supported satisfactory cosmetic results in patients who have insufficient autograft and improvement in range of motion in patients who were treated with Integra for burn reconstruction. There was an overall complication rate of 13%; primarily due to infection, graft loss, hematoma formation, and contracture.

An infection rate of 18% was noted in a systematic review of complication rates in 10 studies that used Integra dermal regeneration template for burns. (59)

Omega3 Wound

Luze et al. (2022) conducted a systematic review of the use of acellular fish skin grafts in burn wound management. (60) The reviewers identified 5 studies of Omega3 Wound but no RCTs. The identified studies were preclinical (animal), case series, retrospective observational, and 1 small (N = 21) cohort study. The review authors concluded that while the approach is promising, large-cohort studies are needed.

ReCell Autologous Cell Harvesting Device

Two RCTs have evaluated ReCell for deep dermal burns (Table 23). (61, 62)

In both studies, 2 similar areas with a burn injury in the same individual were randomized to the control or treatment intervention (i.e., all participants received both treatments). The studies differed in their populations, interventions, and outcome measures. In the earlier study, participants all had deep partial thickness burns, while in the 2019 study the population included individuals with mixed-depth, full thickness burns. Holmes 2018 was a head-to-head comparison of ReCell alone versus skin grafting alone, and Holmes et al (2019) compared ReCell in combination with skin grafting. In the earlier study, the primary effectiveness endpoints were the incidence of wound closure at 4 weeks and the incidence of complete donor site healing at 1 week. In the 2019 trial, the co-primary effectiveness endpoints were non-inferiority of the incidence of RECELL-treated site closure by week 8 when compared to the control, and the superiority of the 37% relative reduction in donor skin for the ReCell treatment when compared with the control.

Study results are detailed in Table 24 and limitations in Tables 25 and 26. Although the ReCell device was comparable to standard care on outcomes such as complete wound closure; confidence in the strength of the overall body of evidence is limited by individual study limitations and heterogeneity of populations, interventions, and outcome measures across studies. Additional RCT evidence in the intended use population is needed.

Table 23. Randomized Controlled Trials of ReCell for Thermal Burns-Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Holmes et al. (2018) (62) NCT 01138917	US	9	2010-2015	Individuals ages 18 to 65 years, with acute, deep partial-thickness thermal burns from 1% to 20% TBSA that required autografting for definitive closure.	ReCell device N=101	Meshed STSG Treatment N=101
Holmes et al. (2019) (61) NCT 02380612	US	6	2015-2017	Individuals ages 5 years or older, with acute thermal burn involving 5% to 50% of TBSA that underwent autografting for definitive closure	ReCell device treatment applied over STSG N=30	Meshed STSG Treatment Alone N=30

STSG:: Split-thickness skin grafts; TBSA: total body surface area.

Table 24a. Randomized Controlled Trials of ReCell for Thermal Burns-Results

Study	Wound Closure (95% re-epithelialization) at 4 weeks	Wound Closure (95% re-epithelialization) at 8 weeks	Complete donor site healing at 1 week (100% reepithelialization)
Holmes et al. (2018) (62) NCT01138917			
ReCell	81/83 (97.6%)		21.8%
STSG	83/83 (100%)		10.0%
Between-group difference	-2.4% (95% CO: -8.4% to 2.3%)		p=.04
Holmes et al. (2019) (61) NCT02380612			
ReCell plus STSG	50%	24/26 (92%)	
STSG alone	48%	22/26 (85%)	
Between-group difference		-7.7% Upper limit of the 97.5% CI 6.4% (i.e., within the pre-defined non-inferiority margin 10%)	

AE: adverse events; CI: confidence interval; ND: no significant difference.

Table 24b. Randomized Controlled Trials of ReCell for Thermal Burns-Results

Study	Relative Reduction in Donor Skin	Pain (VAS)	Participant Satisfaction and Scar Assessment	Adverse Events (Incidence)
Holmes et al. (2018) (62) NCT01138917				
ReCell		NSD at 16 weeks (data in figure)	NSD in subject satisfaction with appearance or in scarring at 16, 24, and 52 weeks (data in figures)	Treatment site: 35.6% Donor site: 4.0%
STSG				Treatment site: 21.8% Donor site: 6.9%
Between-group difference				Treatment site: p =.0013 Donor site: 6.9% p =.25
Holmes et al. (2019) (61) NCT02380612				
ReCell plus STSG	368 (SD 150) cm ²	NSD between groups in treatment area pain from week 1 to week 12 or week 52	NSD in subject satisfaction with appearance or in scar assessment at any time point	NSD between groups in pre-specified safety events 17 individuals (57%) experienced AEs at control and ReCell sites; 27% had mild AEs, 37% moderate AEs. 1 death, attributed to underlying condition
STSG alone	264 (SD 119) cm ²			
Between-group difference	32%; p<.001			

AE: adverse events; CI: confidence interval; ND: no significant difference.

Table 25. Randomized Controlled Trials of ReCell for Thermal Burns-Study Relevance

Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Holmes et al. (2018) (62) NCT 01138917					
Holmes et al.	2. Participants had mixed			5. Unclear if 32% reduction in donor	

(2019) (61) NCT 02380612	depth full-thickness burns			site skin is clinically meaningful	
--------------------------------	-------------------------------	--	--	---------------------------------------	--

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^aPopulation key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^bIntervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5: Other.

^cComparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^eFollow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 26. Randomized Controlled Trials of ReCell for Thermal Burns-Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Holmes et al. (2018) (62) NCT 01138917				83/101 participants evaluated in modified per protocol analysis	Non- inferiority margin based on 90 subjects	
Holmes et al. (2019) (61) NCT 02380612				26/30 participants evaluated in per protocol analysis		3. confidence intervals not reported

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^aAllocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^bBlinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^cSelective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^ePower key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^fStatistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

Section Summary: Deep Dermal Burns

Epicel is FDA-approved under a humanitarian device exemption (HDE) for the treatment of deep dermal or full-thickness burns comprising a TBSA of 30% or more, with patient survival of 90%. Integra Dermal Regeneration Template has been compared with autograft in a within-subject study and with autograft-allograft in a small RCT with 10 patients per group. Outcomes are at least as good as with autograft or allograft, with a reduction in scarring and without risks associated with cadaver skin. This product has also been studied in a large series with over 222 burn patients, showing a take rate of 76% and with a take rate of epidermal autograft placed over Integra of 87.7%.

The ReCell device has been evaluated in 2 RCTs. One RCT evaluated ReCell as an adjunct to meshed autologous skin grafting and the other compared ReCell head-to-head with skin grafting. Although the ReCell device was comparable to standard care on outcomes such as complete wound closure, confidence in the strength of the overall body of evidence is limited by individual study limitations and heterogeneity of populations, interventions, and outcome measures across studies. Additional RCT evidence in the intended use population is needed.

Other Indications

Dystrophic Epidermolysis Bullosa

OrCel was approved under a HDE for use in patients with dystrophic epidermolysis bullosa undergoing hand reconstruction surgery, to close and heal wounds created by the surgery, including those at donor sites. HDE status has been withdrawn for Dermagraft for this indication.

Fivenson et al. (2003) reported the off-label use of Apligraf in 5 patients with recessive dystrophic epidermolysis bullosa who underwent syndactyly release. (64)

Subsection Summary: Dystrophic Epidermolysis Bullosa

Dystrophic epidermolysis bullosa is a rare disorder. Because this is a rare disorder, it is unlikely that RCTs will be conducted to evaluate whether OrCel improves health outcomes for this condition.

Punch Biopsy Wounds

Baldursson et al. (2015) reported a double-blinded RCT with 81 patients (162 punch biopsy wounds) that compared Kerecis Omega3 Wound (derived from fish skin) with Oasis SIS ECM (porcine small intestinal submucosa extracellular matrix). (65) The primary outcome (the percentage of wounds healed at 28 days) was similar for the fish skin ADM (95%) and the porcine SIS ECM (96.3%). The rate of healing was faster with Kerecis Omega3 ($p=0.041$). At 21 days, 72.5% of the fish skin ADM group had healed compared with 56% of the porcine SIS ECM

group. Interpretation of this study is limited because it did not include an accepted control condition for this indication.

Split-Thickness Donor Sites

There is limited evidence to support the efficacy of OrCel compared with SOC for the treatment of split-thickness donor sites in burn patients. Still et al. (2003) examined the safety and efficacy of bilayered OrCel to facilitate wound closure of split-thickness donor sites in 82 severely burned patients. (66) Each patient had 2 designated donor sites that were randomized to a single treatment of OrCel or standard dressing (Biobrane-L). The healing time for OrCel sites was significantly shorter than for sites treated with a standard dressing, enabling earlier re-cropping. OrCel sites also exhibited a nonsignificant trend for reduced scarring. Additional studies are needed to evaluate the effect of this product on health outcomes.

Pressure Ulcers

Brown-Etris et al. (2019) reported an RCT of 130 patients with stage 3 or stage 4 pressure ulcers who were treated with Oasis Wound Matrix (extracellular collagen matrix derived from porcine small intestinal submucosa) plus SOC or SOC alone. (66) At 12 weeks, the proportion of wounds healed in the collagen matrix group was 40% compared to 29% in the SOC group. This was not statistically significant ($p=0.111$). There was a statistical difference in the proportion of patients who achieved 90% wound healing (55% vs. 38% $p=0.037$), but complete wound healing is the preferred and most reliable measure. It is possible that longer follow-up may have identified a significant improvement in the percent of wounds healed. The study did include 6-month follow-up, but there was high loss to follow-up and an insufficient number of patients at this time point for statistical comparison.

In the propensity matched study by Gurtner et al. (2020) described above, Theraskin improved the healing rate of pressure ulcers by 20% (66.7% vs 46.8%). (67)

Miscellaneous

In addition to indications previously reviewed, off-label uses of bioengineered skin substitutes have included inflammatory ulcers (e.g., pyoderma gangrenosum, vasculitis), scleroderma digital ulcers, post-keloid removal wounds, genetic conditions, and variety of other conditions. (68) Products that have been FDA-approved or -cleared for one indication (e.g., lower-extremity ulcers) have also been used off-label in place of other FDA-approved or -cleared products (e.g., for burns). (69) No controlled trials were identified for these indications.

Summary of Evidence

Breast Reconstruction

For individuals who are undergoing breast reconstruction who receive allogeneic acellular dermal matrix (ADM) products, the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are symptoms, morbid events, functional outcomes, quality of life, and treatment-related morbidity. A systematic review found no difference in overall complication rates with ADM allograft compared with standard procedures for breast reconstruction. Reconstructions with ADM have been reported to have higher seroma,

infection, and necrosis rates than reconstructions without ADM. However, capsular contracture and malposition of implants may be reduced. Thus, in cases where there is limited tissue coverage, the available evidence may inform patient decision making about reconstruction options. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Tendon Repair

For individuals who are undergoing tendon repair who receive GraftJacket, the evidence includes an RCT. Relevant outcomes are symptoms, morbid events, functional outcomes, quality of life, and treatment-related morbidity. The RCT identified found improved outcomes with the GraftJacket ADM allograft for rotator cuff repair. Although these results were positive, additional study with a larger number of patients is needed to evaluate the consistency of the effect. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Surgical Repair of Hernias or Parastomal Reinforcement

For individuals who are undergoing surgical repair of hernias or parastomal reinforcement who receive acellular collagen-based scaffolds, the evidence includes RCTs. Relevant outcomes are symptoms, morbid events, functional outcomes, quality of life, and treatment-related morbidity. Several comparative studies including RCTs have shown no difference in outcomes between tissue-engineered skin substitutes and either standard synthetic mesh or no reinforcement. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Diabetic Lower-Extremity Ulcers

For individuals who have diabetic lower-extremity ulcers who receive AlloPatch, Apligraf, Dermagraft, Integra, mVASC or TheraSkin the evidence includes RCTs. Relevant outcomes are disease-specific survival, symptoms, change in disease status, morbid events, and quality of life. RCTs have demonstrated the efficacy of AlloPatch (reticular ADM), Apligraf and Dermagraft (living cell therapy), Integra (biosynthetic), mVASC, and TheraSkin over the standard of care. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have diabetic lower-extremity ulcers who receive ADM products other than AlloPatch, Apligraf, Dermagraft, Integra, mVASC, or TheraSkin, the evidence includes RCTs. Relevant outcomes are disease-specific survival, symptoms, change in disease status, morbid events, and quality of life. Results from a multicenter RCT showed some benefit of DermACELL that was primarily for the subgroup of patients who only required a single application of the ADM. Studies are needed to further define the population who might benefit from this treatment. Additional study with a larger number of subjects is needed to evaluate the effect of GraftJacket, TheraSkin, DermACELL, Cytal, PriMatrix, and Oasis Wound Matrix, compared with current SOC or other advanced wound therapies. An RCT of Omega3 Wound (Kerecis) has been published and 2 larger RCTs are registered and reported as completed but have not been

published. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Lower-Extremity Ulcers due to Venous Insufficiency

For individuals who have lower-extremity ulcers due to venous insufficiency who receive Apligraf or Oasis Wound Matrix, the evidence includes RCTs. Relevant outcomes are disease-specific survival, symptoms, change in disease status, morbid events, and quality of life. RCTs have demonstrated the efficacy of Apligraf living cell therapy and xenogenic Oasis Wound Matrix over the standard of care. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have lower-extremity ulcers due to venous insufficiency who receive bioengineered skin substitutes other than Apligraf or Oasis Wound Matrix, the evidence includes RCTs. Relevant outcomes are disease-specific survival, symptoms, change in disease status, morbid events, and quality of life. In a moderately large RCT, Dermagraft was not shown to be more effective than controls for the primary or secondary end points in the entire population and was only slightly more effective than controls (an 8%-15% increase in healing) in subgroups of patients with ulcer durations of 12 months or less or size of 10 cm or less. Additional study with a larger number of subjects is needed to evaluate the effect of the xenogenic PriMatrix skin substitute vs the current standard of care. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Dystrophic Epidermolysis Bullosa

For individuals who have dystrophic epidermolysis bullosa who receive OrCel, the evidence includes case series. Relevant outcomes are symptoms, change in disease status, morbid events, and quality of life. OrCel was approved under a humanitarian drug exemption for use in patients with dystrophic epidermolysis bullosa undergoing hand reconstruction surgery, to close and heal wounds created by the surgery, including those at donor sites. Outcomes have been reported in small series (e.g., 5 patients). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Deep Dermal Burns

For individuals who have deep dermal burns who receive bioengineered skin substitutes (i.e., Epicel, Integra Dermal Regeneration Template), the evidence includes RCTs. Relevant outcomes are disease-specific survival, symptoms, change in disease status, morbid events, functional outcomes, quality of life, and treatment-related morbidity. Overall, few skin substitutes have been approved, and the evidence is limited for each product. Epicel (living cell therapy) has received U.S. Food and Drug Administration (FDA) approval under a humanitarian device exemption for the treatment of deep dermal or full-thickness burns comprising a total body surface area of 30% or more. Comparative studies have demonstrated improved outcomes for biosynthetic skin substitute Integra Dermal Regeneration Template for the treatment of burns. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Practice Guidelines and Position Statements

National Institute for Health and Care Excellence

In 2023, NICE updated its guidance on the prevention and management of diabetic foot problems. (70) The Institute recommended that clinicians “consider dermal or skin substitutes as an adjunct to standard care when treating diabetic foot ulcers, only when healing has not progressed and on the advice of the multidisciplinary foot care service.”

In 2019, NICE published guidance on the ReCell system for treating skin loss, scarring, and depigmentation after burn injury. (71) The guidance recommended that additional research was needed to address the uncertainties regarding the potential benefits of ReCell.

Ongoing and Unpublished Clinical Trials

Some currently ongoing or unpublished trials that might influence this policy are listed in Table 27.

Table 27. Summary of Key Trials

NCT Number	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT05291169	A Randomized, Multicenter, Open Label Study Comparing Omeza Combination Therapy with Standard of Care to Standard of Care alone for Chronic Venous Leg Ulcers over the course of 4 weeks	110	Oct 2023
NCT05084183	An Adaptive, Randomized, Controlled Trial Evaluating the Effectiveness of PermeaDerm® (PD) as Compared to Mepilex Ag® Used as Standard of Care in the Treatment of Adult and Pediatric Partial Thickness Burns	68	Nov 2023
NCT05439746	Clinical Trial to Assess the Efficacy of Microlyte Matrix on the Healing of Surgically Created Partial Thickness Donor Site Wounds on Patients Requiring Split-thickness Skin Grafting	53	Jan 2024
NCT05506215	A Prospective, Multicenter, Open Label, Randomized, Controlled Clinical Study Evaluating the Effect of NovoSorb® SynPath™ Dermal Matrix Compared to Standard of Care (SOC) In the Treatment of Nonresponsive, Chronic Diabetic Foot Ulcers.	138	Mar 2024

NCT05372809	Closure Obtained With Vascularized Epithelial Regeneration for DFUs With SkinTE®	100	Jun 2024
NCT02587403 ^a	A Randomized, Prospective Study Comparing Fortiva™ Porcine Dermis vs. Strattice™ Reconstructive Tissue Matrix in Patients Undergoing Complex Open Primary Ventral Hernia Repair	120	Feb 2024
NCT04927702	Assessment of Wound Closure Comparing Synthetic Hybrid-Scale Fiber Matrix (Restrata®) With Standard of Care in Treating Diabetic Foot Ulcers (DFU) and With Living Cellular Skin Substitute (Apligraf®) in Treating Venous Leg Ulcers (VLU)	170	Jul 2024
NCT06035536	A Multi-Center, Randomized Controlled Clinical Investigation Evaluating Wound Closure With Symphony™ Versus Standard of Care in the Treatment of Non-Healing Diabetic Foot Ulcers	120	Dec 2024
NCT05517902	A Phase 3 Multicenter, Single-Arm, Open-Label Study Evaluating the Safety, Tolerability and Efficacy of StrataGraft® Construct in Pediatric Subjects With Deep Partial Thickness (DPT) Thermal Burns	50	Jun 2025
NCT04090424	A Pivotal Study to Assess the Safety and Effectiveness of NovoSorb® Biodegradable Temporizing Matrix (BTM) in the Treatment of Severe Burn Skin Injuries	150	Dec 2025
NCT03394612	A Phase II, Prospective, Intra-patient Randomised Controlled, Multicentre Study to Evaluate the Safety and Efficacy of an Autologous Bio-engineered Dermo-epidermal Skin Substitute (EHSK-KF; denovoSkin) for the Treatment of Full-Thickness Defects in Adults and Children in Comparison to Autologous Split-thickness Skin Grafts (STSG)	20	Dec 2026
Unpublished			

NCT02322554	The Registry of Cellular and Tissue Based Therapies for Chronic Wounds and Ulcers	50,000	Jan 2020
NCT03925286 ^a	A Prospective Randomized Clinical Trial Comparing Multi-layer Bandage Compression Therapy With and Without a Biologically Active Human Skin Allograft (Theraskin) for the Treatment of Chronic Venous Leg Ulcers	100	Dec 2020
NCT03589586 ^a	An Open-Label Trial to Assess the Clinical Effectiveness of DermACELL AWM in Subjects With Chronic Venous Leg Ulcers	100	Jan 2021
NCT03881254	A Multi-center, Randomized Controlled Clinical Trial Evaluating the Effects of SkinTE™ in the Treatment of Wagner One Diabetic Foot Ulcers	100	Jul 2021
NCT04198441	A Randomized, Multicenter, Open Label Study Comparing the Omeza® Products Bundle to Standard of Care for Chronic Venous Leg Ulcers and Chronic Diabetic Foot Ulcers	78	Dec 2021
NCT04257370 ^a	An Open Label, Randomized Controlled Study to Compare Healing of Severe Diabetic Foot Ulcers and Forefoot Amputations in Diabetics With and Without Moderate Peripheral Arterial Disease Treated With Kerecis Omega3 Wound and SOC vs. SOC Alone	330	Oct 2022
NCT04537520 ^a	Interventional Multi-Center Post Market Randomized Controlled Open-Label Clinical Trial Comparing Kerecis Omega3 Wound Versus SOC in Hard to Heal Diabetic Foot Wounds	180	Dec 2022
NCT04918784	Assessment of Wound Closure Comparing Synthetic Hybrid-Scale Fiber Matrix (Restrata®, Acera Surgical, Inc.) With Standard of Care in Treating Diabetic Foot Ulcer	46	Dec 2022
NCT05883098	Effectiveness of Supra SDRM® vs. Fibracol Plus Collagen in the Treatment of Diabetic Foot Ulcers: a Pilot Randomized Controlled Trial	30	Jun 2023

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member's benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

CPT Codes	15271, 15272, 15273, 15274, 15275, 15276, 15277, 15278, 15777
HCPCS Codes	A2002, A2004, A2005, A2006, A2007, A2008, A2009, A2010, A2011, A2012, A2013, A2014, A2015, A2016, A2017, A2018, A2019, A2020, A2021, A2022, A2023, A2024, A2025, A2026, A2027, A2028, A2029, A4100, A6460, A6461, C1832, C5271, C5272, C5273, C5274, C5275, C5276, C5277, C5278, C9354, C9356, C9358, C9360, C9363, C9364, Q4100, Q4101, Q4102, Q4103, Q4104, Q4105, Q4106, Q4107, Q4108, Q4110, Q4111, Q4112, Q4113, Q4114, Q4115, Q4116, Q4117, Q4118, Q4121, Q4122, Q4123, Q4124, Q4125, Q4126, Q4127, Q4128, Q4130, Q4134, Q4135, Q4136, Q4141, Q4142, Q4143, Q4146, Q4147, Q4149, Q4152, Q4158, Q4161, Q4164, Q4165, Q4166, Q4167, Q4175, Q4179, Q4182, Q4193, Q4195, Q4196, Q4197, Q4200, Q4202, Q4203, Q4220, Q4222, Q4226 (Deleted 01/23: C1849)

*Current Procedural Terminology (CPT®) ©2023 American Medical Association: Chicago, IL.

References

1. Snyder DL, Sullivan N, Margolis DJ, Schoelles K. Skin substitutes for treating chronic wounds. Technology Assessment Program Project ID No. WNDR0818. (Prepared by the ECRI Institute-Penn Medicine Evidence-based Practice Center under Contract No. HHS 290-2015-00005-I) Rockville, MD: Agency for Healthcare Research and Quality. February 2020. Available at: <<https://effectivehealthcare.ahrq.gov>> (accessed November 13, 2023).
2. U.S. Food and Drug Administration. Regulatory Considerations for Human Cells, Tissues, and Cellular and Tissue-Based Products: Minimal Manipulation and Homologous Use. Dec 2017. Available at: <<https://www.fda.gov>> (accessed November 13, 2023).
3. U.S. Food and Drug Administration. Executive Summary Breast Implant Special Topics. Mar 2019. Available at: <<https://www.fda.gov>> (accessed November 13, 2023).
4. U.S. Food & Drug Administration. Acellular Dermal Matrix (ADM) Products Used in Implant-Based Breast Reconstruction Differ in Complication Rates: FDA Safety Communication. Mar 2021. Available at: <<https://www.fda.gov>> (accessed November 13, 2023).

5. Davila AA, Seth AK, Wang E, et al. Human acellular dermis versus submuscular tissue expander breast reconstruction: a multivariate analysis of short-term complications. *Arch Plast Surg*. Jan 2013; 40(1):19-27. PMID 23362476
6. Lee KT, Mun GH. Updated evidence of acellular dermal matrix use for implant-based breast reconstruction: a meta-analysis. *Ann Surg Oncol*. Feb 2016; 23(2):600-610. PMID 26438439
7. McCarthy CM, Lee CN, Halvorson EG, et al. The use of acellular dermal matrices in two-stage expander/implant reconstruction: a multicenter, blinded, randomized controlled trial. *Plast Reconstr Surg*. Nov 2012; 130(5 Suppl 2):57S-66S. PMID 23096987
8. Hinchcliff KM, Orbay H, Busse BK, et al. Comparison of two cadaveric acellular dermal matrices for immediate breast reconstruction: A prospective randomized trial. *J Plast Reconstr Aesthet Surg*. May 2017; 70(5):568-576. PMID 28341592
9. Mendenhall SD, Anderson LA, Ying J, et al. The BREASTrial Stage II: ADM breast reconstruction outcomes from definitive reconstruction to 3 months postoperative. *Plast Reconstr Surg Glob Open*. Jan 2017; 5(1):e1209. PMID 28203509
10. Mendenhall SD, Moss WD, Graham EM, et al. The BREASTrial Stage III: Acellular Dermal Matrix Breast Reconstruction Outcomes from 3 Months to 2 Years Postoperatively. *Plast Reconstr Surg*. Jan 01 2023; 151(1):17-24. PMID 36194057
11. Dikmans RE, Negenborn VL, Bouman MB, et al. Two-stage implant-based breast reconstruction compared with immediate one-stage implant-based breast reconstruction augmented with an acellular dermal matrix: an open-label, phase 4, multicentre, randomised, controlled trial. *Lancet Oncol*. Feb 2017; 18(2):251-258. PMID 28012977
12. Barber FA, Burns JP, Deutsch A, et al. A prospective, randomized evaluation of acellular human dermal matrix augmentation for arthroscopic rotator cuff repair. *Arthroscopy*. Jan 2012; 28(1):8-15. PMID 21978432
13. Rashid MS, Smith RDJ, Nagra N, et al. Rotator cuff repair with biological graft augmentation causes adverse tissue outcomes. *Acta Orthop*. Jul 21 2020: 1-7. PMID 32691656
14. Bellows CF, Smith A, Malsbury J, et al. Repair of incisional hernias with biological prosthesis: a systematic review of current evidence. *Am J Surg*. Jan 2013; 205(1):85-101. PMID 22867726
15. Espinosa-de-los-Monteros A, de la Torre JJ, Marrero I, et al. Utilization of human cadaveric acellular dermis for abdominal hernia reconstruction. *Ann Plast Surg*. Mar 2007; 58(3):264-267. PMID 17471129
16. Gupta A, Zahriya K, Mullens PL, et al. Ventral herniorrhaphy: experience with two different biosynthetic mesh materials, Surgisis and Alloderm. *Hernia*. Oct 2006; 10(5):419-425. PMID 16924395
17. Bochicchio GV, De Castro GP, Bochicchio KM, et al. Comparison study of acellular dermal matrices in complicated hernia surgery. *J Am Coll Surg*. Oct 2013; 217(4):606-613. PMID 23973102
18. Roth JS, Zachem A, Plymale MA, et al. Complex ventral hernia repair with acellular dermal matrices: clinical and quality of life outcomes. *Am Surg*. Feb 01 2017; 83(2):141-147. PMID 28228200
19. Bellows CF, Shadduck P, Helton WS, et al. Early report of a randomized comparative clinical trial of Strattice reconstructive tissue matrix to lightweight synthetic mesh in the repair of inguinal hernias. *Hernia*. Apr 2014; 18(2):221-230. PMID 23543334

20. Fleshman JW, Beck DE, Hyman N, et al. A prospective, multicenter, randomized, controlled study of non-cross-linked porcine acellular dermal matrix fascial sublay for parastomal reinforcement in patients undergoing surgery for permanent abdominal wall ostomies. *Dis Colon Rectum*. May 2014; 57(5):623-631. PMID 24819103
21. Kalaiselvan R, Carlson GL, Hayes S, et al. Recurrent intestinal fistulation after porcine acellular dermal matrix reinforcement in enteric fistula takedown and simultaneous abdominal wall reconstruction. *Hernia*. Jun 2020; 24(3):537-543. PMID 31811593
22. Santema TB, Poyck PP, Ubbink DT. Skin grafting and tissue replacement for treating foot ulcers in people with diabetes. *Cochrane Database Syst Rev*. Feb 11 2016; 2:CD011255. PMID 26866804
23. Veves A, Falanga V, Armstrong DG, et al. Graftskin, a human skin equivalent, is effective in the management of noninfected neuropathic diabetic foot ulcers: a prospective randomized multicenter clinical trial. *Diabetes Care*. Feb 2001; 24(2):290-295. PMID 11213881
24. Marston WA, Hanft J, Norwood P, et al. The efficacy and safety of Dermagraft in improving the healing of chronic diabetic foot ulcers: results of a prospective randomized trial. *Diabetes Care*. Jun 2003; 26(6):1701-1705. PMID 12766097
25. Frykberg RG, Marston WA, Cardinal M. The incidence of lower-extremity amputation and bone resection in diabetic foot ulcer patients treated with a human fibroblast-derived dermal substitute. *Adv Skin Wound Care*. Jan 2015; 28(1):17-20. PMID 25407083
26. Zelen CM, Orgill DP, Serena T, et al. A prospective, randomised, controlled, multicentre clinical trial examining healing rates, safety and cost to closure of an acellular reticular allogenic human dermis versus standard of care in the treatment of chronic diabetic foot ulcers. *Int Wound J*. Apr 2017; 14(2):307-315. PMID 27073000
27. Zelen CM, Orgill DP, Serena TE, et al. An aseptically processed, acellular, reticular, allogenic human dermis improves healing in diabetic foot ulcers: A prospective, randomised, controlled, multicentre follow-up trial. *International wound journal*. Oct 2018; 15(5):731-739. PMID 29682897
28. Driver VR, Lavery LA, Reyzelman AM, et al. A clinical trial of Integra Template for diabetic foot ulcer treatment. *Wound Repair Regen*. Nov 12 2015; 23(6):891-900. PMID 26297933
29. Campitiello F, Mancone M, Della Corte A, et al. To evaluate the efficacy of an acellular Flowable matrix in comparison with a wet dressing for the treatment of patients with diabetic foot ulcers: a randomized clinical trial. *Updates Surg*. Dec 2017; 69(4):523-529. PMID 28497218
30. Gould LJ, Orgill DP, Armstrong DG, et al. Improved healing of chronic diabetic foot wounds in a prospective randomised controlled multi-centre clinical trial with a microvascular tissue allograft. *Int Wound J*. May 2022; 19(4):811-825. PMID 34469077
31. Armstrong DG, Galiano RD, Orgill DP, et al. Multi-centre prospective randomised controlled clinical trial to evaluate a bioactive split thickness skin allograft vs standard of care in the treatment of diabetic foot ulcers. *Int Wound J*. May 2022; 19(4):932-944. PMID 35080127
32. Sanders L, Landsman AS, Landsman A, et al. A prospective, multicenter, randomized, controlled clinical trial comparing a bioengineered skin substitute to a human skin allograft. *Ostomy Wound Manage*. Sep 2014; 60(9):26-38. PMID 25211605

33. DiDomenico L, Landsman AR, Emch KJ, et al. A prospective comparison of diabetic foot ulcers treated with either a cryopreserved skin allograft or a bioengineered skin substitute. *Wounds*. Jul 2011; 23(7):184-189. PMID 25879172
34. Brigido SA, Boc SF, Lopez RC. Effective management of major lower extremity wounds using an acellular regenerative tissue matrix: a pilot study. *Orthopedics*. Jan 2004; 27(1 Suppl):S45-149. PMID 14763548
35. Reyzelman A, Crews RT, Moore JC, et al. Clinical effectiveness of an acellular dermal regenerative tissue matrix compared to standard wound management in healing diabetic foot ulcers: a prospective, randomised, multicentre study. *Int Wound J*. Jun 2009; 6(3):196-208. PMID 19368581
36. Reyzelman AM, Bazarov I. Human acellular dermal wound matrix for treatment of DFU: literature review and analysis. *J Wound Care*. Mar 2015; 24(3):128; 129-134. PMID 25764957
37. Brigido SA. The use of an acellular dermal regenerative tissue matrix in the treatment of lower extremity wounds: a prospective 16-week pilot study. *Int Wound J*. Sep 2006; 3(3):181-187. PMID 16984575
38. Walters J, Cazzell S, Pham H, et al. Healing rates in a multicenter assessment of a sterile, room temperature, acellular dermal matrix versus conventional care wound management and an active comparator in the treatment of full-thickness diabetic foot ulcers. *Eplasty*. Mar 2016; 16:e10. PMID 26933467
39. Cazzell S, Vayser D, Pham H, et al. A randomized clinical trial of a human acellular dermal matrix demonstrated superior healing rates for chronic diabetic foot ulcers over conventional care and an active acellular dermal matrix comparator. *Wound Repair Regen*. May 2017; 25(3):483-497. PMID 28544150
40. Frykberg RG, Cazzell SM, Arroyo-Rivera J, et al. Evaluation of tissue engineering products for the management of neuropathic diabetic foot ulcers: an interim analysis. *J Wound Care*. Jul 2016; 25 Suppl 7:S18-25. PMID 27410467
41. Lantis JC, Snyder R, Reyzelman AM, et al. Fetal bovine acellular dermal matrix for the closure of diabetic foot ulcers: a prospective randomised controlled trial. *J Wound Care*. Jul 01 2021; 30(Sup7):S18-S27. PMID 34256588
42. Niezgodna JA, Van Gils CC, Frykberg RG, et al. Randomized clinical trial comparing OASIS Wound Matrix to Regranex Gel for diabetic ulcers. *Adv Skin Wound Care*. Jun 2005; 18(5 Pt 1):258-266. PMID 15942317
43. Uccioli L, Giurato L, Ruotolo V, et al. Two-step autologous grafting using HYAFF scaffolds in treating difficult diabetic foot ulcers: results of a multicenter, randomized controlled clinical trial with long-term follow-up. *The international journal of lower extremity wounds*. Jun 2011; 10(2):80-85. PMID 21693443
44. Lullove EJ, Liden B, Winters C, et al. A Multicenter, Blinded, Randomized Controlled Clinical Trial Evaluating the Effect of Omega-3-Rich Fish Skin in the Treatment of Chronic, Nonresponsive Diabetic Foot Ulcers. *Wounds*. Jul 2021; 33(7):169-177. PMID 33872197
45. Lullove EJ, Liden B, McEneaney P, et al. Evaluating the effect of omega-3-rich fish skin in the treatment of chronic, nonresponsive diabetic foot ulcers: penultimate analysis of a multicenter, prospective, randomized controlled trial. *Wounds*. Apr 2022; 34(4):E34-E36. PMID 35797557

46. Lantis Li JC, Lullove EJ, Liden B, et al. Final efficacy and cost analysis of a fish skin graft vs standard of care in the management of chronic diabetic foot ulcers: a prospective, multicenter, randomized controlled clinical trial. *Wounds*. Apr 2023; 35(4):71-79. PMID 37023475
47. O'Meara S, Cullum N, Nelson EA et al. Compression for venous leg ulcers. *Cochrane Database Syst Rev*, 2012 Nov 16; 11:CD000265. PMID 23152202
48. Falanga V, Margolis D, Alvarez O, et al. Rapid healing of venous ulcers and lack of clinical rejection with an allogeneic cultured human skin equivalent. Human Skin Equivalent Investigators Group. *Arch Dermatol*. Mar 1998; 134(3):293-300. PMID 9521027
49. Mostow EN, Haraway GD, Dalsing M, et al. Effectiveness of an extracellular matrix graft (OASIS Wound Matrix) in the treatment of chronic leg ulcers: a randomized clinical trial. *J Vasc Surg*. May 2005; 41(5):837-843. PMID 15886669
50. Romanelli M, Dini V, Bertone M, et al. OASIS wound matrix versus Hyaloskin in the treatment of difficult-to-heal wounds of mixed arterial/venous aetiology. *Int Wound J*. Mar 2007; 4(1):3-7. PMID 17425543
51. Romanelli M, Dini V, Bertone MS. Randomized comparison of OASIS wound matrix versus moist wound dressing in the treatment of difficult-to-heal wounds of mixed arterial/venous etiology. *Adv Skin Wound Care*. Jan 2010; 23(1):34-38. PMID 20101114
52. Harding K, Sumner M, Cardinal M. A prospective, multicentre, randomised controlled study of human fibroblast-derived dermal substitute (Dermagraft) in patients with venous leg ulcers. *Int Wound J*. Apr 2013; 10(2):132-137. PMID 23506344
53. Cazzell S. A randomized controlled trial comparing a human acellular dermal matrix versus conventional care for the treatment of venous leg ulcers. *Wounds*. Mar 2019; 31(3):68-74. PMID 30720443
54. Carsin H, Ainaud P, Le Bever H, et al. Cultured epithelial autografts in extensive burn coverage of severely traumatized patients: a five-year single-center experience with 30 patients. *Burns*. Jun 2000; 26(4):379-387. PMID 10751706
55. Lagus H, Sarlomo-Rikala M, Bohling T, et al. Prospective study on burns treated with Integra(R), a cellulose sponge and split-thickness skin graft: comparative clinical and histological study--randomized controlled trial. *Burns*. Dec 2013; 39(8):1577-1587. PMID 23880091
56. Branski LK, Herndon DN, Pereira C, et al. Longitudinal assessment of Integra in primary burn management: a randomized pediatric clinical trial. *Crit Care Med*. Nov 2007; 35(11):2615-2623. PMID 17828040
57. Heimbach DM, Warden GD, Luterman A, et al. Multicenter postapproval clinical trial of Integra dermal regeneration template for burn treatment. *J Burn Care Rehabil*. Jan-Feb 2003; 24(1):42-48. PMID 12543990
58. Hicks KE, Huynh MN, Jeschke M, et al. Dermal regenerative matrix use in burn patients: A systematic review. *J Plast Reconstr Aesthet Surg*. Nov 2019; 72(11):1741-1751. PMID 31492583
59. Gonzalez SR, Wolter KG, Yuen JC. Infectious Complications Associated with the Use of Integra: A Systematic Review of the Literature. *Plast Reconstr Surg Glob Open*. Jul 2020; 8(7):e2869. PMID 32802634

60. Luze H, Nischwitz SP, Smolle C, et al. The Use of Acellular Fish Skin Grafts in Burn Wound Management-A Systematic Review. *Medicina (Kaunas)*. Jul 09 2022; 58(7). PMID 35888631
61. Holmes JH, Molnar JA, Shupp JW, et al. Demonstration of the safety and effectiveness of the RECELL[®] System combined with split-thickness meshed autografts for the reduction of donor skin to treat mixed-depth burn injuries. *Burns*. Jun 2019; 45(4):772-782. PMID 30578048
62. Holmes Iv JH, Molnar JA, Carter JE, et al. A Comparative Study of the ReCell[®] Device and Autologous Spit-Thickness Meshed Skin Graft in the Treatment of Acute Burn Injuries. *J Burn Care Res*. Aug 17 2018; 39(5):694-702. PMID 29800234
63. Fivenson DP, Scherschun L, Cohen LV. Apligraf in the treatment of severe mitten deformity associated with recessive dystrophic epidermolysis bullosa. *Plast Reconstr Surg*. Aug 2003; 112(2):584-588. PMID 12900618
64. Baldursson BT, Kjartansson H, Konradsdottir F, et al. Healing rate and autoimmune safety of full-thickness wounds treated with fish skin acellular dermal matrix versus porcine small intestine submucosa: a noninferiority study. *Int J Low Extrem Wounds*. Mar 2015; 14(1):37-43. PMID 25759413
65. Still J, Glat P, Silverstein P, et al. The use of a collagen sponge/living cell composite material to treat donor sites in burn patients. *Burns*. Dec 2003; 29(8):837-841. PMID 14636761
66. Brown-Etris M, Milne CT, Hodde JP. An extracellular matrix graft (Oasis wound matrix) for treating full-thickness pressure ulcers: A randomized clinical trial. *J Tissue Viability*, 2018 Dec 5; 28(1):21-26. PMID 30509850
67. Gurtner GC, Garcia AD, Bakewell K, et al. A retrospective matched-cohort study of 3994 lower extremity wounds of multiple etiologies across 644 institutions comparing a bioactive human skin allograft, TheraSkin, plus standard of care, to standard of care alone. *Int Wound J*. Feb 2020; 17(1):55-64. PMID 31729833
68. Lazic T, Falanga V. Bioengineered skin constructs and their use in wound healing. *Plast Reconstr Surg*. Jan 2011; 127 Suppl 1:75S-90S. PMID 21200276
69. Saffle JR. Closure of the excised burn wound: temporary skin substitutes. *Clin Plast Surg*. Oct 2009; 36(4):627-641. PMID 19793557
70. National Institute for Health and Care Excellence (NICE). Diabetic Foot Problems: Prevention and Management [NG19]. 2023. Available at: <<https://www.nice.org.uk>> (accessed April 19, 2024).
71. Peirce SC, Carolan-Rees G. ReCell[®] Spray-On Skin System for Treating Skin Loss, Scarring and Depigmentation after Burn Injury: A NICE Medical Technology Guidance. *Appl Health Econ Health Policy*. Apr 2019; 17(2):131-141. PMID 30635844

Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been changed since this medical policy document was written. See Medicare's National Coverage at <<http://www.cms.hhs.gov>>.

Policy History/Revision	
Date	Description of Change
07/01/2024	Document updated with literature review. The following changes were made to coverage: 1) mVASC and TheraSkin added to medically necessary statement for diabetic lower-extremity ulcers; 2) Artacent Wound, DeNovoSkin, SimpliDerm and Tutomesh Fenestrated Bovine Pericardium added to example list of products considered experimental, investigational and/or unproven. References 10, 30-31 and 46 added; others removed.
04/15/2024	Document updated with literature review. The following change was made to the example list of products considered experimental, investigational and/or unproven under Coverage: Recell® and OvaTex added. References 53, 67-69 and 77 added; others updated.
10/01/2022	Document updated. The following changes were made to the example list of products considered experimental, investigational and/or unproven under Coverage: Added Omeza® Collagen Matrix, Phoenix Wound Matrix, PermeaDerm B, PermeaDerm Glove, and PermeaDerm C. No new references added.
06/01/2022	Document updated with literature review. The following changes were made to products list under the experimental, investigational and/or unproven list under Coverage: added BellaCell HD or Surederm; Matrix HD™; Micromatrix®; Microlyte® Matrix; Mirragen® Advanced Wound Matrix, Skin substitute, FDA-cleared as a device, not otherwise specified, Innovamatrix FS, SUPRATHEL, and Supra SDRM. References 4, 47 and 52 added.
07/15/2021	Document updated with literature review. The following changes were made to products list under the experimental, investigational and/or unproven list under Coverage: added Bio-ConneKt® Wound Matrix; Coll-e-derm; Dermagine; Duragen® XS, Duragen™ Plus; FlowerDerm™; Geistlich Derma-Gine™; Integra® Matrix Wound Dressing (previously Avagen); InteguPly®; Keroxx™; MatriStem; Microderm®; MyOwn skin; Ologen™ Collagen Matrix; Omega3 Wound (originally Merigen wound dressing); Progenamatrix; Puracol® and Puracol® Plus Collagen Wound Dressings; and Skin TE™. References 1, 14, 22, 42, 61, and 62 added.
07/15/2020	Document updated with literature review. Coverage unchanged. Rationale and references revised; some references removed; added references 1, 2, 46, 52, and 62.
08/01/2019	New medical document. AlloDerm®, AlloMend®, Cortiva® [AlloMax™], DermACELL™, DermaMatrix™, FlexHD®, FlexHD® Pliable™, Graftjacket® may be considered medically necessary for breast reconstruction surgery when criteria are met. AlloPatch®, Apligraf®, Dermagraft®, Integra® Omnigraft™

Dermal Regeneration Matrix (also known as Omnigraft™) or, Integra Flowable Wound Matrix may be considered medically necessary in the treatment of chronic, non-infected, full-thickness diabetic lower-extremity ulcers. Apligraf® or Oasis™ Wound Matrix may be considered medically necessary for the treatment of chronic, non-infected, partial- or full-thickness lower-extremity skin ulcers due to venous insufficiency which have not adequately responded to a 1-month period of conventional ulcer therapy. Orcel™ may be considered medically necessary for the treatment of dystrophic epidermolysis bullosa when standard wound therapy has failed and in accordance with the humanitarian device exemption specification of the U.S. Food and Drug Administration. Epicel® may be considered medically necessary in the treatment of deep dermal or full-thickness burns comprising a total body surface area of $\geq 30\%$ when provided in accordance with the humanitarian device exemptions specifications of the FDA. Integra® Dermal Regeneration Template may be considered medically necessary for the treatment of second-and third-degree burns. All other uses of the bioengineered skin and soft tissue substitutes listed in the coverage of this policy are considered experimental, investigational and/or unproven. All other skin and soft tissue substitutes not listed are considered experimental, investigational and/or unproven.