

| | |
|------------------------------|-------------------|
| Policy Number | SUR705.029 |
| Policy Effective Date | 08/15/2025 |

Surgical Treatment of Femoroacetabular Impingement

| Table of Contents | Related Policies (if applicable) |
|-----------------------------------|----------------------------------|
| Coverage | None |
| Policy Guidelines | |
| Description | |
| Rationale | |
| Coding | |
| References | |
| Policy History | |

Disclaimer

Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

Coverage

NOTE 1: This medical policy does not apply to surgical treatment when FAI is absent.

Open or arthroscopic treatment of femoroacetabular impingement (FAI) **may be medically necessary** when **ALL** of the following conditions have been met:

1. Age
 - Candidates should be skeletally mature with documented closure of growth plates (e.g., ≥ 15 years of age).
2. Symptoms
 - Moderate-to-severe hip-related pain worsened by flexion activities (e.g., squatting or prolonged sitting) that significantly limits activities; AND
 - Unresponsive to conservative therapy for at least 3 months (e.g., nonsteroidal anti-inflammatory drugs [NSAIDs], physical therapy, activity modifications, restriction of athletic pursuits, and avoidance of symptomatic motion); AND
 - Positive impingement sign on clinical examination (pain elicited with 90° of flexion and internal rotation and adduction of the femur).

3. Imaging

- Morphology indicative of cam or pincer FAI (e.g., pistol-grip deformity, femoral head-neck offset with an alpha angle >50°, a positive wall sign, acetabular retroversion [over-coverage with crossover sign]), coxa profunda or protrusion, or damage of the acetabular rim; AND
- High probability of a causal association between the FAI morphology and damage (e.g., a pistol-grip deformity with a tear of the acetabular labrum and articular cartilage damage in the anterosuperior quadrant); AND
- No evidence of advanced osteoarthritis, defined as Tönnis grade 2 or 3, or joint space of less than 2 mm; AND
- No evidence of severe (Outerbridge grade IV) chondral damage.

Treatment of FAI is considered experimental, investigational, and/or unproven in all other situations.

Policy Guidelines

If femoroacetabular impingement morphology is identified, individuals should be advised not to play aggressive sports. No more frequent than annual follow-up with magnetic resonance arthrography may be indicated for femoroacetabular impingement morphology to evaluate cartilage changes before damage becomes severe. It should be noted that current imaging techniques limit the early identification of cartilage defects, whereas delay in the surgical correction of bony abnormalities may lead to disease progression to the point at which joint preservation is no longer appropriate. Confirmation of subtle femoroacetabular impingement morphology may require 3-dimensional computed tomography. Some clinicians may also use local anesthetic injection into the joint to assist in confirming femoroacetabular impingement pathology.

Treatment of femoroacetabular impingement should be restricted to centers experienced in treating this condition and staffed by surgeons adequately trained in techniques addressing femoroacetabular impingement. Because of the differing benefits and risks of open and arthroscopic approaches, individuals should make an informed choice between the procedures.

Some individuals may require a revision procedure if symptoms recur or persist. Published studies have indicated that all sources of impingement might not have been identified before surgery, and those that had might not have been adequately treated. The risk of additional surgical procedures can be reduced by intraoperative assessment of impingement after bone debridement and reshaping.

There are specific CPT codes for arthroscopic treatment of femoroacetabular impingement (CPT 29914-29916), and these codes should be used accordingly. CPT codes 29862 and 29863 are for arthroscopic surgery on the articular cartilage, labrum, and/or synovium of the hip joint,

not for repair of femoroacetabular impingement pathology and, therefore, should not be used to report arthroscopic femoroacetabular impingement surgery.

There are no specific CPT codes for the open treatment of femoroacetabular impingement. The procedure might be coded using code 27299 (unlisted procedure, pelvis or hip joint).

Description

Femoroacetabular Impingement

Femoroacetabular impingement arises from an anatomic mismatch between the head of the femur and the acetabulum, causing compression of the labrum or articular cartilage during flexion. The mismatch can arise from subtle morphologic alterations in the anatomy or orientation of the ball and socket components (e.g., a bony prominence at the head-neck junction or acetabular over-coverage), with articular cartilage damage initially occurring from abutment of the femoral neck against the acetabular rim, typically at the anterosuperior aspect of the acetabulum. Although hip joints can possess the morphologic features of femoroacetabular impingement without symptoms, femoroacetabular impingement may become pathologic with repetitive movement and/or increased force on the hip joint. High-demand activities may also result in pathologic impingement in hips with normal morphology.

Two types of impingement, cam and pincer, may occur alone or, more frequently, together. Cam impingement is associated with an asymmetric or nonspherical contour of the head or neck of the femur jamming against the acetabulum, resulting in cartilage damage and delamination (detachment from the subchondral bone). Deformity of the head/neck junction that looks like a pistol grip on radiographs is associated with damage to the anterosuperior area of the acetabulum. Symptomatic cam impingement is found most frequently in young male athletes. Pincer impingement is associated with over coverage of the acetabulum and pinching of the labrum, with pain more typically beginning in women of middle age. In cases of isolated pincer impingement, the damage may be limited to a narrow strip of the acetabular cartilage.

Epidemiologic and radiographic studies have found correlations between hip osteoarthritis and femoroacetabular impingement lesions, supporting the theory that prolonged contact between the anatomically mismatched acetabulum and femur may lead not only to cam and pincer lesions, but eventually to further cartilage damage and subsequent joint deterioration. It is believed that osteoplasty of the impinging bone is needed to protect the cartilage from further damage and to preserve the natural joint. Therefore, if femoroacetabular impingement morphology is shown to be an etiology of osteoarthritis, a strategy to reduce the occurrence of idiopathic hip osteoarthritis could be early recognition and treatment of femoroacetabular impingement before cartilage damage and joint deterioration occurs.

An association between femoroacetabular impingement and athletic pubalgia, sometimes called sports hernia, has been proposed. Athletic pubalgia is an umbrella term for a large variety of musculoskeletal injuries involving attachments and/or soft tissue support structures

of the pubis, and is not addressed on this policy (see medical policy SUR705.036 on the surgical treatment of athletic pubalgia).

Treatment

A technique for hip dislocation with open osteochondroplasty that preserved the femoral blood supply was reported by Ganz. Visualization of the entire joint with this procedure led to the identification and acceptance of femoroacetabular impingement as an etiology of cartilage damage and the possibility of correcting the abnormal femoroacetabular morphology. Open osteochondroplasty of bony abnormalities and treatment of the symptomatic cartilage defect is considered the criterion standard for complex bony abnormalities. However, open osteochondroplasty is invasive, requiring transection of the greater trochanter (separation of the femoral head from the femoral shaft) and dislocation of the hip joint to provide full access to the femoral head and acetabulum. In addition to the general adverse effects of open surgical procedures, open osteochondroplasty with dislocation has been associated with nonunion and neurologic and soft tissue lesions.

Less invasive hip arthroscopy and an arthroscopy-assisted mini-approach were developed in 2004. Arthroscopy requires specially designed instruments and is considered technically more difficult due to reduced visibility and limited access to the joint space. Advanced imaging techniques, including computed tomography and fluoroscopy, have been used to improve visualization of the 3-dimensional head/neck morphology during arthroscopy.

Femoroacetabular Impingement can also be a source of hip-related pain and decreased hip internal rotation in the pediatric population. When nonoperative management of femoroacetabular impingement in children and adolescents is ineffective, surgical procedures may be indicated. Surgical techniques include arthroscopy, open hip dislocation, limited open with arthroscopy, and osteotomy.

Slipped Capital Femoral Epiphysis

Patients with slipped capital femoral epiphysis have a displaced femoral head in relation to the femoral neck within the confines of the acetabulum, which can result in hip pain, thigh pain, knee pain, and onset of a limp. Slipped capital femoral epiphysis occurs most frequently in children between the ages of 10 to 16. Upon reaching skeletal maturity, patients diagnosed with slipped capital femoral epiphysis, 32% were found to have clinical signs of impingement. It is not uncommon for patients with slipped capital femoral epiphysis to develop premature osteoarthritis and require total hip arthroplasty within 20 years.

Treatment

The standard treatment for slipped capital femoral epiphysis is stabilization across the physis by in situ pinning. Alternative treatments proposed for pediatric patients with slipped capital femoral epiphysis related femoroacetabular impingement include osteoplasty without dislocation, or with the open dislocation technique described by Ganz. The Ganz technique (capital realignment with open dislocation) is technically demanding, with a steep learning

curve and a high-risk of complications, including avascular necrosis. Therefore, early treatment to decrease impingement must be weighed against increased risk of adverse events.

Regulatory Status

Surgery for the treatment of femoroacetabular impingement is a procedure and, as such, is not subject to regulation by the United States (U.S.) Food and Drug Administration (FDA).

Rationale

Medical policies assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, two domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Adults with Asymptomatic Femoroacetabular Impingement

Clinical Context and Therapy Purpose

The purpose of femoroacetabular impingement surgery is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as observation, in adults with asymptomatic femoroacetabular impingement.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals who are adults with asymptomatic femoroacetabular impingement.

Femoroacetabular impingement is more prevalent in the athletic population. The true prevalence of femoroacetabular impingement is difficult to obtain as there is disagreement

whether it should be assessed on the basis of radiographic or clinical findings suggestive of femoroacetabular impingement, as a large percentage of individuals in both the athletic and general population may have radiographic findings indicative of femoroacetabular impingement but remains clinically asymptomatic. (1) Frank et al. conducted a systematic review of 26 studies to determine the prevalence of femoroacetabular impingement in asymptomatic individuals (N=2114). (2) The prevalence of cam-type deformity was 23.1% in the general population, compared with 54.8% in the athletic population.

Interventions

The therapy being considered is femoroacetabular impingement surgery. Femoroacetabular impingement results from localized compression in the joint due to an anatomic mismatch between the head of the femur and the acetabulum. Symptoms of impingement typically occur in young to middle-aged adults before the onset of osteoarthritis but may be present in younger individuals with developmental hip disorders.

The objective of surgical treatment of femoroacetabular impingement is to provide symptom relief and reduce further joint damage.

Comparators

Comparators of interest include observation.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, health status measures, quality of life, and change in disease status.

The existing literature evaluating femoroacetabular impingement surgery as a treatment for patients who are adults with asymptomatic femoroacetabular impingement has varying lengths of follow up. At least 1 year of follow-up is desirable to assess outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Currently, there are no studies providing evidence on the efficacy of femoroacetabular impingement for adults with asymptomatic femoroacetabular impingement morphology for the prevention of osteoarthritis. Indirect evidence consists of observational studies that demonstrate a relation between femoroacetabular impingement and osteoarthritis.

Observational Studies

Oner et al. (2016) conducted a retrospective study to determine the prevalence of femoroacetabular impingement as an etiologic factor for osteoarthritis in the hip joint among patients who had undergone total hip arthroscopy (THA). (3) Radiographs of 1,004 patients who had undergone THA between 2005 and 2010 were reviewed by 3 authors. Intra- and interobserver consistencies were calculated. The predisposing etiologic factor leading to end-stage degenerative hip disease was undetermined in 26 of the radiographs. Among the remaining 978 patients, 99 patients were diagnosed with femoroacetabular impingement by all 3 reviewers, 83 with a cam-type femoroacetabular impingement, and 16 with pincer-type femoroacetabular impingement. Interobserver agreement was high, with a contingency coefficient between observers of 0.71 for the diagnosis of femoroacetabular impingement.

A population-based cohort study by Thomas et al. (2014) found that subclinical deformities of the hip, including cam-type femoroacetabular impingement, were significant predictors of radiographic evidence of osteoarthritis and total hip replacement in women. (4) A cohort of 1,003 women underwent pelvis radiographs at years 2 and 20. At 20 years, blinded radiographic analysis was available for 670 (46%) hips, of which 70 (11%) showed osteoarthritis. For total hip replacement, data at the 20-year assessment were available for 1,455 (99%) hips, of which 40 (3%) had undergone replacement. Pincer-type femoroacetabular impingement at year 2 was not significantly associated with radiographic evidence of osteoarthritis. Cam-type femoroacetabular impingement at year 2 was significantly associated with the development of radiographic osteoarthritis and total hip replacement. Each degree increased in the alpha angle above 65° was associated with an increased risk of 5% for radiographic evidence of osteoarthritis and 4% for total hip replacement. These findings were limited by the low rate of participants having both baseline and follow-up radiographs.

Ganz and colleagues began a cross-sectional population-based natural history cohort in 2005 that included over 1,000 young men to determine whether morphologic alterations are associated with an increased rate of early osteoarthritis. Reichenbach et al. (2011) reported on 1080 asymptomatic young men in the Sumiswald Cohort had undergone clinical examination and completed the Western Ontario and McMaster Universities Osteoarthritis Index and the EuroQoL health-related quality of life questionnaire. (5) Of these, 244 randomly selected subjects (mean age, 19.9 years) underwent magnetic resonance imaging to evaluate cam-type deformities, labral lesions, cartilage thickness, and impingement pits. Definite cam-type deformities were detected in 67 (27%) asymptomatic men. Logistic regression models, adjusted for age and body mass index, found for patients with cam-type deformities odds ratios of 2.77 for labral lesions, 2.91 for impingement pits, and 2.45 for labral deformities. Cartilage thickness was -0.19 mm lower in subjects with cam-type deformities.

Gosvig et al. (2010) published findings from a cross-sectional radiographic and questionnaire database of 4,151 individuals from the Copenhagen Osteoarthritis study. (6) The study group consisted of 1,332 men (mean age, 60.0 years; range, 22-90 years), and 2,288 women (mean age, 60.8 years; range, 21-90 years). The hips were categorized as being without malformations

or as having an abnormality consisting of a deep acetabular socket, a pistol-grip deformity, or a combination of the two based on radiographic criteria. Male and female prevalence of hip joint malformations was 71% and 36.6%, respectively. The prevalence of hip osteoarthritis, radiographically defined, was 9.5% in men and 11.2% in women. A deep acetabular socket or a pistol-grip deformity was a significant risk factor in the development of hip osteoarthritis (relative risk, 2.4 and 2.2, respectively).

A study by Takeyama et al. (2009) from Asia reviewed records of 817 patients (946 hips) who underwent primary surgery for osteoarthritis or other hip diseases to determine the prevalence of femoroacetabular impingement. (7) Most (73%) patients were diagnosed with osteoarthritis secondary to developmental hip dysplasia. Only 17 (1.8%) patients were considered to have had primary osteoarthritis. Of these, 6 patients (average age, 63 years; range, 32-79 years) were determined to have femoroacetabular impingement from preoperative radiographs, resulting in a possible etiology of femoroacetabular impingement for 0.6% of the total population undergoing surgery for osteoarthritis and 35% in the population with primary osteoarthritis.

Bardakos and Villar (2009) retrospectively examined 43 patients (43 hips) younger than 55 years of age with a history of symptomatic idiopathic arthritis, who exhibited pistol-grip deformity of the femur and mild-to-moderate osteoarthritis (Tönnis grade I or II). (8) Radiographs showed progression of osteoarthritis in two-thirds of the patients, with 12 receiving hip replacement or resurfacing after more than 10 years. Logistic regression analysis showed the medial proximal femoral angle and the posterior wall sign as significant independent predictors for progression of osteoarthritis. A reduction of 1° in the medial proximal angle increased the odds of osteoarthritis progression by 21 times, while hip osteoarthritis with a positive posterior wall sign was 10 times more likely to progress than a hip that had a negative posterior wall sign. Of note, one-third of the patients with a pistol-grip deformity did not progress rapidly within the assessment period.

Kim et al. (2007) reviewed outcomes for 43 patients (mean age, 40 years; range, 18-68 years) with labral tears and early osteoarthritis (Tönnis grade 0 to I; average Japanese Orthopedic Association scores, <1 [The Japanese Orthopedic Association scale ranges from 0 [severe pain] to 3 [no pain]] and symptoms lasting 3 months or more who had been treated with debridement]. (9) At an average 50-month follow-up (range, 12-96 months), 74% of patients reported symptom improvement. Blinded evaluation of preoperative radiographs and magnetic resonance arthrograms indicated 42% of patients had femoroacetabular impingement. Patients treated only with débridement were less likely to improve if early-stage osteoarthritis or femoroacetabular impingement was present at the time of surgery (Japanese Orthopedic Association score, 1.67). Patients without either femoroacetabular impingement or osteoarthritis scored 2.6 while patients with femoroacetabular impingement scored 1.83.

A frequently cited paper by Beck et al. (2005) with Ganz as coauthor has described the potential relation between hip morphology and acetabular damage. (10) In this report, 26 patients with pure pistol-grip deformity and 16 patients with isolated coxa profunda were identified from

302 hips treated for intra-articular pathology between 1996 and 2001. Among the 26 hips with isolated cam impingement on preoperative radiographs, all showed acetabular cartilage damage in the anterosuperior area of the acetabulum with separation between the acetabular cartilage and the labrum. In the 16 hips with isolated pincer impingement, the damage was located more circumferentially, usually including only a narrow strip of the acetabular cartilage. The report illustrated that, in carefully selected patients with early-stage osteoarthritis and well-defined hip configurations, a strong association exists between specific hip morphology and the pattern of cartilage damage.

Tanzer and Noiseux (2004) reported on 3 separate populations when investigating anterior hip impingement as a common etiology of hip disorders. (11) The 3 populations of interest were patients who had undergone hip arthroscopy for labral tears (n=38), patients who had undergone cheilectomy for anterior femoroacetabular impingement (n=10), and patients who had THA due to idiopathic arthritis (n=200). Radiographic findings showed a pistol-grip deformity in 97% of the patients with labral tears and 100% of the patients with idiopathic arthritis.

Section Summary: Adults With Asymptomatic Femoroacetabular Impingement

There is no direct evidence that performing femoroacetabular impingement surgery on asymptomatic adults with femoroacetabular impingement morphology will prevent the development of osteoarthritis. There is indirect evidence from retrospective studies that individuals with cam-type impingement related to a pistol-grip deformity will experience labral damage, which can lead to the subsequent development of osteoarthritis.

Adults with Symptomatic Femoroacetabular Impingement

Clinical Context and Therapy Purpose

The purpose of femoroacetabular impingement surgery is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conservative management in adults with symptomatic femoroacetabular impingement.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals who are adults with symptomatic femoroacetabular impingement. Symptoms of impingement typically occur in young to middle-aged adults before the onset of osteoarthritis and may be present in younger individuals with developmental hip disorders.

Interventions

The therapy being considered is femoroacetabular impingement surgery. Femoroacetabular impingement results from localized compression in the joint due to an anatomic mismatch between the head of the femur and the acetabulum. The objective of surgical treatment of femoroacetabular impingement is to provide symptom relief and reduce further joint damage.

Comparators

Comparators of interest include conservative management. Conservative management includes activity changes, non-steroidal anti-inflammatory medications, and physical therapy.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, health status measures, quality of life, and change in disease status.

The existing literature evaluating femoroacetabular impingement surgery as a treatment for patients who are adults with asymptomatic femoroacetabular impingement has varying lengths of follow up. At least 1 year of follow-up is desirable to assess outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Surgical options for the treatment of adults with symptomatic femoroacetabular impingement include: open, arthroscopic, mini-open, and mixed open/arthroscopic.

Surgery (nonspecific)

Systematic Reviews

A systematic review and meta-analysis by Reiman et al. (2018) evaluated surgery for the treatment of femoroacetabular impingement among athletes as reported in 35 studies (1634 athletes; 1828 hips). (12) Evidence was graded limited to moderate strength. Athletes were reported to return to sport at preinjury level post-intervention at a rate of only 74% (95% confidence interval [CI], 67% to 81%). The mean follow-up post-surgery was 28.1 ± 15.5 months. Professional athletes returned to sport at a higher level than collegiate athletes ($p=0.0002$). It is difficult to assess effects of surgery on athletic performance as only 14% of studies reported on athletic performance before and after surgery.

A Cochrane review by Wall et al. (2014) evaluated surgery for the treatment of femoroacetabular impingement, conducting a literature search for randomized and quasi-randomized trials assessing surgical intervention compared with placebo treatment, nonoperative treatment, or no treatment in adults with femoroacetabular impingement. (13) No studies met these inclusion criteria. Four ongoing studies were identified at the time of publication.

A systematic review by Harris et al. (2013) evaluating the treatment of femoroacetabular impingement included literature through April 2013, identifying 29 studies (N=2369 patients): 83% had level IV evidence (case series), 14% had level III (cohort), and 3.4% had level I (RCT). (14) An arthroscopic approach was used in 59% of studies. Study interventions included nonoperative treatment, arthroscopy, surgical open dislocation, mixed open/arthroscopic, and mini-open. Both Nonarthritic Hip Score and modified Harris Hip Score (mHHS) values improved significantly, regardless of the surgical procedure compared with nonsurgical management. Differences between surgical techniques could not be assessed due to heterogeneity across surgical groups and inconsistent outcome measures.

Open Surgery

Systematic Reviews

A systematic review by Bedi et al. (2008) evaluated the management of labral tears and femoroacetabular impingement. (15) Seven of the 19 studies assessed were case series of patients with femoroacetabular impingement treated with open hip dislocation.

Observational Studies

Espinosa et al. (2006) compared the effect of reattaching (n=35) or removing (n=25) the labrum during treatment for femoroacetabular impingement. (16) Patients were 20 to 40 years of age and had no prior surgery; all had preoperative evidence of acetabular damage. Independent evaluations at 2-year follow-up indicated improved Merle d'Aubigné scores for both groups. The study also reported a reduction in osteoarthritis progression.

Peters and Erickson (2006) reported on 29 patients (30 hips) in a prospective study with minimum 2-year follow-up. (17) The specific diagnoses were primary femoroacetabular impingement in 25 patients (26 hips), Legg-Calve-Perthes disease (n=3) and slipped capital femoral epiphysis (n=1). The average age of the patients was 31 years (range, 16-51 years). Twenty-nine of the 30 hips had cam-type impingement (n=14) or mixed cam and pincer-type impingement (n=15). The Harris Hip Score improved from 70 at baseline to 87 at an average 32-month follow-up. No progression to osteoarthritis has observed in 68% of patients. There was nonunion in 8 (27%) hips; 5 (17%) hips were expected to convert to THA due to progressive pain, and 4 (13%) had progressed to osteoarthritis. Radiographic signs of progression of osteoarthritis and clinical failure requiring conversion to THA were seen only in patients with severe damage to the acetabular articular cartilage.

Beck et al. (2004) reported on 19 of 22 patients (average age, 36 years; range, 21-52 years) with confirmed clinical, radiographic, and magnetic resonance arthrographic diagnosis of femoroacetabular impingement, treated with surgical dislocation of the hip. (18) Follow-up duration was at least 4 years. All had labral damage, and 18 had acetabular damage. Using the Merle d'Aubigné hip score, 13 of the hips were rated excellent to good and pain scores improved from 2.9 to 5.1. By the 4- to 5-year follow-up, 5 (26%) patients had undergone THA due to cartilage damage.

Arthroscopic Surgery

The evidence on arthroscopic surgery for femoroacetabular impingement consists of systematic reviews, RCTs, and stand-alone observational studies.

Systematic Reviews

A systematic review by Jan et al. (2023) evaluated 5-year outcomes and survival rate of hip arthroscopy for femoroacetabular impingement syndrome. (19) A total of 15 articles were included, and labral repair was the most commonly performed procedure. The survival rate of hip arthroscopy at 5 years was generally high, with ranges of 0.0% to 17.9% for conversion to total hip arthroplasty (duration, 28.8 to 87.1 months) and 1.3% to 26.7% for revision surgery (duration, 14.8 to 83.7 months). All studies reported statistically significant improvement ($p<.05$) in patient-reported outcomes at the 5-year time point, with the most frequent patient-reported outcome being the mHHS. The authors noted that a lack of prospective data limited the quality of this review and also prevented the meta-analysis of pooled data.

Lamo-Espinosa et al. (2023) conducted a systematic review and meta-analysis comparing the efficacy and outcomes of patients with femoroacetabular impingement treated with hip arthroscopy versus physiotherapy with joint lavage. (20) The study included 6 RCTs (N=839). Results show that arthroscopy was superior to physiotherapy with joint lavage at 6 and 12 months in terms of the International Hip Outcome Tool (iHOT)-33 score (mean difference [MD], 3.98 [95% CI, 0.19 to 7.77] and MD, 10.65 [95% CI, 6.54 to 4.76], respectively) and Hip Outcome Score-Activities of Daily Living (HOS-ADL) subscale score (MD, 5.19 [95% CI, 0.77 to 9.61] and MD, 8.09 [95% CI, 3.11 to 13.07], respectively). However, this superiority did not exceed the minimum clinically important difference for HOS-ADL. For the outcome of Hip Outcome Score-Sport subscale score (HOS-S) at 6 and 12 months, arthroscopy did not exceed the minimum clinically important difference at 6 or 12 months (numerical results were not provided). Concerning safety, hip arthroscopy increased the risk of additional surgery (OR, 11.11; 95% CI, 1.42 to 86.89), OA (OR, 6.18; 95% CI, 1.06 to 36.00), and numbness (OR, 73.73; 95% CI, 10.00 to 543.92), while there were no significant differences between groups in the rates of infection or nerve injury.

Dwyer et al. (2020) published a systematic review and meta-analysis comparing efficacy and outcomes of patients with femoroacetabular impingement treated with hip arthroscopy versus physical therapy alone. (21) Reviewers identified 3 RCTs that included 650 patients (323 randomized to surgery; 327 randomized to physical therapy) with a follow-up rate of 90% and mean duration of 11.5 months. Patients treated with arthroscopic surgery had improved scores on the International Hip Outcome Tool 33 compared with the nonoperative group (standardized mean difference, 3.46; 95% CI, 0.07 to 6.86; $p<0.05$). The degree of statistical heterogeneity for this result was low ($I^2 = 41\%$; $p=0.18$). Pooled analyses of other outcome measures were conducted due to reporting in 2 or less studies. Casartelli and coworkers (2021) subsequently published a systematic review and meta-analysis assessing the iHOT-33 pooled MD in the same 3 RCTs (10.9; 95% CI, 4.7 to 17.0) and noted that this difference exceeded the minimum clinically important difference of 10 points, favoring hip arthroscopy. (22)

Minkara et al. (2019) published a systematic review and meta-analysis analyzing risk factors and outcomes after patients with femoroacetabular impingement had undergone hip arthroscopy. (23) Reviewers identified 29 relevant articles that included 1,911 patients (1,981 hips). Reviewers conducted a meta-analysis assessing return to play, revision rate, surgical and nonsurgical complications, change in α -angle, intraoperative bone resection, and patient-reported outcome measures after hip arthroscopy in femoroacetabular impingement. However, all but 2 studies (1 RCT, 1 prospective cohort) in the meta-analysis were case series. Reviewers also sought to identify risk factors associated with intervention success and/or failure. The data on reoperation and complication rates are most relevant. The cumulative risk of reoperation after hip arthroscopy, including revision surgery or subsequent THA, was 5.5% (95% CI, 3.6% to 7.5%). For patients requiring a secondary procedure, 77% underwent THA, and 13% required revision arthroscopy. A single study was the source for 19% of patients requiring a second procedure, which assessed hip arthroscopy exclusively among patients who were 50 years of age and older (mean, 57 years; range, 50-77 years). The risk of clinically reported complications was 1.7% (95% CI, 0.9% to 2.5%) The most frequent complication was heterotopic ossification, followed by transient neurapraxia, typically of the lateral femoral cutaneous nerve and sciatic nerve.

Kierkegaard et al. (2017) published a systematic review and meta-analysis of patients with femoroacetabular impingement who had undergone hip arthroscopy. (24) Outcomes were pain, activities of daily living (ADLs), and sports function. Databases were searched through September 2015. Nineteen studies were included in the meta-analysis (15 case series, 3 cohorts, 1 RCT). The RCT by Krych et al. (2013) is described in the next section. (25) The total number of patients included in the 19 studies was 2,322 (mean age, 36 years; range, 18-57 years) and 42% were women. Weighted mean differences between pre- and postoperative outcomes were evaluated in the meta-analysis. Detectable pain reduction was achieved in less than 3 months and maintained through 5 years. Improved ADLs were evident between 3 and 6 months and maintained through at least 3 years of follow-up. Sports function improvements were detected between 6 and 12 months after arthroscopy and were maintained through follow-up over several years. Patients who received femoroacetabular impingement continued to have some pain postsurgery.

Randomized Controlled Trials

An RCT comparing arthroscopic labral repair with labral debridement was reported by Krych et al. (2013). (25) This nonblinded RCT included 36 females with pincer-type or combined-type femoroacetabular impingement. At a mean 32-month follow-up (range, 12-48 months), both treatment groups showed significant improvements in the Hip Outcome Score vs baseline. Compared with the debridement group, the arthroscopic repair group had better outcomes on the Hip Outcome Score ADLs scale (91.2 vs 80.9) and Hip Outcome Score sports scale (88.7 vs 76.3). Most patients in the arthroscopic repair group also rated their hip function as normal or nearly normal (94% vs 78%).

Palmer et al. (2025) conducted an assessor-blind RCT comparing arthroscopic surgery to physical therapy in patients (N=222) with femoroacetabular impingement. (26) The primary

outcomes were minimum joint space width (mJSW) and HOS-ADL. At 38 months follow up, there was no difference in mJSW between groups. HOS-ADL was higher in the arthroscopy group than in the physical therapy group (84.2 vs 74.2; difference, 8.9; 95% CI, 7.0 to 10.8).

Open Surgery versus Arthroscopic Surgery

Systematic Reviews

Zhang et al. (2016) published a systematic review of studies comparing the efficacy and safety of hip arthroscopy with open surgical dislocation for the treatment of femoroacetabular impingement. (27) Five comparative studies published through August 2016 were included, evaluating a total of 352 hips. All studies were considered good or high quality based on the Newcastle-Ottawa Scale. Length of follow-up among the studies ranged from 12 to 25 months. At the 3-month follow-up, patients undergoing open dislocation experienced significant improvements in alpha angle (-4.45; 95% CI, -8.22 to -0.67) compared with patients undergoing arthroscopy, while patients undergoing arthroscopy reported significantly better Nonarthritic Hip Score (16.58; 95% CI, 9.54 to 23.61) compared with patients undergoing open dislocation. At 12-month follow-up, Nonarthritic Hip Score remained significantly better in the arthroscopy group, though the mHHS and Hip Outcome Score scales for ADLs and sports were equivalent between groups. Complications were also similar between groups, though reoperation rates were significantly lower in patients undergoing arthroscopy (relative risk, 0.4; 95% CI, 0.17 to 0.95).

Nwachukwu et al. (2016) published a systematic review and meta-analysis comparing open with arthroscopic surgical techniques for the treatment of femoroacetabular impingement. (28) The literature search included studies published through October 2014, which had a mean follow-up of at least 3 years. Sixteen studies met inclusion criteria - 9 open surgical hip dislocation studies and 7 hip arthroscopy studies. Pooled cohort analyses were conducted on data from 600 hips with a mean follow-up of 58 months from the open surgery studies and 1,484 hips with a mean follow-up of 51 months from the arthroscopy studies. Conversion to THA was the outcome endpoint, with an overall survival rate of 93% for patients undergoing open surgery and 90.5% for patients undergoing arthroscopy ($p=0.06$). Scores on the 12-Item Short Form Health Survey were significantly better among patients undergoing arthroscopy. Direct comparisons of other outcomes were limited by outcome instrument heterogeneity. Both surgical techniques demonstrated favorable outcomes using their respective measuring systems.

Several other systematic reviews comparing open with arthroscopic surgery for femoroacetabular impingement have been identified. (29-31) Matsuda et al. (2011) included 18 level III or IV studies (controlled cohort or case series) with a minimum 1-year follow-up. (29) Selected were 6 studies on open surgical dislocation, 4 on mini-open procedures, and 8 arthroscopic studies. All 3 approaches were effective in reducing pain and improving function in short-term to mid-term studies. Open dislocation surgery had a comparatively higher major complication rate primarily because of trochanteric osteotomy-related issues. The mini-open method showed comparable efficacy but significant incidence of iatrogenic injury to the lateral femoral cutaneous nerve. Botser et al. (2011) included 26 level II to IV articles totaling 1,462

hips in 1,409 patients. (30) Of these, 900 hips were treated arthroscopically, 304 with the open dislocation method, and 258 by the mini-open method. Mean time from onset of symptoms to surgery was 28 months. Overall complication rates were 1.7% for the arthroscopic group, 9.2% for the open surgical dislocation group, and 16% for the combined approach group.

Mini-Open and Mixed Open/Arthroscopic Approaches

The evidence for mixed-open and open/arthroscopic approaches for the treatment of femoroacetabular impingement consists of observational studies. This technique permits direct visualization of the anterior femoral head-neck junction without dislocation.

Observational Studies

Wu et al. (2019) reported on a study of anterolateral mini-open and arthroscopic osteochondroplasty in the treatment of cam-type femoroacetabular impingement (36 patients; 39 hips). (32) The mean Harris Hip Score significantly improved from 61.1 to 84.2 (p<0.01). Due to progression of hip osteoarthritis, 9 hips (23%) underwent total hip arthroplasty (THA) at a mean of 22 months. A survival analysis with the endpoint of conversion to THA or any femoroacetabular impingement revision surgery indicated a 74.9% rate for hip preservation at 5-years.

A study of the mini-open surgical technique performed on 118 patients with femoroacetabular impingement was described by Chiron et al. (2012). (33) Fifty-eight percent had cam-type impingement, and 42% had mixed-type impingement. The average follow-up was 2.2 years. Nonarthritic Hip Score, internal rotation, and alpha angles significantly improved following surgery. Eight revisions were performed, 2 patients experienced residual pain and eventually underwent THA, and 2 progressed rapidly to osteoarthritis.

A mixed open/arthroscopic approach for the treatment of femoroacetabular impingement was reported by Laude et al. (2009) for 97 patients (100 hips). (34) The average age of patients was 33 years (range, 16-56 years). Ninety-one (94%) were available for follow-up at an average 58 months (range, 29-104 months). Scores on the Nonarthritic Hip Score improved from 55 at baseline to 84 at the last follow-up. One patient had a femoral neck fracture 3 weeks postoperatively, and 13 (14%) required revision due to persistent pain. Eleven (12%) hips required THA at a mean of 40 months (range, 5-75 months). The best results were observed in patients younger than 40 years with Tönnis grade 0.

Section Summary: Adults With Symptomatic Femoroacetabular Impingement

The evidence for the use of open dislocation for the treatment of adults with symptomatic femoroacetabular impingement consists of systematic reviews of observational studies. The evidence for the use of arthroscopy to treat adults with femoroacetabular impingement includes systematic reviews of RCTs, and 2 additional RCTs. Systematic reviews comparing open dislocation and arthroscopy have shown that both procedures successfully reduce pain and improve functional outcomes, with arthroscopy showing more favorable satisfaction ratings. Although the evidence is mostly observational, cumulatively studies have reported on thousands of individuals and outcomes have been positive.

Adolescents and Children with Symptomatic Femoroacetabular Impingement

Clinical Context and Therapy Purpose

The purpose of femoroacetabular impingement surgery is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conservative management, in children ≤15 years of age with symptomatic femoroacetabular impingement.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals who are children ≤15 years of age with symptomatic femoroacetabular impingement. Symptoms of impingement typically occur in young to middle-aged adults before the onset of osteoarthritis but may be present in younger individuals with developmental hip disorders.

Interventions

The therapy being considered is femoroacetabular impingement surgery. Femoroacetabular impingement results from localized compression in the joint due to an anatomic mismatch between the head of the femur and the acetabulum. The objective of surgical treatment of femoroacetabular impingement is to provide symptom relief and reduce further joint damage.

Comparators

Comparators of interest include conservative management. Conservative management includes activity changes, non-steroidal anti-inflammatory medications, and physical therapy.

Outcomes

The general outcomes of interest are symptoms functional outcomes, health status measures, quality of life, and change in disease status.

The existing literature evaluating femoroacetabular impingement surgery as a treatment for individuals who are children ≤15 years of age with symptomatic femoroacetabular impingement has varying lengths of follow up, ranging from 2 to 5 years. At least 1 year of follow-up is desirable to assess outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

The evidence for the surgical management of adolescents and children with symptomatic femoroacetabular impingement consists of a meta-analysis and observational studies.

Systematic Review/Meta-Analysis

Huang et al. (2022) published a systematic review and meta-analysis evaluating the efficacy of arthroscopic treatment for femoroacetabular impingement syndrome in adolescents. (35) Data from 832 hips in 753 patients (mean age, 16.2 years; mean follow-up, 40.8 months) were retrieved from 6 prospective studies and 7 retrospective studies with moderate to high methodological quality. Following hip arthroscopy, all patient- reported outcomes improved significantly ($p<.0001$ for all measures). Specifically, 13 studies reported a 26.3- point improvement in the mean weighted mHHS, 9 studies reported a 37.0-point improvement in the HOS-S subscale, 7 studies reported a 23.7-point improvement in the HOS-ADL subscale, 4 studies reported a 23.0- point improvement in the Nonarthritic Hip Score (NAHS), and 5 studies reported a 42.7-point decrease in Visual Analogue Scale (VAS) pain scores. The alpha angle decreased by 22.0° from 62.9° to 40.9° following arthroscopic surgery. Across all studies, complications were reported for 10 hips (1.2%), including 3 cases of transient pudendal nerve neuropraxias, 4 cases of numbness, 2 minor infections, and 1 portal site wound dehiscence. Revision surgery was reported for 28 hips (3.4%), of which 8 (28.6%) were caused by new injuries and 20 (71.4%) were attributed to unrelieved symptoms or complications.

Observational Studies

Guindani et al. (2017) published results from patients less than 18 years of age who were retrospectively identified as having undergone surgical dislocation for several indications at a single institution. (36) Among the 51 patients (53 hips) in the study, 18 (34%) hips had a diagnosis of femoroacetabular impingement. Patients with femoroacetabular impingement reported significant improvements in the following pre- and post-measurements: mHHS, Nonarthritic Hip Score, and 12-Item Short Form Health Survey. No significant improvements were found in sphericity deviation score or on alpha angles (both anteroposterior and Lauenstein views).

Nwachukwu et al. (2017) reviewed an institutional hip preservation registry of patients with femoroacetabular impingement who underwent hip arthroscopy. (37) The authors sought to define the minimal clinically important difference and the substantial clinical benefit for adolescents undergoing hip arthroscopy. Data from 47 adolescents (68.1% female; mean age, 16.5 years) were obtained on the patients' mHHS, the Hip Outcome Score, and the international Hip Outcome Tool. Overall adolescent patients reported a minimal clinically important difference for the various patient-related outcomes but not substantial clinical benefit. The authors discussed the potential limitations of patient-related outcomes for adolescents compared with adults. They noted that adolescents might have higher expectations, and greater physical activity demands that influence their scores.

Included in the Huang et al. (2022) meta-analysis was a multicenter prospective study by Tran et al. (2013) who assessed arthroscopic treatment for cam-type femoroacetabular

impingement in 34 skeletally immature adolescents with open growth plates (41 hips). (38) At a mean follow-up of 14 months (range, 1-2 years), mHHS improved from 77.39 to 94.15 and Nonarthritic Hip Score improved from 76.34 to 93.18. Return to full sporting activity was reported by 78% of patients. No complications (e.g., avascular necrosis, slipped capital femoral epiphysis, fracture, growth plate arrest) were observed.

Section Summary: Adolescents and Children With Symptomatic Femoroacetabular Impingement

The evidence consists of a meta-analysis and observational studies. All studies reported favorable outcomes in pain reduction and functional improvements, but all had relatively small sample sizes ranging between 11 and 116 hips per study. No serious adverse events were reported. Studies comparing surgery to conservative management in this population are lacking.

Children with Slipped Capital Femoral Epiphysis-Associated Femoroacetabular Impingement Clinical Context and Therapy Purpose

The purpose of femoroacetabular impingement surgery is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as surgical repair of slipped capital femoral epiphysis alone, in individuals who are children ≤ 15 years of age with slipped capital femoral epiphysis associated femoroacetabular impingement.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals who are children ≤ 15 years of age with slipped capital femoral epiphysis associated femoroacetabular impingement.

Interventions

The therapy being considered is femoroacetabular impingement surgery. Femoroacetabular impingement results from localized compression in the joint due to an anatomic mismatch between the head of the femur and the acetabulum. Symptoms of impingement typically occur in young to middle-aged adults before the onset of osteoarthritis but may be present in younger individuals with developmental hip disorders. The objective of surgical treatment of femoroacetabular impingement is to provide symptom relief and reduce further joint damage.

Comparators

Comparators of interest include surgical repair of slipped capital femoral epiphysis alone.

Outcomes

The general outcomes of interest are symptoms functional outcomes, health status measures, quality of life, and change in disease status. The existing literature evaluating femoroacetabular impingement surgery as a treatment for individuals who are children ≤ 15 years of age with slipped capital femoral epiphysis associated femoroacetabular impingement has varying lengths of follow up. At least 1 year of follow-up is desirable to assess outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

Oduwole et al. (2017) reviewed 15 case series identified in a literature search from 2005 to 2016 that reported on the efficacy of surgical management in patients with femoroacetabular impingement secondary to slipped capital femoral epiphysis. (39) A total of 261 patients (266 hips) underwent both arthroscopic and open procedures (arthroscopic osteochondroplasty, 85 patients [88 hips]; surgical hip dislocation, 131 patients [133 hips]; open osteotomy, 45 patients [45 hips]). Mean alpha angle corrections observed for arthroscopy were 32.14°; for surgical hip dislocation, 41.45°; and for open osteotomy, 6.0° (p<0.05). Surgical hip dislocation resulted in the most improved correction of the alpha angle.

Observational Studies

Guindani et al. (2017) published results for patients less than 18 years of age undergoing surgical dislocation for several indications. (36) Among the 51 patients (53 hips) in the study, 13 (24%) hips had the diagnosis of slipped capital femoral epiphysis. Mean age at surgery for the whole population was 14 years and mean follow-up was 3 years. Outcomes postsurgery differed by indication. Slipped capital femoral epiphysis patients reported significant improvements in the following pre- and post-measurements: Nonarthritic Hip Score and on alpha angles (both anteroposterior and Lauenstein views). No significant improvements were found in mHHS, 12-Item Short Form Health Survey, or sphericity deviation scores.

Sink et al. (2010) retrospectively reviewed data from 2 United States (U.S.) centers evaluating 36 patients (39 hips) with stable slipped capital femoral epiphysis who were treated with open surgical hip dislocation for chronic symptoms. (40) The average time between in situ pinning and surgical hip dislocation was 20 months (range, 6-48 months). Most patients had partial or complete relief of symptoms immediately after initial pinning followed by a recurrence of symptoms consistent with impingement. All but 1 patient had either a labral or a cartilage injury, with labral injury observed in 34 of 39 hips and cartilage injury in 33 of 39 hips; the average depth of cartilage damage was 5 mm (range, 2-10 mm). There was no correlation between slip severity or duration of symptoms and the type of cartilage injury.

Ziebarth et al. (2009) with Ganz as coauthor conducted a joint 2-center retrospective review that assessed data from their Swiss institution (n=30) and a children's hospital in Boston (n=10).

Follow-up was 1 to 8 years for patients between 9 and 18 years of age with moderate-to-severe slipped capital femoral epiphysis who were treated with surgical dislocation. (41) No patients from either institution developed osteonecrosis, infection, deep venous thrombosis, or nerve palsies. Three patients developed delayed unions; none developed nonunions. Five patients required additional surgery for heterotopic ossification (n=1), residual impingement (n=1), or breakage of screw or wire fixation (n=3). The short-term postoperative clinical outcomes were found to be near normal, with similar scores between the operative and nonoperative hips.

As reported by Spencer et al. (2006), the same U.S. institution evaluated 19 patients (age range, 12-43 years) who underwent femoral neck osteoplasty (n=13) or osteoplasty with intertrochanteric osteotomy (n=6) via Ganz-type surgical dislocation. (42) Of 12 patients with a history of slipped capital femoral epiphysis (age range, 12-38 years), 9 reported improved symptom control at 8 to 25-month follow-up. Of the 7 patients (age range, 17-43 years) without slipped capital femoral epiphysis who underwent open surgical dislocation for pistol-grip deformities, 5 reported worse symptoms or minimal relief. Outcomes for patients with a chondral flap were worse than for patients without a chondral flap.

Section Summary: Children with Slipped Capital Femoral Epiphysis-Associated Femoroacetabular Impingement

The evidence for the use of femoroacetabular impingement surgical management for children with slipped capital femoral epiphysis associated femoroacetabular impingement consists of a systematic review and observational studies. Currently, there is no method to determine which children with slipped capital femoral epiphysis will develop femoroacetabular impingement. While most patients experienced symptom relief following femoroacetabular impingement surgery, the surgery is invasive, and complications (e.g., delayed union) have been reported.

Revision Arthroscopic Surgery for Femoroacetabular Impingement

Clinical Context and Therapy Purpose

The purpose of revision arthroscopic surgery is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conservative management, in individuals with residual femoroacetabular impingement symptoms following primary surgery.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with residual femoroacetabular impingement symptoms following primary surgery.

Interventions

The therapy being considered is revision arthroscopic surgery. The objective of surgical treatment of femoroacetabular impingement is to provide symptom relief and reduce further joint damage.

Comparators

Comparators of interest include conservative management. Conservative management includes activity changes, non-steroidal anti-inflammatory medications, and physical therapy.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, health status measures, quality of life, and change in disease status.

The existing literature evaluating revision arthroscopic surgery as a treatment for residual femoroacetabular impingement symptoms following primary surgery has varying lengths of follow-up. While studies described below all reported at least 1 outcome of interest, longer follow-up was necessary to fully observe outcomes. At least 1 year of follow-up is desirable to adequately assess outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

O'Connor et al. (2020) published a systematic review and meta-analysis on revision hip arthroscopy, which included 15 studies for review identifying 4,765 hips in 4,316 patients. (43) The most common indication for revision surgery was inadequate bony resection during the index procedure. Meta-analysis indicated significant improvements of patient-reported outcomes from baseline to final follow-up after revision surgery based on the mHHS (increase of 17.20 points), the Hip Outcome Score-Activities of Daily Living (improvement by 13.98 points), and the Visual Analog Scale (decrease of 3.16 points). However, while patient-reported outcomes improve significantly after revision, they remain lower than those of patients undergoing primary hip arthroscopic surgery.

Sardana et al. (2015) published a systematic review on revision hip arthroscopy, considering articles published through July 2014. (44) Three prospective case-control studies and 3 retrospective chart reviews, providing information on 448 hips, were selected. The most common indications for revision surgery were residual femoroacetabular impingement, labral tears, and chondral lesions. The mean interval between index and revision procedures was 25.6 months (range, 20.5-36 months). Patients most often requiring revision surgery were women (60%) and younger patients (mean age, 33.4 years). Revision hip arthroscopy improved functional outcomes (33.6% improvement in Harris Hip Score) and pain relief. Reviewers noted that the studies were low-quality (level III and IV).

A systematic review by Cvetanovich et al. (2015) evaluated revision hip arthroscopy. (45) Reviewers included 5 studies, with a total of 348 revision hip arthroscopies. The mean age of patients was 31.4 years, and 60% were female. The mean interval between index and revision procedures was 27.8 months. The most common indication for revision surgery was residual femoroacetabular impingement (81%). At a mean of 22.4-month follow-up, revision hip arthroscopy resulted in improved functional outcomes, as measured by the Harris Hip Score (weighted mean difference, 56.8 preoperative vs 72.0; $p=0.01$), Nonarthritic Hip Score, Hip Outcome Score, and 12-Item Short Form Health Survey.

Section Summary: Revision Arthroscopic Surgery for Femoroacetabular Impingement

The evidence for revision arthroscopic surgery for individuals with residual femoroacetabular impingement symptoms consists of 3 systematic reviews of observational studies. The observational studies, although low-quality, showed consistently favorable functional outcomes following revision surgery. The evidence for revision arthroscopic surgery for children consists of one observational study. Results have shown that children receiving revision surgery have functional outcomes comparable to children receiving primary arthroscopic surgery.

Summary of Evidence

For individuals who are adults with asymptomatic femoroacetabular impingement who receive femoroacetabular impingement surgery, there is no direct evidence that the surgical treatment will prevent the development of osteoarthritis. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. Indirect evidence consists of observational studies. In retrospective studies of patients with osteoarthritis, the relevant outcomes were radiographic evidence of hip joint malformations. In prospective studies of patients with femoroacetabular impingement, the relevant outcome is progression to osteoarthritis. Several large observational studies (>1000 patients), as well as smaller studies, have shown radiographic evidence of relationships between abnormal hip morphology and the development of osteoarthritis. There have been no studies in which femoroacetabular impingement surgery was performed on patients with femoroacetabular impingement morphology but no symptoms. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are adults with symptomatic femoroacetabular impingement who receive femoroacetabular impingement surgery, the evidence includes mostly systematic reviews of large and small observational studies and systematic a small randomized controlled trial (RCTs). Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. Open hip dislocation surgery and arthroscopic surgery are the most common surgical techniques performed on patients with femoroacetabular impingement. Systematic reviews have evaluated open hip dislocation surgery and arthroscopic surgery, compared with no comparator, nonsurgical management, and other surgical techniques. Compared with nonsurgical management, all types of surgical techniques have resulted in significant improvements in functional outcomes, pain, and radiographic measurements. The reviews were limited when comparing surgical techniques with each other because patient

characteristics and outcome measurements were heterogeneous among studies. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are children 15 years of age or younger with symptomatic femoroacetabular impingement who receive femoroacetabular impingement surgery, the evidence includes a meta-analysis evaluating small observational studies and individual observational studies. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. While the studies reported reductions in pain and improvements in functional outcomes, the sample sizes were relatively small, ranging between 11-116 hips per study. Additionally, comparative studies were not identified. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are children 15 years of age or younger with slipped capital femoral epiphysis-associated femoroacetabular impingement who receive femoroacetabular impingement surgery, the evidence includes a systematic review and small observational studies (range, 19-51 patients). Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. While most patients experienced symptom relief following femoroacetabular impingement surgery, the surgery is invasive and complications (e.g., nonunions) were reported. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have residual femoroacetabular impingement symptoms following a primary surgery who receive revision arthroscopic surgery, the evidence includes systematic reviews of observational studies (>400 patients). Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. Though the studies were of low-quality, consistent improvements in functional outcomes, pain relief, and patient satisfaction were reported, in some cases beyond 3 years. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Practice Guidelines and Position Statements

American Academy of Orthopaedic Surgeons (AAOS)

In 2020, the AAOS published a consensus-based best practice guidelines checklist for preoperative, intraoperative, and postoperative hip arthroscopy considerations in patients with femoroacetabular impingement. (46)

The guidelines define conservative care treatment as a trial of rest, non-steroidal anti-inflammatory drugs, activity modification or restriction, and physical therapy - without concomitant use of opioids. Prior to completion of the full duration of conservative treatment, assessment of the following joint parameters is recommended: high alpha angle, low Tönnis grade, cam or combined impingement, large range of motion limitations with pain, high baseline mental health status, large cam ($>65^\circ$ alpha angle) or combined deformity in absence of osteoarthritis changes. A shorter duration of conservative treatment is permissible in

professional or out-of-season athletes, patients completing physical therapy with no or marginal improvement, high baseline mental health status, and/or successful surgery on the contralateral side. Contraindications for hip arthroscopy include joint space narrowing <2 mm along the sourcil or osteoarthritis, Tönnis grade 2 or higher, severe femoral retroversion or anteversion with gait abnormality, obesity hindering access, broken Shenton line, pain not localizing to the hip or out of proportion due to psychiatric issue, inclination Tönnis angle >13 to 15°, or failed arthroscopy with dysplastic features. Hypermobility (Beighton Hypermobility Score ≥5) is not considered a contraindication for hip arthroscopy.

National Institute for Health and Care Excellence (NICE)

In 2011, the NICE issued guidance on arthroscopic femoroacetabular impingement surgery for hip impingement syndrome. (47) The NICE considered the evidence on the efficacy of arthroscopic femoroacetabular impingement surgery for hip impingement syndrome to be adequate for symptom relief in the short and medium term.

The NICE (2011) guidance on open femoroacetabular surgery for hip impingement syndrome indicated that evidence for this procedure was adequate for symptom relief in the short and medium term. (48)

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this policy are listed in Table 1.

Table 1. Summary of Key Trials

| NCT Number | Trial Name | Planned Enrollment | Completion Date |
|--------------------------|--|--------------------|-----------------------|
| <i>Ongoing</i> | | | |
| NCT04243447 | Identification of Predictors for Clinical Outcomes in Femoroacetabular Impingement Surgery (DoD FAI-2) | 800 | Sep 2025 |
| NCT02692807 | Arthroscopic Surgical Procedures Versus Sham Surgery for Patients with Femoroacetabular Impingement and/or Labral Tears: A Multicenter, International, Double-Blinded, Randomized Controlled Trial (HIPARTI) | 140 | Dec 2035 |
| NCT05746533 ^a | A Comparison of Total Hip Arthroplasty and Hip Preservation Outcomes | 10,000 | Jun 2053 (recruiting) |

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member's benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

| | |
|--------------------|-----------------------------------|
| CPT Codes | 27299, 29914, 29915, 29916, 29999 |
| HCPCS Codes | None |

*Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.

References

1. Egger AC, Frangiamore S, Rosneck J. Femoroacetabular Impingement: A Review. *Sports Med Arthrosc Rev*. Dec 2016; 24(4):e53-e58. PMID 27811519
2. Frank JM, Harris JD, Erickson BJ, et al. Prevalence of Femoroacetabular Impingement Imaging Findings in Asymptomatic Volunteers: A Systematic Review. *Arthroscopy*. Jun 2015; 31(6):1199-1204. PMID 25636988
3. Oner A, Koksal A, Sofu H, et al. The prevalence of femoroacetabular impingement as an aetiological factor for end- stage degenerative osteoarthritis of the hip joint: analysis of 1,000 cases. *Hip Int*. Mar-Apr 2016; 26(2):164-168. PMID 26916653
4. Thomas GE, Palmer AJ, Batra RN, et al. Subclinical deformities of the hip are significant predictors of radiographic osteoarthritis and joint replacement in women. A 20 year longitudinal cohort study. *Osteoarthritis Cartilage*. Oct 2014; 22(10):1504-1510. PMID 25047637
5. Reichenbach S, Leunig M, Werlen S, et al. Association between cam-type deformities and magnetic resonance imaging-detected structural hip damage: a cross-sectional study in young men. *Arthritis Rheum*. Dec 2011; 63(12):4023-4030. PMID 21904996
6. Gosvig KK, Jacobsen S, Sonne-Holm S, et al. Prevalence of malformations of the hip joint and their relationship to sex, groin pain, and risk of osteoarthritis: a population-based survey. *J Bone Joint Surg Am*. May 2010; 92(5):1162-1169. PMID 20439662
7. Takeyama A, Naito M, Shiramizu K, et al. Prevalence of femoroacetabular impingement in Asian patients with osteoarthritis of the hip. *Int Orthop*. Oct 2009; 33(5):1229-1232. PMID 19277653
8. Bardakos NV, Villar RN. Predictors of progression of osteoarthritis in femoroacetabular impingement: a radiological study with a minimum of ten years follow-up. *J Bone Joint Surg Br*. Feb 2009; 91(2):162-169. PMID 19190047
9. Kim KC, Hwang DS, Lee CH, et al. Influence of femoroacetabular impingement on results of hip arthroscopy in patients with early osteoarthritis. *Clin Orthop Relat Res*. Mar 2007; 456:128-132. PMID 17106273
10. Beck M, Kalhor M, Leunig M, et al. Hip morphology influences the pattern of damage to the acetabular cartilage: femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br*. Jul 2005; 87(7):1012-1018. PMID 15972923

11. Tanzer M, Noiseux N. Osseous abnormalities and early osteoarthritis: the role of hip impingement. *Clin Orthop Relat Res.* Dec 2004; (429):170-177. PMID 15577483
12. Reiman MP, Peters S, Sylvain J, et al. Femoroacetabular impingement surgery allows 74% of athletes to return to the same competitive level of sports participation but their level of performance remains unreported: a systematic review with meta-analysis. *Br J Sports Med.* Aug 2018; 52(15):972-981. PMID 29581142
13. Wall PD, Brown JS, Parsons N, et al. Surgery for treating hip impingement (femoroacetabular impingement). *Cochrane Database Syst Rev.* Sep 8 2014; 9(9):CD010796. PMID 25198064
14. Harris JD, Erickson BJ, Bush-Joseph CA, et al. Treatment of femoroacetabular impingement: a systematic review. *Curr Rev Musculoskelet Med.* Sep 2013; 6(3):207-218. PMID 23743861
15. Bedi A, Chen N, Robertson W, et al. The management of labral tears and femoroacetabular impingement of the hip in the young, active patient. *Arthroscopy.* Oct 2008; 24(10):1135-1145. PMID 19028166
16. Espinosa N, Rothenfluh DA, Beck M, et al. Treatment of femoro-acetabular impingement: preliminary results of labral refixation. *J Bone Joint Surg Am.* May 2006; 88(5):925-935. PMID 16651565
17. Peters CL, Erickson JA. Treatment of femoro-acetabular impingement with surgical dislocation and debridement in young adults. *J Bone Joint Surg Am.* Aug 2006; 88(8):1735-1741. PMID 16882895
18. Beck M, Leunig M, Parvizi J, et al. Anterior femoroacetabular impingement: part II. Midterm results of surgical treatment. *Clin Orthop Relat Res.* Jan 2004; (418):67-73. PMID 15043095
19. Jan K, Fenn TW, Kaplan DJ, et al. Patients maintain clinically significant outcomes at 5-year follow-up after hip arthroscopy for femoroacetabular impingement syndrome: A systematic review. *Arthroscopy.* Aug 2023; 39(8):1869-1881.e1. PMID 37207920
20. Lamo-Espinosa JM, Mariscal G, Gómez-Álvarez J, et al. Efficacy and safety of arthroscopy in femoroacetabular impingement syndrome: a systematic review and meta-analysis of randomized clinical trials. *Sci Rep.* Oct 01 2023; 13(1):16493. PMID 37779117
21. Dwyer T, Whelan D, Shah PS, et al. Operative versus nonoperative treatment of femoroacetabular impingement syndrome: A Meta-analysis of Short-Term Outcomes. *Arthroscopy.* Jan 2020; 36(1):263-273. PMID 31864588
22. Casartelli NC, Valenzuela PL, Maffiuletti NA, et al. Effectiveness of hip arthroscopy on treatment of femoroacetabular impingement syndrome: A meta-analysis of randomized controlled trials. *Arthritis Care Res (Hoboken).* Aug 2021; 73(8):1140-1145. PMID 32339441
23. Minkara AA, Westermann RW, Rosneck J, et al. Systematic review and meta-analysis of outcomes after hip arthroscopy in femoroacetabular impingement. *Am J Sports Med.* Feb 2019; 47(2):488-500. PMID 29373805
24. Kierkegaard S, Langeskov-Christensen M, Lund B, et al. Pain, activities of daily living and sport function at different time points after hip arthroscopy in patients with femoroacetabular impingement: a systematic review with meta- analysis. *Br J Sports Med.* Apr 2017; 51(7):572-579. PMID 27845683
25. Krych AJ, Thompson M, Knutson Z, et al. Arthroscopic labral repair versus selective labral debridement in female patients with femoroacetabular impingement: a prospective randomized study. *Arthroscopy.* Jan 2013; 29(1):46-53. PMID 23276413

26. Palmer A, Fernquest S, Rombach I, et al. Medium-term results of arthroscopic hip surgery compared with physiotherapy and activity modification for the treatment of femoroacetabular impingement syndrome: a multi-centre randomised controlled trial. *Br J Sports Med.* Jan 02 2025; 59(2):109-117. PMID 39592214
27. Zhang D, Chen L, Wang G. Hip arthroscopy versus open surgical dislocation for femoroacetabular impingement: A systematic review and meta-analysis. *Medicine (Baltimore).* Oct 2016; 95(41):e5122. PMID 27741133
28. Nwachukwu BU, Rebolledo BJ, McCormick F, et al. Arthroscopic versus open treatment of femoroacetabular impingement: a systematic review of medium- to long-term outcomes. *Am J Sports Med.* Apr 2016; 44(4):1062-1068. PMID 26059179
29. Matsuda DK, Carlisle JC, Arthurs SC, et al. Comparative systematic review of the open dislocation, mini-open, and arthroscopic surgeries for femoroacetabular impingement. *Arthroscopy.* Feb 2011; 27(2):252-269. PMID 21266276
30. Botser IB, Smith TW, Jr., Nasser R, et al. Open surgical dislocation versus arthroscopy for femoroacetabular impingement: a comparison of clinical outcomes. *Arthroscopy.* Feb 2011; 27(2):270-278. PMID 21266277
31. Papalia R, Del Buono A, Franceschi F, et al. Femoroacetabular impingement syndrome management: arthroscopy or open surgery? *Int Orthop.* 2012 May; 36(5):903-914. PMID 22190060
32. Wu CT, Mahameed M, Lin PC, et al. Treatment of cam-type femoroacetabular impingement using anterolateral mini-open and arthroscopic osteochondroplasty. *J Orthop Surg Res.* Jul 2019; 14(1):222. PMID 31315654
33. Chiron P, Espie A, Reina N, et al. Surgery for femoroacetabular impingement using a minimally invasive anterolateral approach: analysis of 118 cases at 2.2-year follow-up. *Orthop Traumatol Surg Res.* Feb 2012; 98(1):30-38. PMID 22257764
34. Laude F, Sariali E, Nogier A. Femoroacetabular impingement treatment using arthroscopy and anterior approach. *Clin Orthop Relat Res.* Mar 2009; 467(3):747-752. PMID 19089524
35. Huang HJ, Zhou X, Huang ZG, et al. Arthroscopic treatment for femoroacetabular impingement syndrome in adolescents: A systematic review and meta-analysis. *Clin J Sport Med.* Nov 01 2022; 32(6):608-616. PMID 36315820
36. Guindani N, Eberhardt O, Wirth T, et al. Surgical dislocation for pediatric and adolescent hip deformity: clinical and radiographical results at 3 years follow-up. *Arch Orthop Trauma Surg.* Apr 2017; 137(4):471-479. PMID 28197752
37. Nwachukwu BU, Chang B, Kahlenberg CA, et al. Arthroscopic treatment of femoroacetabular impingement in adolescents provides clinically significant outcome improvement. *Arthroscopy.* Oct 2017; 33(10):1812-1818. PMID 28623078
38. Tran P, Pritchard M, O'Donnell J. Outcome of arthroscopic treatment for cam type femoroacetabular impingement in adolescents. *ANZ J Surg.* Sep 3 2013; 83(5):382-386. PMID 22943465
39. Oduwole KO, de Sa D, Kay J, et al. Surgical treatment of femoroacetabular impingement following slipped capital femoral epiphysis: A systematic review. *Bone Joint Res.* Aug 2017; 6(8):472-480. PMID 28790036

40. Sink EL, Zaltz I, Heare T, et al. Acetabular cartilage and labral damage observed during surgical hip dislocation for stable slipped capital femoral epiphysis. *J Pediatr Orthop.* Jan-Feb 2010; 30(1):26-30. PMID 20032738
41. Ziebarth K, Zilkens C, Spencer S, et al. Capital realignment for moderate and severe SCFE using a modified Dunn procedure. *Clin Orthop Relat Res.* Mar 2009; 467(3):704-716. PMID 19142692
42. Spencer S, Millis MB, Kim YJ. Early results of treatment of hip impingement syndrome in slipped capital femoral epiphysis and pistol grip deformity of the femoral head-neck junction using the surgical dislocation technique. *J Pediatr Orthop.* May-Jun 2006; 26(3):281-285. PMID 16670535
43. O'Connor M, Steinl GK, Padaki AS, et al. Outcomes of Revision Hip Arthroscopic Surgery: A Systematic Review and Meta-analysis. *Am J Sports Med.* Apr 2020; 48(5):1254-1262. PMID 31503501
44. Sardana V, Philippon MJ, de Sa D, et al. Revision hip arthroscopy indications and outcomes: a systematic review. *Arthroscopy.* Oct 2015; 31(10):2047-2055. PMID 26033461
45. Cvetanovich GL, Harris JD, Erickson BJ, et al. Revision hip arthroscopy: a systematic review of diagnoses, operative findings, and outcomes. *Arthroscopy.* Jul 2015; 31(7):1382-1390. PMID 25703289
46. Lynch TS, Minkara A, Aoki S, et al. Best practice guidelines for hip arthroscopy in femoroacetabular Impingement: Results of a Delphi Process. *J Am Acad Orthop Surg.* Jan 15, 2020; 28(2):81-89. PMID 31181030
47. National Institute for Health and Clinical Excellence (NICE). Arthroscopic femoro-acetabular surgery for hip impingement syndrome [IPG408]. 2011. Available at <<https://www.nice.org.uk>> (accessed May 13, 2025).
48. National Institute for Health and Care Excellence (NICE). Open femoro-acetabular surgery for hip impingement syndrome [IPG403]. 2011. Available at <<https://www.nice.org.uk>> (accessed May 13, 2025).

Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does not have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been developed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

Policy History/Revision

| Date | Description of Change |
|------|-----------------------|
|------|-----------------------|

| | |
|------------|--|
| 08/15/2025 | Document updated with literature review. The following changes were made in Coverage: 1) Added “nonsteroidal anti-inflammatory drugs, [NSAIDs], physical therapy” as examples of conservative therapy, 2) Removed the following procedures from this policy: Hip arthroscopy to repair degenerative labral tears (e.g., due to early osteoarthritis); Labrum reconstruction of the hip; Capsular plication; Autologous osteochondral mosaicplasty in combination with femoral neck osteochondroplasty. 3) Removed NOTE 2 defining labral reconstruction versus labral repair 4) Removed separate statement related to revision of open or arthroscopic surgical procedures as the same criteria would apply as the primary procedure. Added reference 26; others updated and/or removed. |
| 08/15/2024 | Document updated with literature review. The following changes were made in Coverage: 1) Updated term “patient” to “candidate”; 2) Added examples of mechanical symptoms associated with labral tears. 3) Added references 24, 25, 27, 39, 54, 59, 60, 62, 63; others updated; some removed. |
| 08/15/2023 | Reviewed. No changes. |
| 05/01/2022 | Document updated with literature review. Coverage unchanged. Added reference 56; others updated; some removed. |
| 11/01/2021 | Reviewed. No changes. |
| 10/01/2020 | Document updated with literature review. Coverage unchanged. Added references 17, 24, 25, 28, 37, 40, 43, 48, 51, 54, 55; some removed. |
| 08/01/2019 | Reviewed. No changes. |
| 08/01/2018 | Document updated with literature review. The following was added to Coverage: 1) Hip arthroscopy to repair a labral tear may be considered medically necessary for traumatic labral tears causing mechanical symptoms and as an adjunct to FAI surgery. 2) The following procedures are considered experimental investigational and/or unproven for the treatment of FAI: Hip arthroscopy to repair degenerative labral tears (e.g., due to early osteoarthritis); or Labrum reconstruction of the hip; or Capsular plication; or Autologous osteochondral mosaicplasty in combination with femoral neck osteochondroplasty. 3) Revision of open or arthroscopic surgical treatment of FAI may be medically necessary when ALL the following conditions have been met: The above criteria (section 1, 2, 3) are met, the patient has residual symptoms following a primary surgery and the patient remains unresponsive to conservative therapy for at least 3 months postoperatively (including activity modifications, restriction of athletic pursuits, and avoidance of symptomatic motion). 4) NOTE 2: Labral reconstruction uses a graft to reconstruct the native labrum. This is distinct from a labral repair, which is to repair the torn tissue by sewing it back together and/ or to its attachment site. Title changed from Femoroacetabular Impingement (FAI) Syndrome. Added references 1-6, 21, 32, 33, 39, 43, 47-55, 58. |

| | |
|------------|---|
| 09/15/2016 | Reviewed. No changes. |
| 10/15/2015 | Document updated with literature review. Coverage unchanged. |
| 10/15/2014 | Reviewed. No changes. |
| 07/15/2013 | Document updated with literature review. The following was changed in Coverage: Criteria for age now states "Patient should be skeletally mature with documented closure of growth plates (e.g., 15 years or older)". Also, "(Hip Impingement Syndrome)" was removed from the policy title. |
| 08/15/2011 | Document updated with literature review. Coverage unchanged. Rationale completely revised. |
| 08/15/2009 | Updated document with literature review. Revised coverage from "experimental, investigational and unproven" for all indications to "medically necessary" under specified conditions. |
| 07/15/2008 | New medical document |