

<b>Policy Number</b>	<b>SUR713.034</b>
<b>Policy Effective Date</b>	<b>02/01/2025</b>
<b>Policy End Date</b>	<b>12/31/2025</b>

## Aqueous Shunts and Stents for Glaucoma

<b>Table of Contents</b>
<a href="#"><u>Coverage</u></a>
<a href="#"><u>Policy Guidelines</u></a>
<a href="#"><u>Description</u></a>
<a href="#"><u>Rationale</u></a>
<a href="#"><u>Coding</u></a>
<a href="#"><u>References</u></a>
<a href="#"><u>Policy History</u></a>

<b>Related Policies (if applicable)</b>
None

### Disclaimer

#### **Carefully check state regulations and/or the member contract.**

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

### Coverage

#### **\*CAREFULLY CHECK STATE REGULATIONS AND/OR THE MEMBER CONTRACT\***

#### **Ab Externo Aqueous Shunts**

Insertion of ab externo aqueous shunts approved by the U.S. Food and Drug Administration **may be considered medically necessary** as a method to reduce intraocular pressure in individuals with glaucoma where medical therapy has failed to adequately control intraocular pressure.

Use of an ab externo aqueous shunt for all other conditions, including in individuals with glaucoma when intraocular pressure is adequately controlled by medications, **is considered experimental, investigational and/or unproven.**

#### **Ab Interno Aqueous Stents**

Insertion of ab interno aqueous stents approved by the U.S. Food and Drug Administration as a method to reduce intraocular pressure in individuals with glaucoma where medical therapy has failed to adequately control intraocular pressure **may be considered medically necessary**.

Implantation of 1 or 2 U.S. Food and Drug Administration approved ab interno stents in conjunction with cataract surgery **may be considered medically necessary** in individuals with mild-to-moderate open-angle glaucoma treated with ocular hypotensive medication.

**NOTE 1:** Refer to the Description Section for mild to moderate open-angle glaucoma definition.

Use of ab interno stents for all other conditions **is considered experimental, investigational and/or unproven**.

**NOTE 2:** Shunts and stents are only able to reduce intraocular pressure (IOP) to the mid-teens and may be inadequate when very low IOP is needed to reduce glaucoma damage.

## Policy Guidelines

None.

## Description

Glaucoma surgery is intended to reduce intraocular pressure when the target IOP cannot be reached using medications. Due to complications with established surgical approaches (e.g., trabeculectomy), a variety of shunts are being evaluated as alternative surgical treatments for patients with inadequately controlled glaucoma. Microstents are also being evaluated in patients with mild-to-moderate open-angle glaucoma (OAG) currently treated with ocular hypotensive medication.

### Glaucoma

Glaucoma is the leading cause of irreversible blindness worldwide and is characterized by elevated intraocular pressure (IOP). In 2020, glaucoma affected approximately 52.7 million individuals globally, with a projected increase to 79.8 million in 2040. (1) Glaucoma has been reported to be 7 times more likely to cause blindness and 15 times more likely to cause visual impairment in Black individuals as compared to White individuals. In the U.S. in 2010, Black individuals had the highest prevalence rate of primary open angle glaucoma at 3.4% compared to 1.7% among White individuals.

In the primary (conventional) outflow pathway from the eye, aqueous humor passes through the trabecular meshwork, enters a space lined with endothelial cells (Schlemm canal), drains into collector channels, and then into the aqueous veins. Increases in resistance in the trabecular meshwork and/or the inner wall of the Schlemm canal can disrupt the balance of aqueous humor inflow and outflow, resulting in an increase in IOP and glaucoma risk.

The 2020 American Academy of Ophthalmology (AAO) defines the severity of glaucoma according to the following categories (56):

- Mild: definite optic disc or retinal nerve fiber layer (RNFL), or macular imaging abnormalities consistent with glaucoma and a normal visual field as tested with standard automated perimetry (SAP);
- Moderate: definite optic disc or RNFL, or macular imaging abnormalities consistent with glaucoma and visual field abnormalities in one hemifield that are not within 5 degrees of fixation as tested with SAP;
- Severe: definite optic disc or RNFL, or macular imaging abnormalities consistent with glaucoma and visual field abnormalities in both hemifields and/or loss within 5 degrees of fixation in at least one hemifield as tested with SAP;
- Indeterminate: definite optic disc or RNFL, or macular imaging abnormalities consistent with glaucoma inability of patient to perform visual field testing, unreliable/uninterpretable visual field test results, or visual fields not performed.

## Treatment

### *Ocular Medication*

First-line treatment typically involves pharmacologic therapy. Topical medications either increase aqueous outflow (prostaglandins, alpha-adrenergic agonists, cholinergic agonists, Rho kinase inhibitors) or decrease aqueous production (alpha-adrenergic agonists, beta blockers, carbonic anhydrase inhibitors). Pharmacologic therapy may involve multiple medications, have potential side effects, and may be inconvenient for older adults or incapacitated patients.

### *Surgery*

Surgical intervention may be indicated in patients with glaucoma when the target IOP cannot be reached pharmacologically. Surgical procedures for glaucoma aim to reduce IOP from impaired aqueous humor drainage in the trabecular meshwork and/or Schlemm canal.

Trabeculectomy (guarded filtration surgery) is the most established surgical procedure for glaucoma, which involves dissecting the conjunctiva, creating a scleral flap and scleral ostomy then suturing down the flap and closing the conjunctiva, allowing aqueous humor to directly enter the subconjunctival space. This procedure creates a subconjunctival reservoir, which can effectively reduce IOP, but commonly results in filtering “blebs” on the eye, and is associated with numerous complications (e.g., hemorrhage, scarring, hypotony, infection, leaks, bleb-related endophthalmitis) and long-term failure. Other surgical procedures (not addressed herein) include trabecular laser ablation, deep sclerectomy (which removes the outer wall of the Schlemm canal and excises deep sclera and peripheral cornea), and viscodanalostomy (which unroofs and dilates the Schlemm canal without penetrating the trabecular meshwork or anterior chamber). Canaloplasty involves dilation and tension of the Schlemm canal with a suture loop between the inner wall of the canal and the trabecular meshwork. This ab externo procedure uses the iTrack illuminated microcatheter (iScience Interventional) to access and dilate the entire length of the Schlemm canal and to pass the suture loop through the canal.

Insertion of shunts from outside the eye (ab externo) is another surgical option to lower IOP. Examples of ab externo devices cleared by the United States (U.S.) Food and Drug Administration (FDA) include the Ahmed, Baerveldt, Molteno, and EX-PRESS mini-shunt, which shunt aqueous humor between the anterior chamber and the suprachoroidal space. These devices differ by explant surface areas, shape, plate thickness, presence or absence of a valve, and details of surgical installation. Generally, the risk of hypotony (low pressure) is reduced with aqueous shunts compared with trabeculectomy, but IOP outcomes are worse than after standard guarded filtration surgery. The risk of postoperative infection is lower with shunts than with trabeculectomy, and failure rates are similar (10% of devices fail annually). The primary indication for aqueous shunts is for failed medical or surgical therapy, although some ophthalmologists have advocated their use as a primary surgical intervention, particularly for selected conditions such as congenital glaucoma, trauma, chemical burn, or pemphigoid.

#### *Minimally Invasive Glaucoma Surgeries*

Minimally invasive glaucoma surgeries (MIGS) are alternative, less invasive techniques that are being developed and evaluated. MIGS, which use microscopic-sized equipment and smaller incisions, involves less surgical manipulation of the sclera and the conjunctiva compared with other surgical techniques. There are several categories of MIGS: miniaturized trabeculectomy, trabecular bypass, milder laser photocoagulation, and totally internal or suprachoroidal stents. Shunts and stents can be administered through an external flap of the conjunctiva and sclera (ab externo) or in a small incision in the cornea with the devices inserted through the anterior chamber of the eye (ab interno). Some ab interno microstents may be inserted with injectors.

Examples of ab interno devices either approved or given marketing clearance by the FDA include the iStent, which is a 1-mm long stent inserted into the end of the Schlemm canal through the cornea and anterior chamber, iStent inject, iStent infinite, and XEN gelatin stent.

Because aqueous humor outflow is pressure-dependent, the pressure in the reservoir and venous system is critical for reaching the target IOP. Therefore, some devices may be unable to reduce IOP below the pressure of the distal outflow system used (e.g., <15 mm Hg) and are not indicated for patients for whom very low IOP is desired (e.g., those with advanced glaucoma). It has been proposed that stents such as the iStent, iStent inject, and Hydrus Microstent may be useful in patients with early-stage glaucoma to reduce the burden of medications and problems with compliance. One area of investigation is patients with glaucoma who require cataract surgery. An advantage of ab interno stents is that they may be inserted into the same incision and at the same time as cataract surgery. Also, most devices do not preclude subsequent trabeculectomy if needed. It may also be possible to insert more than 1 stent to achieve desired IOP.

#### **Regulatory Status**

The regulatory status of the various ab externo and ab interno aqueous shunts and microstents is summarized in Table 1.

The first-generation Ahmed™ (New World Medical), Baerveldt® (Advanced Medical Optics), Krupin (Eagle Vision), and Molteno® (Molteno Ophthalmic) ab externo aqueous shunts were cleared for marketing by the FDA through the 510(k) process between 1989 and 1993; modified Ahmed and Molteno devices were cleared in 2006. They are indicated for use “in patients with intractable glaucoma to reduce intraocular pressure where medical and conventional surgical treatments have failed.” The AquaFlow™ Collagen Glaucoma Drainage Device (STAAR Surgical) was approved by the FDA through the premarket approval (PMA) process for the maintenance of the subscleral space following nonpenetrating deep sclerectomy. In 2003, the ab externo EX-PRESS® Mini Glaucoma Shunt was cleared for marketing by the FDA through the 510(k) process.

In 2016, the XEN® Glaucoma Treatment System (Allergan), which consists of the XEN45 Gel Stent preloaded into the XEN Injector, was cleared for marketing by the FDA through the 510(k) process as an ab interno aqueous stent for management of refractory glaucoma. The approval was for patients with refractory glaucoma who failed previous surgical treatment or for patients with primary open-angle glaucoma (POAG) unresponsive to maximum tolerated medical therapy. The FDA determined that this device was substantially equivalent to existing devices, specifically the Ahmed™ Glaucoma Valve and the EX-PRESS® Glaucoma Filtration Device.

In 2018, the first microstent, the iStent® Trabecular Micro-Bypass Stent preloaded into the iStent *inject* device (Glaukos) was approved by the FDA through the 515(d) process for use in conjunction with cataract surgery for the reduction of IOP in adults with mild-to-moderate OAG currently treated with ocular hypotensive medication. In 2022, iStent infinite® was FDA-approved for primary OAG when medical and surgical treatment have failed. Notably, this device is not required to be performed in conjunction with cataract surgery and contains 3 stents preloaded into an injector system.

In August 2018, Alcon announced an immediate voluntary recall of the CyPass microstent, which had been approved by the FDA in 2016 for use in conjunction with cataract surgery in adults with mild-to-moderate OAG. The recall was based on 5-year postsurgery data from the COMPASS-XT long-term safety study. Results showed a statistically significant increase in endothelial cell loss among patients receiving the CyPass microstent compared with patients receiving cataract surgery alone.

In September 2023, a randomized controlled trial (NCT01881425) reported two-year follow-up outcomes comparing the PRESERFLO MicroShunt (Santen) to trabeculectomy in patients with mild to severe primary OAG inadequately controlled by maximum tolerated medical therapy. (2) As of October 2024, FDA approval of the device is still pending.

**Table 1. Regulatory Status of Aqueous Shunts and Stents**

Device	Manufacturer	Type	FDA Status	Date
AquaFlow™	STAAR Surgical	Drainage device	PMA	2001
Ahmed™	New World Medical	Aqueous glaucoma shunt, ab externo	510(k)	<1993

Baerveldt®	Advanced Medical Optics	Aqueous glaucoma shunt, ab externo	510(k)	<1993
Krupin	Eagle Vision	Aqueous glaucoma shunt, ab externo	510(k)	<1993
Molteno®	Molteno Ophthalmic	Aqueous glaucoma shunt, ab externo	510(k)	<1993
EX-PRESS®	Alcon	Mini-glaucoma shunt, ab externo	510(k)	2003
XEN® Gel Stent; XEN injector	AqueSys/Allergan	Aqueous glaucoma shunt, ab interno	510(k)	2016
iStent®; iStent inject®	Glaukos	Microstent, ab interno	515(d) in conjunction with cataract surgery	2018
iStent supra®	Glaukos	Suprachoroidal stent	Not approved; in clinical trial	N/A
CyPass®	Alcon	Suprachoroidal stent, ab interno	Company voluntarily recalled	2018
Hydrus™	Ivantis	Microstent, ab interno	PMA approval	2018
Beacon Aqueous Microshunt	MicroOptx	Micro-Shunt, ab externo	Not approved; in clinical trial	N/A
PRESERFLO® MicroShunt (previously InFocus)	Santan	Micro-Shunt, ab externo	Not approved; in clinical trial	N/A
iStent Infinate®	Glaukos	Microstent, ab interno	510(k)	2022

FDA: Food and Drug Administration; N/A: non-applicable; PMA: premarket approval.

FDA product codes: OGO, KYF.

## Rationale

Medical policies assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

## **Aqueous Shunts and Stents for Glaucoma**

### Clinical Context and Therapy Purpose

The purpose of aqueous shunts and stents in individuals who have glaucoma is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

#### *Populations*

The relevant populations of interest are:

- Individuals with refractory open-angle glaucoma (OAG);
- Individuals with mild-to-moderate primary open-angle glaucoma (POAG) who are undergoing cataract surgery;
- Individuals with indications for glaucoma treatment other than cataract surgery or refractory OAG.

#### *Interventions*

The therapies being considered are:

- For individuals with refractory OAG:
  - Ab externo aqueous shunts;
  - Ab interno aqueous stents;
- For individuals with mild-to-moderate OAG undergoing cataract surgery: ab interno aqueous stents;
- For individuals with indications for glaucoma treatment other than cataract surgery or refractory OAG: ab externo aqueous shunts or ab interno aqueous stents.

#### *Comparators*

Comparators include medical therapies and trabeculectomy.

#### *Outcomes*

The general outcomes of interest are change in intraocular pressure (IOP) and medication use. Changes in IOP and medication use are measured for at least 12 months. Safety measures involve longer follow-up, for several years.

### Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

### Ab Externo Aqueous Shunts

A Cochrane review by Minckler et al. (2006) included 15 randomized or pseudo-RCTs (N=1153) evaluating the Ahmed, Baerveldt, Molteno, and Schocket shunts. (3) Trabeculectomy was found to lower mean IOP by 3.8 mm Hg more than the Ahmed shunt at 1 year. This systematic review did not compare complications, because reviewers considered them to be too variably reported to permit comparative tabulation. There was no evidence of the superiority of 1 shunt over another. An update by Tseng et al. (2017) identified 27 studies, 4 of these studies compared Ahmed or Baerveldt shunts to trabeculectomy and 2 compared different types of shunts. (4) There was some evidence that Baerveldt and Molteno implants may reduce eye pressure more than Ahmed, and Molteno may lower eye pressure better than the Shocket.

A technology assessment on commercially available aqueous shunts, including the Ahmed, Baerveldt, Krupin, and Molteno devices, from the American Academy of Ophthalmology (AAO) was published by Minckler et al. (2008). (5) It indicated that IOP would generally settle at higher levels ( $\geq 18$  mm Hg) with aqueous shunts than with standard trabeculectomy (14-16 mm Hg) or trabeculectomy with antifibrotic agents 5-fluorouracil or mitomycin C (8-10 mm Hg). In a single study, mean IOPs with the Baerveldt shunt and adjunct medications were equivalent to trabeculectomy with mitomycin C (13 mm Hg). Five-year success rates for the 2 procedures were similar (50%). The assessment concluded that, based on level 1 evidence, aqueous shunts were comparable to trabeculectomy for IOP control and duration of benefit. The risk of postoperative infection was lower with aqueous shunts than with trabeculectomy.

Complications of aqueous shunts included: immediate hypotony after surgery, excessive capsule fibrosis and clinical failure, erosion of the tube or plate edge, strabismus, and, very rarely, infection. The most problematic long-term consequence of anterior chamber tube placement was accelerated damage to the corneal endothelium.

Zhang et al. (2022) compared the effectiveness of trabeculectomy and Ahmed and EX-PRESS implants in the treatment of primary and secondary glaucoma via a systematic review and network meta-analysis. (6) The review included 14 RCTs, involving 866 eyes of 808 patients. Overall, there were 339 eyes in the trabeculectomy group, 368 eyes in the EX-PRESS group, and

159 eyes in the Ahmed group. Results revealed that after 3 months, trabeculectomy was associated with similar improvement in IOP as compared to Ahmed (weighted mean difference [WMD], 0.014; 95% confidence interval [CI], -0.14 to 0.18) and EX-PRESS (WMD, 0.014; 95% CI, -0.072 to 0.097). However, at 1-year, EX-PRESS was associated with a significant improvement in IOP (WMD, 0.097; 95% CI, 0.008 to 0.18) as well as complete success (relative risk [RR], 0.73; 95% CI, 0.57 to 0.93) as compared to trabeculectomy. In a comparison of EX-PRESS and Ahmed implants, EX-PRESS was found to be superior to Ahmed with regard to reduction in the number of post-operative medications. Limitations of this meta-analysis included the presence of publication bias and heterogeneity of the included data.

#### *Baerveldt Glaucoma Shunt*

##### *Randomized Controlled Trials*

Results from the open-label, multicenter, randomized Tube versus Trabeculectomy study were reviewed in the 2008 AAO technology assessment and by Gedde et al. (2012) who reported on the 5-year follow-up. (5, 7, 8) That study included 212 eyes of 212 patients (age range, 18-85 years) from 17 study centers, who had trabeculectomy and/or cataract extraction with intraocular lens implantation and uncontrolled glaucoma with IOP of 18 mm Hg or greater and 40 mm Hg or lower on maximally tolerated medical therapy, randomized to tube (Baerveldt shunt) or trabeculectomy. Excluding patients who had died, the study had an 82% follow-up rate at 5 years, with a similar proportion of patients in the tube and trabeculectomy groups. At 5 years, neither IOP (14.3 mm Hg in the shunt group vs 13.6 mm Hg in the trabeculectomy group) nor the number of glaucoma medications (1.4 in the shunt group vs 1.2 in the trabeculectomy group) differed significantly based on intention-to-treat analysis. The cumulative probability of failure over the 5 years was lower in the shunt group (29.8%) than in the trabeculectomy group (46.9%), and the rates of reoperation were lower (9% vs 29%, respectively). The rates of loss of 2 or more lines of visual acuity were similar (46% in the shunt group vs 43% in the trabeculectomy group).

Subsequent publications have reported no significant differences between the groups for vision-related quality of life or visual field outcomes from the Tube vs Trabeculectomy study. (9-11)

#### *EX-PRESS Mini Shunt*

##### *Systematic Reviews*

A Cochrane review by Wang et al. (2015) evaluated the efficacy of adjunctive procedures for trabeculectomy. (12) Three RCTs were included which compared trabeculectomy alone with trabeculectomy plus EX-PRESS Mini Shunt. These trials were rated as having a high or unclear risk of bias using the Cochrane criteria. None of the RCTs reported a significant improvement for the EX-PRESS group. However, in the pooled analysis, IOP was lower in the combination group than in the trabeculectomy alone group (weighted mean difference, -1.58; 95% confidence interval [CI], -2.74 to -0.42). The pooled analysis also showed that subsequent cataract surgery was less frequent in the combination group than in trabeculectomy alone (relative risk, 0.34; 95% CI, 0.14 to 0.74). The combination group had a lower rate of some complications (e.g., hyphema, needling). An updated analysis by Park et al. (2023) identified a

total of 8 studies (7 with EX-PRESS and 1 with PreserFlo MicroShunt). (13) Low-certainty evidence showed that adjunct EX-PRESS resulted in lower IOP at 1 year (MD, -1.76; 95% CI, -.81 to -.70).

#### Randomized Controlled Trials

A U.S. multicenter randomized trial by Netland et al. (2014), compared trabeculectomy with EX-PRESS implantation in 120 patients (120 eyes) (see Table 2). (14) Comparator groups were similar at baseline. Throughout a 2-year postsurgical follow-up, average IOP and number of medications were similar between groups (see Table 3). Surgical success was 90% and 87% at 1 year and 83% and 79% at 3 years in the EX-PRESS and trabeculectomy groups, respectively. Visual acuity returned to near baseline levels at 1 month after EX-PRESS implantation (median, 0.7 months) and at 3 months after trabeculectomy (median, 2.2 months;  $p=0.041$ ). Postoperative complications were higher after trabeculectomy (41%) than after EX-PRESS implantation (18.6%).

Additional single center RCTs have corroborated the results of the multicenter trial. (15-20)

**Table 2. Summary of Key RCT Characteristics for EX-PRESS**

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
de Jong et al. (2009) (15); de Jong et al. (2011) (16)	Netherlands	1	2003-2004	Patients with primary OAG not controlled by IOP medication.	EX-RESS (n=39)	Trabeculectomy (n=39)
Netland et al. (2014) (14)	U.S., Canada	7	NR	Patients with OAG treated with IOP medications who were candidates for glaucoma surgery.	EX-RESS (n=59)	Trabeculectomy (n=61)
Wagschal et al. (2015) (17); Gonzalez-Rodriguez et al. (2016) (18)	Canada	1	2011-2012	Patients with primary OAG not controlled by IOP medication.	EX-RESS (n=33)	Trabeculectomy (n=31)
Konopinska et al. (2021) (19) (NCT04335825)	Poland	1	2016-2019	Patients with OAG not controlled by IOP medication who qualified for both cataract and OAG surgery.	Phaco EX-PRESS (n=43)	Phaco – Trabeculectomy (n=38)

IOP: intraocular pressure; NR: not reported; OAG: open-angle glaucoma; Phaco: phacoemulsification; RCT: randomized controlled trial; U.S.: United States.

**Table 3. Summary of Key RCT Results for Ex-PRESS**

Study	Mean IOP (SD), mm Hg		p	Mean Medication Use (SD)	
	EX-PRESS	Trabeculectomy		EX-PRESS	Trabeculectomy
<b>Netland et al. (2014) (14)</b>					
Baseline	25.1 (6.0)	26.4 (6.9)	0.27	3.1 (1.1)	3.1 (1.2)
Month 6	13.8 (4.7)	11.9 (4.6)	0.03	NR	NR
Year 2	14.7 (4.6)	14.6 (7.1)	0.93	0.9 (1.3)	0.7 (1.2)

IOP: intra-ocular pressure; NR: not reported; SD: standard deviation; RCT: randomized controlled trial.

### Comparative Effectiveness Analyses

Five-year results of 2 RCTs comparing the Ahmed and Baerveldt shunts have been published. The Ahmed Baerveldt Comparison (ABC) study was a multicenter international RCT evaluating the comparative safety and efficacy of the Ahmed Glaucoma Valve and Baerveldt Glaucoma Implant in 276 adults with previous incisional eye surgery or refractory glaucoma. (21, 22) The ABC was funded by National Eye Institute, Research to Prevent Blindness, and New World Medical. The Ahmed Versus Baerveldt (AVB) study, reported by Christakis et al. (2016), was an international, multicenter RCT enrolling 238 patients with uncontrolled glaucoma despite maximally tolerated medical therapy that was funded by the Glaucoma Research Society of Canada. (23)

Christakis et al. (2017) analyzed 5-year pooled data from the ABC and AVB trials comparing the relative efficacy of the 2 implants. (24) At year 5, mean IOP was 15.8 mm Hg in the Ahmed group and 13.2 mm Hg in the Baerveldt group ( $p=.007$ ). The cumulative failure rate in the Ahmed group was 49%; in the Baerveldt group, it was 37%. Mean glaucoma medication use was significantly lower in patients receiving the Baerveldt implant than in patients receiving the Ahmed implant ( $p=0.007$ ). Visual acuity was similar between both groups. While efficacy measures were significantly better in the Baerveldt group, these patients experienced more hypotony (4.5%) than patients in the Ahmed group (0.4%;  $p=.002$ ).

### Section Summary: Ab Externo Aqueous Shunts

Evidence for the use of ab externo aqueous shunts for the treatment of OAG uncontrolled by medications consists of RCTs comparing shunts with trabeculectomy. Outcomes of interest are IOP and antiglaucoma medication use. Follow-up among the trials ranged from 1 to 5 years. Results from ab externo aqueous shunts are similar to trabeculectomy, while adverse event rates were higher among patients undergoing trabeculectomy.

The comparative effectiveness of 2 ab externo devices (the Ahmed and Baerveldt shunts) has been evaluated in 2 trials, the AVB, and the ABC trials. These trials reported similar results, with both devices lowering IOP significantly. Compared with patients receiving the Ahmed shunt, patients receiving the Baerveldt shunt experienced lower IOP and needed fewer medications. However, patients receiving the Baerveldt shunt experienced higher rates of hypotony-related complications.

## Ab Interno Aqueous Stents

This section reviews the evidence for ab interno stents with the FDA approval or marketing clearance.

### *Xen Glaucoma Treatment System*

#### Systematic Reviews

Lim et al. (2022) conducted a systematic review and meta-analysis of 14 studies (N=963 eyes) involving the stand alone XEN45 gel stent ab interno device implant. (25) The review included 7 prospective and 7 retrospective studies. The mean age of included patients was 66 years and the maximum follow-up duration ranged from 6 to 30 months. A variety of surgical techniques were employed across the studies; however, surgical steps were largely consistent. Results revealed that implantation of the XEN45 gel stent significantly decreased IOP ( $p<.001$ ) across all timepoints (1 day, 1 week, 1, 3, 6, 12, 18, and 24 months) with a mean decrease of 7.44 mm Hg at 24 months. The use of IOP-lowering medications was also reduced significantly ( $p<.001$ ) post-implantation across all timepoints (1 week, 1, 3, 6, 12, 18, and 24 months) with a mean reduction of 1.67 medications at 24 months. Serious adverse events occurred rarely with transient numerical hypotony the most common postoperative complication. Postoperative needling procedures were required in 38% of eyes during the entire follow-up period. The overall quality of the evidence within the systematic review was low, with most included studies being case series with relatively short follow-up durations and a lack of standardized definitions of treatment success and failure. Additional RCTs with a clinically meaningful definition of success and failure are needed.

Another systematic review and meta-analysis that evaluated the efficacy of the XEN gel stent implant in 78 eligible studies reported similar conclusions. (26) Following XEN stent implantation, there was a significant reduction in IOP ( $p<.001$ ), and the number of anti-glaucoma medications used ( $p<.001$ ) through 48 months post-surgery. However, the quality of included studies was noted to be relatively low, and the definition of outcomes was inconsistent across the included studies.

#### Randomized Controlled Trial

Sheybani et al. (2023) conducted a randomized, noninferiority trial comparing XEN45 gel stent to trabeculectomy in patients (N=139) with an IOP of 15 to 44 mm Hg while receiving topical IOP medication. (27) At 12 months XEN45 was noninferior to trabeculectomy in terms of surgical success which was defined as at least a 20% reduction in IOP without a medication increase, clinical hypotony, vision loss, or secondary surgical intervention (between group difference, -6.1%; 95% CI, -22.9% to 10.8%). XEN45 resulted in fewer postoperative interventions and faster visual recovery than trabeculectomy.

#### Nonrandomized Comparative Studies

Schlenker et al. (2017) published a multicenter, retrospective comparative study that compared the risk, safety, and efficacy for stand-alone ab interno microstent implantation with mitomycin C (MMC) and trabeculectomy plus MMC (Table 4). (28) Implantations of the ab interno XEN 45 gelatin microstent is a less invasive surgery than trabeculectomy. The primary outcome was the

hazard ratio (HR) of failure, defined as 2 consecutive IOP readings of less than 6 mm Hg, including vision loss. Success was measured by the withdrawal of glaucoma-related medications at 1-month postsurgery. The adjusted HR of failure of the microstent relative to trabeculectomy was 1.2 for complete success (95% CI, 0.7 to 2.0). Both surgeries had a 75% survival of approximately 10 months for complete success. During the last reported follow-up (varying times), antiglaucoma medications were being used by 25% of patients who received the microstent implantation and 33% of trabeculectomy patients. Patients in both groups reported similar numbers of postoperative interventions, such as laser suture lysis and needling. The need for reoperation was higher among those who had undergone microstent implantation, but this difference was not statistically significant. The authors concluded that the ab interno gelatin microstent with MMC was noninferior to trabeculectomy plus MMC. Changes in IOP and medication use appear in Table 5.

Wagner et al. (2020) also reported similar success rates for trabeculectomy (65.5%, 95% CI, 55.6 to 75.9%) and XEN Implant (58.5%, 95% CI, 47.6 to 69.4%,  $p = .16$ ; adjusted odds ratio 0.66, 95% CI, 0.32 to 1.37) but a greater reduction in IOP with trabeculectomy (10.5 mm Hg) compared to the XEN implant (7.2 mm Hg;  $p = .003$ ). (29) Baseline measurements showed older age (73.0 vs 67.2) and a lower number of medication classes (2.0 vs 3.0) for the XEN group. A regression mixed model that adjusted for gender, age, preoperative IOP, and medications did not indicate a difference in the proportion of success for the 2 groups.

Stoner et al. (2021) conducted a retrospective comparative study of 100 eyes that had undergone either XEN or EX-PRESS standalone shunt implantation at a single center. (30) Surgical success was defined as IOP between 6- and 18-mm Hg without reoperation, loss of light perception, device removal, or use of glaucoma medications. The incidence of adverse effects during the first 3 months was lower with the XEN implant, but the failure rate at 1 year was higher (HR 3.94, 95% CI, 1.73 to 9.00,  $p = .001$ ) compared to EX-PRESS. Sensitivity analysis to adjust for differences in baseline characteristics between the groups in this retrospective study achieved similar results.

#### Non-Comparative Observational Studies

The largest study with a follow-up of longer than 1 year was by Gabbay et al. (2021), who reported a retrospective analysis of 205 patients/eyes that had received an XEN implant. (31) At 3 years, 25% of eyes met the criteria for success, with a failure rate of 25% and requirement for needling in 36.6%. For eyes that retained an XEN implant, IOP decreased from an average of 22.6 mmHg (standard deviation [SD], 7.0) before surgery to 14.0 (SD, 2.9) at 3 years; the number of medications decreased from an average of 2.6 (SD, 1.1) to 0.6 (SD, 1.0) at 3 years. The failure rate was higher in non-Caucasians (74% of 13) compared to Caucasians (21% of 188,  $p < .001$ ), with Caucasians comprising 93.5% of the study population.

**Table 4. Summary Characteristics for Non-Randomized Comparative Studies Using the XEN Implant for Refractory Open-Angle Glaucoma**

Study	Country	Participants	Treatment Delivery	FU
-------	---------	--------------	--------------------	----

Schlenker et al. (2017) (28)	Austria, Belgium, Canada, Germany	Patients with OAG, pseudoexfoliation, pigment dispersion, normal-tension, angle-recession, combined mechanism, history of angle closure, or juvenile glaucoma and no prior incisional surgery	<ul style="list-style-type: none"> <li>• XEN alone (n=185)</li> <li>• Trabeculectomy (n=169)</li> </ul>	Up to 30 months (last visit in chart)
Wagner et al. (2020) (29)	Germany	Consecutive patients with refractory OAG, pseudoexfoliation, pigment dispersion, or normal-tension glaucoma who underwent surgery from January 2016 to February 2018	<ul style="list-style-type: none"> <li>• XEN alone (n=82 eyes)</li> <li>• Trabeculectomy (n=89 eyes)</li> </ul>	1 year
Stoner et al. (2021) (30)	U.S.	Patients with uncontrolled glaucoma with either IOP uncontrolled by medications or progression of glaucoma	<ul style="list-style-type: none"> <li>• XEN (n=52)</li> <li>• EX-PRESS (n=48)</li> </ul>	1 year

FU: follow-up; IOP: intraocular pressure; n: number; OAG: open-angle glaucoma; U.S.: United States.

**Table 5. Summary of Results for the XEN Implant for Refractory Open-Angle Glaucoma**

Study	Population	Median IOP (SD), mm Hg		Medication Use (SD)	
		Baseline	1 Year <sup>a</sup>	Baseline	1 Year <sup>a</sup>
Schlenker et al. (2017) (28)	XEN alone	24.0 (IQR: 19 to 32)	13.0 (IQR: 10 to 15)	3.0 (IQR: 3 to 4)	0.0 (IQR: 0 to 1)
	Trabeculectomy	24.0 (IQR: 19 to 30)	13.0 (IQR: 10 to 16)	3.0 (IQR: 3 to 4)	0.0 (IQR: 0 to 1)
Wagner et al. (2020) (29)	XEN	19.0 (IQR 16.8-25.0)	7.2 (8.2) reduction	2.0 (1.0-3.0)	0.3 (0.5)
	Trabeculectomy	21.0 (IQR 17.0-27.0)	10.5 (9.2) reduction	3.0 (2.0-4.0)	0.2 (0.5)
Stoner et al. (2021) (30)	XEN	21.4 (1.2)	13.0 (0.6)	2.8 (0.2)	1.5 (0.2)
	EX-PRESS	18.9 (1.1)	11.5 (0.8)	3.2 (0.2)	0.5 (0.2)

<sup>a</sup> Follow-up for Schlenker (2017) was not 1 year, but last visit in retrospective chart review.

IOP: intraocular pressure; IQR: interquartile range, NR: not reported; SD: standard deviation.

### Section Summary: Ab Interno Aqueous Stents

Clearance for the XEN gel stent as a stand-alone procedure was based on a review in which the FDA concluded that while there were technical differences between the stent and predicate

devices (shunts), the differences did not affect safety and effectiveness in lowering IOP and medication use. Evidence for the use of the XEN implant consists of systematic reviews, an RCT, and nonrandomized comparative studies which retrospectively reviewed charts of patients either receiving the XEN implant or undergoing a trabeculectomy or implantation of an EX-PRESS shunt. Additional evidence consists of several single-arm studies.

The RCT found XEN45 to be noninferior to trabeculectomy. The nonrandomized comparative studies included patients with different types of glaucoma and found that patients receiving the XEN implant experienced reductions in IOP, and medication use similar to patients undergoing trabeculectomy. A retrospective study compared the XEN implant with the EX-PRESS implant and found fewer adverse events in the first 3 months, but lower efficacy and higher failure rates at 1 year. Although there was little information on how patients were chosen to receive the different treatments in these comparative trials, statistical methods were used to address baseline differences between the groups. The single-arm studies, with up to 3 years of follow-up, consistently show that patients receiving the XEN implant experience reductions in IOP and medication use. RCTs with larger sample sizes and longer follow-up are needed to compare the outcomes of the different surgical treatments.

#### Aqueous Microstents in Conjunction with Cataract Surgery

The iStent and iStent inject, which is preloaded with 2 stents, have FDA approval for use in conjunction with cataract surgery. An additional stent, the CyPass, had FDA approval but was voluntarily recalled by the manufacturer in 2018, as follow-up data has shown significant endothelial cell loss among patients receiving the CyPass in conjunction with cataract surgery compared with patients receiving cataract surgery alone. Studies comparing implantation of stents during cataract surgery with cataract surgery alone are discussed below.

#### *iStent*

##### Systematic Reviews

A 2019 Cochrane review on the iStent in patients with OAG was published by Le et al. (Table 6). (32) The authors identified 7 RCTs, all of which were considered to be at high or unclear risk of bias. Four of the trials compared iStent in combination with cataract surgery to cataract surgery alone, 2 RCTs compared treatment with iStent or iStent inject to medical therapy, and 1 RCT compared 1, 2, or 3 iStents. Results of the meta-analyses on use of the iStent in combination with cataract surgery are shown in Table 7. Implantation of 1 or 2 iStents resulted in a higher proportion of patients who were drop free (relative risk: 1.38) and reduced the mean number of drops when compared to phacoemulsification alone (-0.42 drops). The review concluded that based on the 4 trials, there was very low-quality evidence that iStent may result in a higher proportion of patients who are drop free or achieve better IOP control.

An industry-sponsored meta-analysis of standalone iStents was reported by Healy et al. (2021). (33) The investigators included 4 RCTs and 9 non-randomized or single-arm studies with at least 6 months of follow-up. The number of eyes in the studies ranged from 15 to 99 (total N=778). The pooled weighted reduction in IOP was reported as 31.1% at 6 to 12 months and 32.9% at 60 months with a reduction of approximately 1 medication in the pooled analysis. In the

individual studies, the reduction in IOP ranged from -1.0 to -10.7; the largest reduction in IOP was in a prospective case series (n=44) with 25% loss to follow-up. The lowest reduction in IOP (-1.0) was in a larger RCT (n=77) with low loss to follow-up (2.5%). Notably, the systematic review did not report the number of device failures in these studies. Additional limitations are the inclusion of retrospective case series and the high heterogeneity between studies, which would typically preclude meta-analysis.

**Table 6. Meta-analysis Characteristics**

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Le et al. (2019) (32)	Aug 2018	7	Eyes with open-angle glaucoma	765 (33 to 239)	RCT	42 months

N: number; RCT: randomized controlled trial.

**Table 7. Meta-analysis Results**

Study	Drop Free Compared to Phacoemulsification Alone	Change in Drops Compared to Phacoemulsification Alone	Change in IOP Compared to Phacoemulsification Alone
<b>Le et al. (2019) (32)</b>			
N	239 (2 RCTs)	282 (2 RCTs)	284 (3 RCTs)
Pooled effect (95% CI)	RR: 1.38 (1.18 to 1.63)	-0.42 (-0.60 to -0.23)	-1.24 mmHg
$I^2$ (p)	67% (p)	0%	

CI: confidence interval; IOP: intraocular pressure; RCT: randomized controlled trial; RR: relative risk

#### iStent and iStent inject Pivotal Trials

Included in the Cochrane review were results from the iStent U.S. investigational device exemption, open-label, 29-site, multicenter RCT. Results were reported to the FDA in 2010, with 1-year results published by Samuelson et al. (2011) and 2-year results published by Craven et al. (2012) (Table 8). (34, 35) Trial objectives were to evaluate the incremental effect on IOP of iStent implantation compared to cataract surgery alone and to determine the potential benefit of combining 2 therapeutic treatments into a single surgical event. A total of 240 patients (mean age, 73 years) with cataracts and mild-to-moderate OAG (IOP  $\leq$ 24 mm Hg controlled on 1-3 medications) underwent a medication washout period. Patients were randomized to cataract surgery plus iStent implantation or cataract surgery only. Follow-up visits were performed at 1, 3, 6, and 12 months. Results were assessed by intention-to-treat analysis with the last observation carried forward and per-protocol analysis. The proportion of eyes meeting both the primary (unmedicated IOP  $\leq$ 21 mm Hg) and secondary outcomes (IOP reduction  $\geq$ 20% without medication) was higher in the treatment group than in the control group through 1-year follow-up (72% of treatment eyes vs 50% of control eyes achieved the primary efficacy endpoint,  $p<0.001$ ). The proportion of patients achieving the secondary efficacy endpoint was 66% in the treatment group and 48% in the control group ( $p=0.003$ ). Ocular hypotensive medications were initiated later in the postoperative period and used in a lower proportion of

patients in the treatment group throughout 1-year follow-up (e.g., 15% vs 35% at 12 months). Mean reduction in IOP was similar in both groups, though the control group used slightly more medication (mean, 0.4 medications) than the treatment group (0.2 medications) at 1 year (Table 9). At 2-year follow-up, 199 (83%) patients remained in the study. The primary endpoint (unmedicated IOP  $\leq$ 21 mm Hg) was reached by 61% of patients in the treatment group and 50% of controls ( $p=0.036$ ). (35) Secondary outcomes include IOP reduction of 20% or more without medication (53% vs 44%) and the mean number of medications used (0.3 vs 0.5), no longer differed significantly between groups at 2 years. As noted by the FDA, this study was conducted in a restricted population with an unmedicated IOP of 22 mm Hg or higher and a medicated IOP of 36 mm Hg or lower.

The pivotal trial on the iStent inject was reported by Samuelson et al. (2019). (36) A total of 505 patients undergoing cataract surgery were randomized after lens implantation to insertion of 2 smaller iStents or control. Results were assessed by intention-to-treat analysis and per-protocol analysis, with patients requiring additional surgical procedures considered to be failures. The addition of medications was based on a standardized protocol. At the 2-year follow-up, a greater percentage of patients had achieved at least a 20% reduction in IOP (75.8% vs 61.9%,  $p=0.005$ ), had a greater reduction in IOP (7.0 vs 5.4,  $p<0.001$ ), and required fewer topical medications (0.4 vs 0.8,  $p<0.001$ ).

Limitations of these studies are described in Tables 10 and 11. The 2 main limitations are that there was no masking to treatment and durability of these microstents after 2 years was not reported. Continued patency of the stents and need for additional treatments has been evaluated through 4 years in studies from the Microinvasive Glaucoma Surgery (MIGS) study group and are described below.

**Table 8. Summary of Pivotal RCT Characteristics**

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Samuelson et al. (2011) (34); Craven et al. (2012) (35)	U.S.	29	2005-2007	Patients with mild-to-moderate POAG, unmedicated IOP $\geq$ 22 and $\leq$ 36 mm Hg	iStent plus cataract surgery (n=116)	Cataract surgery alone (n=123)
Samuelson et al. (2019) (36)	U.S.		2011	Patients with mild-to-moderate POAG, unmedicated IOP $\geq$ 21 and $\leq$ 36 mm Hg	iStent inject (2 stents) plus cataract surgery (n=387)	Cataract surgery alone (n=118)

IOP: intraocular pressure; POAG: primary open-angle glaucoma; RCT: randomized controlled trial; U.S. United States.

**Table 9. Summary of Pivotal RCT Results**

Study	≥ 20% Reduction in Unmedicated IOP at 24 mo n (%)	Mean Reduction in IOP at 24 mo mm Hg (SD)	Mean IOP (SD), mm Hg		p	Mean Medication Use (SD)		p
			iStent	Cataract Alone		iStent	Cataract Alone	
<b>Samuelson et al. (2011) (34); Craven et al. (2012) (35)</b>								
Baseline			18.6 (3.4)	17.9 (3.0)	NR	1.6 (0.8)	1.5 (0.6)	
Year 1			17.0 (2.8)	17.0 (3.1)	NR	0.2 (0.6)	0.4 (0.7)	0.016
Year 2			17.1 (2.9)	17.8 (3.3)	NR	0.3 (0.6)	0.5 (0.7)	
<b>Samuelson et al. (2019) (36)</b> <b>iStent inject</b>	288/380 (75.8%)	7.0 (4.0)	17.1 (3.6)			0.4 (0.8)		
Cataract Alone	73/118 (61.9%)	5.4 (3.7)	17.8 (3.5)			0.8 (1.0)		
p-Value	0.005	<0.001				<0.001		

IOP: intraocular pressure; NR: not reported; RCT: randomized controlled trial; SD: standard deviation.

**Table 10. Study Relevance Limitations**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Follow-Up <sup>e</sup>
Samuelson et al. (2011) (34)					Patency after 2 years is unknown
Samuelson et al. (2019) (36)					Patency after 2 years is unknown

The study limitations stated in this table are those notable in the current literature review; this is not a comprehensive gaps assessment.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

<sup>b</sup> Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

<sup>c</sup> Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

<sup>d</sup> Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported.

<sup>e</sup> Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

**Table 11. Study Design and Conduct Limitations**

Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness <sup>d</sup>	Power <sup>e</sup>	Statistical <sup>f</sup>
Samuelson et al. (2011) (34)		2, 3. No blinding of assessors				
Samuelson et al. (2019) (36)		2, 3. No blinding of assessors				

The study limitations stated in this table are those notable in the current literature review; this is not a comprehensive gaps assessment.

<sup>a</sup> Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup> Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

<sup>c</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup> Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

<sup>e</sup> Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

<sup>f</sup> Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Hooshmand et al. (2019) reported a non-randomized comparative study on outcomes with the use of the iStent inject, which simultaneously injects 2 stents through a single ab interno opening, compared to the first generation single iStent. (37) The iStent inject was developed to provide easier ab interno insertion and comes preloaded with 2 stents that are smaller than the first-generation iStent. There was no significant difference between the earlier model and the second-generation device on outcomes at 12 months, but Kaplan-Meier analysis found an earlier time to add topical medications in the iStent inject patients. Limitations of the study include the length of follow-up, which was limited by the time that the iStent inject had been available, and the non-randomized design. In addition, the study compared 2 cohorts from

different time periods, those who had been treated with the first-generation device and those who had been treated with the second-generation device.

Al Yousef et al. (2020) conducted a matched comparison of the iStent inject and ab interno trabeculectomy in 78 eyes. (38) IOP was reduced in both groups at 1-month follow-up but began to rise at 12 months in the iStent inject group. By 24 months, the IOP in the iStent inject group had returned to near preoperative levels. The IOP in the Trabectome study group was lower than the iStent inject group throughout follow-up.

Efficacy of the iStent inject at 3-year follow-up was reported by Salimi et al. (2021) in a consecutive case series of 124 eyes with different glaucoma subtypes and severities. (39) Mean IOP in patients who retained an implant was reduced from 16.9 mm Hg preoperatively to 13.17 mm Hg ( $p<.001$ ) with a reduction in medications from 2.38 to 1.16 ( $p <.001$ ). The 3-year survival rate of the implant was only 74%.

Matsuo et al. (2023) reported on a retrospective cohort study comparing the effectiveness of combined cataract surgery with microhook ab-interno trabeculotomy versus iStent trabecular micro-bypass stent in 75 eyes with primary OAG and preoperative IOP below 15 mmHg. (40) At 1-year follow-up, both groups showed significant reductions in antiglaucoma medications (from 3.4 to 2.5 for trabeculotomy and 2.5 to 2.0 for iStent;  $p<.05$ ). The trabeculotomy group demonstrated higher rates of surgical success compared to the iStent group ( $p<.01$ ).

Gaskin et al. (2024) reported on a prospective, randomized, assessor-masked controlled trial with 87 patients (n=101 eyes) that compared the effectiveness of cataract surgery plus iStent Inject implantation versus cataract surgery alone in patients with mild-to-moderate glaucoma. (41) At the 24-month follow-up, the iStent Inject group used significantly fewer medications compared to the cataract surgery alone group (mean, 0.7 vs. 1.5;  $p=.008$ ) and had a higher proportion of patients taking no glaucoma medications (57% vs. 36%). At 4 weeks post-surgery, a lower IOP was observed in the iStent group (mean difference, -2.8 mmHg [95% CI, -4.7 to -1]) but there was no difference in subsequent follow-up assessments. Both groups showed improvement in patient-reported outcomes from baseline levels (Ocular Surface Disease Index score and Glaucoma Activity Limitation Questionnaire), with no significant differences between groups. The safety profiles were similar between the two groups.

### *Hydrus Microstent* Systematic Reviews

A Cochrane review by Otarola et al. (2020) included 3 studies with 808 participants. (42) Two studies (described below) were conducted in patients with cataracts and OAG (n=653) and compared the Hydrus microstent combined with cataract surgery to cataract surgery alone. (43, 44) They found moderate-certainty evidence that adding the Hydrus microstent to cataract surgery in patients with mild or moderate OAG increased the proportion of participants who were medication-free at 12 month (risk ratio 1.59, 95% CI, 1.39 to 1.83) and 24 month follow-up (risk ratio 1.63, 95% CI, 1.40 to 1.888), and reduced unmedicated IOP by 2 mm Hg, the number of medications by -0.41, and the need for secondary glaucoma surgery.

The third study compared the Hydrus microstent with the iStent in patients without cataract surgery. (45) This study is described in the next section on microstents as a stand-alone procedure.

#### Randomized Controlled Trials

Trials on the Hydrus Microstent are described in Tables 12 and 13.

Pfeiffer et al. (2015) reported on a single-masked, randomized trial with 100 patients (100 eyes) that compared the effectiveness of the Hydrus Microstent plus cataract surgery with cataract surgery alone. (43) At the 24-month follow-up, the proportion of patients with a 20% reduction in IOP was significantly higher with the Hydrus Microstent (80% vs 46%,  $p<0.001$ ) and the mean IOP after medication washout was lower (16.9 mm Hg vs 19.2 mm Hg,  $p=0.009$ ) compared with cataract surgery alone, respectively. The microstent group used significantly fewer medications (0.5 vs 1.0,  $p=0.019$ ) and had a higher proportion of patients taking no hypotensive medications at the time of cataract surgery (73% vs 38%,  $p=0.001$ ).

Samuelson et al. (2019) reported on a multicenter RCT (HORIZON) comparing implantation of a single Hydrus Microstent following cataract surgery versus cataract surgery alone (Table 13). (44) Patients were masked to treatment assignment for the course of the study. The primary endpoint was percent demonstrating a 20% reduction in unmedicated IOP. Significantly more patients receiving the microstent following cataract surgery experienced a 20% reduction in unmedicated IOP compared with patients undergoing cataract surgery alone (77% vs 58%;  $p<0.001$ ). A posthoc analysis of the HORIZON trial found that glaucoma patients receiving cataract surgery with Hydrus microstent implantation showed significantly slower visual field progression (-0.26 dB/year) compared to those receiving cataract surgery alone (-0.49 dB/year), with a reduced proportion of fast progressors over a 5-year follow-up period. (46)

Comparisons of mean washed out IOP and the mean number of medications used are presented in Table 13.

**Table 12. Summary of Key RCT Characteristics for the Hydrus Microstent**

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Pfeiffer (2015) (43)	Germany, Italy, Spain, the Netherlands	7	2011 to 2012	Patients with concurrent OAG and cataract	Cataract surgery plus Hydrus Microstent implantation (n=50)	Cataract surgery alone (n=50)
Samuelson (2019) (44)	Germany, Italy, Mexico,	26	2012 to 2015	Patients with age-related	Cataract surgery plus Hydrus	Cataract surgery

	Philippines, Poland, Spain, U.K., U.S.			cataract and mild to moderate POAG:	Microstent implantation (n=369)	alone (n=187)
--	---	--	--	--	---------------------------------------	------------------

OAG: open-angle glaucoma; POAG: primary open-angle glaucoma; RCT: randomized controlled trial; U.K.: United Kingdom; U.S.: United States.

**Table 13. Summary of Key RCT Results for the Hydrus Microstent**

Study	Mean washed out IOP			Mean medication use		
	Hydrus Microstent	Cataract alone	p	Hydrus Microstent	Cataract alone	p
<b>Pfeiffer (2015) (43)</b>						
Baseline	26.3 +/- 4.4	26.6 +/- 4.2	0.7	2.0 +/- 1.0	2.0 +/- 1.1	0.8
Year 2	16.9 +/- 3.3	19.2 +/- 4.7	0.009	0.5 +/- 1.0	1.0 +/- 1.0	0.02
<b>Samuelson (2019) (44)</b>						
Baseline mean	25.5 +/- 3.0	25.4 +/- 2.9	NS	1.7 +/- 0.9	1.7 +/- 0.9	NS
Year 2	17.4 +/- 3.7	19.2 +/- 3.8	NR	0.3 +/- 0.8	0.7 +/- 0.9	<0.001

IOP: intraocular pressure; NR: not reported; NS: not significant; RCT: randomized controlled trial.

#### Observational Studies

Fea et al. (2017) conducted a retrospective review of 92 patients undergoing cataract surgery plus Hydrus Microstent implantation. (47) Two-year follow-up showed improvements in IOP and medication use. Mean IOP at baseline was 19.4 mm Hg, decreasing significantly by 6 months to 15.6 mm Hg, which was maintained at 2 years of follow-up (15.7 mm Hg). The mean number of medications was 2.1 at baseline, decreasing significantly by 6 months to 0.5, which was maintained through 2 years of follow-up (0.7).

Salimi et al. (2023) conducted a single-surgeon, consecutive case series of 106 glaucomatous eyes undergoing Hydrus Microstent implantation plus cataract surgery. (48) At 3-year follow-up, surgical success rates ranged from 67% to 91% depending on criteria. Mean IOP decreased significantly from 18.9 mmHg at baseline to 13.9 mmHg (-26.5%; p<.001), while the mean number of antiglaucoma medications decreased from 3.0 to 2.0 (-33%; p<.001). Visual acuity improved and was maintained, while structural and functional markers of glaucoma remained stable.

#### CyPass

The FDA evaluated the clinical performance of the CyPass Micro-Stent system based on the pivotal Clinical Study to Assess the Safety and Effectiveness of the Transcend CyPass Glaucoma Implant in Patients With Open-Angle Glaucoma Undergoing Cataract Surgery (COMPASS) trial (NCT01085357). COMPASS was a multicenter RCT comparing the safety and efficacy of CyPass Micro-Stent plus cataract surgery with cataract surgery alone for treating mild-to-moderate primary OAG in patients undergoing cataract surgery. Evidence from the RCT supported the use of the CyPass stent in conjunction with cataract surgery; however, in August 2018, the

manufacturer voluntarily withdrew the device from the market because a long-term study showed that patients receiving CyPass in conjunction with cataract surgery experienced statistically significant endothelial cell loss compared with patients who underwent cataract surgery alone.

#### Section Summary: Ab Interno Aqueous Microstents

Implantation of 1 or 2 microstents has received the FDA approval for use in conjunction with cataract surgery for reduction of IOP in adults with mild-to-moderate OAG currently treated with ocular hypotensive medication. RCTs and meta-analyses of RCTs have compared cataract surgery alone to microstent implantation in conjunction with cataract surgery when IOP is at least partially controlled with medication. When compared to cataract surgery alone, the studies showed modest but statistically significant decreases in IOP, and medication use through the first 2 years when stents were implanted in conjunction with cataract surgery. A decrease in topical medication application is considered to be an important outcome for patients and reduces the problem of non-compliance that can affect visual outcomes.

#### Microstent Implantation as a Stand-Alone Procedure

##### *iStent*

The iStent was approved by the FDA to be used in conjunction with cataract surgery to reduce IOP in patients with mild-to-moderate OAG. The studies described below evaluated the use of the iStent or iStent inject as a stand-alone procedure.

##### Systematic Reviews

The Cochrane review by Le et al. (2019) on the iStent in patients with OAG identified 2 RCTs that compared treatment with iStent or iStent inject to medical therapy and 1 RCT that compared 1, 2, or 3 iStents. (32) Results of the systematic review are shown in Table 14. Meta-analysis was not performed due to heterogeneity. However, in both trials, iStent implantation resulted in a higher proportion of patients who were drop free and reduced the mean number of drops when compared to medical therapy. One RCT indicated that compared to implantation of 1 stent, implantation of 2 or 3 stents resulted in a similar proportion of patients who were drop free at 36 months or less, but a higher proportion of patients who were drop free after 36 months.

The 2 studies included in the 2019 Cochrane review are described in Tables 15 and 16. Limitations of these studies are described in Tables 17 and 18.

**Table 14. Meta-analysis Results**

Study	Drop Free Compared to Medical Therapy	Drop Free with 2 Stents Compared to 1 Stent at 42 months	Drop Free with 3 Stents Compared to 1 Stent at 42 months
<b>Le et al. (2019) (32)</b>			
N	2 RCTs	1 RCT	1 RCT

Pooled effect (95% CI)	90% of patients in the iStent groups were drop free	RR: 0.51 (0.34 to 0.75)	RR: 0.49 (0.34 to 0.73)
------------------------	---	----------------------------	----------------------------

CI: confidence interval; RCT: randomized controlled trial; RR: relative risk.

#### Randomized Controlled Trials

A 2014 industry-sponsored, multicenter, unblinded, randomized trial compared implantation of 2 iStent inject devices to 2 ocular hypotensive agents. (49) The 192 patients enrolled in this unmasked trial had an IOP not controlled by 1 hypotensive medication. At 12-month follow-up, the 2 groups were comparable for IOP reduction of at least 20%, IOP of 18 mm Hg or less, and mean decrease in IOP. A greater proportion of patients in the iStent inject group achieved an IOP reduction of at least 50% (53.2% vs 35.7%, respectively). One patient in the iStent inject group experienced elevated IOP (48 mm Hg) and 4 required ocular hypotensive medication. Longer-term studies are in progress.

Vold et al. (2016) reported results of an RCT comparing 2 stand-alone iStent inject implants to topical travoprost (1:1 ratio) in 101 phakic eyes with an IOP between 21- and 40-mm Hg and newly diagnosed POAG, pseudoexfoliative glaucoma, or ocular hypertension that had not been treated previously. (50) The patients were not undergoing cataract surgery. The trial was unmasked, and methods for allocation concealment and calculation of power were not described. One hundred patients (54 iStent; 47 travoprost) completed 24 months of follow-up and 73 completed 36 months of follow-up. The trial was performed at a single center in Armenia with visiting surgeons from the U.S. Statistical analyses were not provided. Baseline mean IOP was 25 mm Hg in both groups. Mean IOP at 3 years was 15 mm Hg in both groups. Medication (or second medication) was added to 6 eyes in the iStent group and 11 eyes in the travoprost group. Progression of cataract was reported in 11 eyes in the iStent group and 8 eyes in the travoprost group, with cataract surgery being performed in 5 eyes in the iStent group and 1 eye in the travoprost group. The results would suggest that 2 iStents might reduce the number of medications required to maintain target IOP compared with travoprost but also hasten time to cataract surgery. However, the study methods were poorly reported, and statistical analyses were not reported.

Four-year follow-up of iStent inject is reported in 2 phase 4 publications from the MIGS study group. (51, 52) Berdahl et al. (2020) reported on 53 patients who were on 2 preoperative medications who received 2 iStent inject implants and started on travoprost on postoperative Day 1. At the 48-month follow-up, 85% of eyes had reduced IOP ( $\geq 20\%$ ) with a single medication as compared to the baseline IOP on 2 medications. Mean IOP on 1 medication was 11.9 to 13.0 mm Hg, compared to 19.7 on 2 medications preoperatively. Lindstrom et al. (2020) reported on 57 patients who were on 1 preoperative medication before implantation of 2 iStent inject devices. Month 48 IOP without medication was reduced ( $\geq 20\%$ ) in 95% of eyes with iStent inject. There were no adverse events that were considered to be related to the devices.

#### Hydrus versus iStent

Hydrus microstent was compared with the iStent in a double-blind multicenter RCT by Ahmed et al. (COMPARE, 2020). (45) Eyes (n=152) with mild-to-moderate glaucoma and an IOP of 23 to 39 after washout of medication were randomized to either 1 Hydrus stent or 2 iStents as a stand-alone treatment. Both stents have FDA approval in the U.S. when used in conjunction with cataract surgery but not as a stand-alone procedure. Follow-up was performed through 12 months post-operatively with medications added at the investigator's discretion. The Hydrus outperformed 2 iStents in nearly every measure (Table 16). Eyes implanted with the Hydrus microstent were able to maintain IOP < 18 mm Hg on fewer medications and a greater percentage of patients were medication-free compared to the iStent group (46.6% vs 24.0%, p<0.001). The decision to increase medications was up to the investigator and not pre-specified, but posthoc analysis indicated that the IOP at which medications were increased was similar in the 2 groups.

**Table 15. Summary of RCT Characteristics**

<b>Study; Trial</b>	<b>Countries</b>	<b>Sites</b>	<b>Dates</b>	<b>Participants</b>	<b>Interventions</b>	
					<b>Active</b>	<b>Comparator</b>
Fea et al. (2014) (49)	E.U., Armenia	8		Patients with OAG not controlled on one medication, Post-washout IOP >22 and <38 mmHg	iStent inject (n=94)	Two medications (n=98)
Vold et al. (2016) (50)	Armenia with U.S. surgeons	1		Patients with OAG or PEX who were naive to therapy with IOP > 21 and < 40 mmHg	Two iStents (n=54)	One medication (n=47)
Ahmed et al. (2019) (45)	U.S., E.U., Canada, Asia	12	2013- 2015	Patients with mild-to- moderate glaucoma (OAG, PEX, or PG and IOP 23 to 39 mmHg after washout	Hydrus (n=75)	Two iStents (n=77)

IOP: intraocular pressure; OAG: open-angle glaucoma; PEX: pseudoexfoliative glaucoma; PG: pigmentary glaucoma; RCT: randomized controlled trial.

**Table 16. Summary of RCT Results**

Study	>20% reduction in IOP, n (%)	IOP < 18 mm Hg, n (%)	Mean IOP mm Hg (SD)	Mean reduction in IOP from baseline mm Hg (SD)	Mean number of medications at 12 months	Percent Medication Free at 12 months, n (%)
Fea et al. (2014) (49)	<i>at 12 months</i>	<i>at 12 months</i>	<i>at 12 months</i>			
iStent inject	89/94 (94.7)	87/94 (92.6)	13.0 (2.3)	8.1 (2.6)		
Medical therapy	88/98 (91.8)	88/98 (89.8)	13.2 (2.0)	7.3 (2.2)		
p-Value	0.02	NR	NR	0.43		
Vold et al. (2016) (50)	<i>IOP &lt; 18 mm Hg at 24 months, n (%)</i>	<i>at 36 months</i>	<i>at 36 months</i>			
iStent	90%	91%	14.6 mmHg			
Medical therapy	87%	79%	15.3 mmHg			
p-Value						
Ahmed et al. (2020) (45)		<i>without medication</i>				
Hydrus	39.7%	30.1%	17.3 (3.7)	-8.2 (3.7)	1.0	34 (46.6)
2 iStents	13.3%	9.3%	19.2 (2.4)	-5.1 (2.9)	1.7	18 (24.0)
p-Value	<0.001	<0.001	0.037	0.003	<0.001	0.006

IOP: intraocular pressure; NR: not reported; RCT: randomized controlled trial; SD: standard deviation.

**Table 17. Study Relevance Limitations**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Follow-Up <sup>e</sup>
Fea et al. (2014) (49)					1. Follow-up was limited to 12 months. Monitoring for occlusion of the stents at longer follow-up is needed

Vold et al. (2016) (50)		4. Not the currently marketed device			
Ahmed et al. (2019) (45)		4. Not the currently marketed device		1. Follow-up was through 12 months, longer follow-up is continuing.	

The study limitations stated in this table are those notable in the current literature review; this is not a comprehensive gaps assessment

<sup>a</sup>Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

<sup>b</sup>Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

<sup>c</sup>Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

<sup>d</sup>Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

<sup>e</sup>Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

**Table 18. Study Design and Conduct Limitations**

Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness <sup>d</sup>	Power <sup>e</sup>	Statistical <sup>f</sup>
Fea et al. (2014) (49)	3. Randomization procedure was not described	1, 2, 3. Study could not be blinded		1. Unequal loss to follow-up in the 2 groups, and the subjects lost to follow-up were treated as failures	1. Power calculation s not reported	
Vold et al. (2016) (50)	3. Randomization procedure was not described	1, 2, 3. Study could not be blinded		1. There was 27% loss to follow-up at 36 months	1. Power calculation s not reported	4. Statistical analysis not reported
Ahmed et al. (2020) (45)		2, 3. Investigators were not blinded and there				2. Did not use repeated measures for multiple

		was not independent adjudication or preset criteria for increase in medication				assessments
--	--	--	--	--	--	-------------

The study limitations stated in this table are those notable in the current literature review; this is not a comprehensive gaps assessment.

<sup>a</sup> Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup> Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

<sup>c</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup> Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

<sup>e</sup> Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

<sup>f</sup> Statistical key: 1. Analysis is not appropriate for outcome type: a) continuous; b) binary; c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

### *Greater Than Two Stents*

An RCT comparing the efficacy of 1 iStent with multiple iStent devices was published by Katz et al. (2015). (53) This trial, from a single-institution in Armenia, randomized 119 patients with mild-to-moderate OAG and an IOP between 22- and 38-mm Hg (off medications) to 1 stent (n=38), 2 stents (n=41), or 3 stents (n=40). Randomization was performed using a pseudorandom number generator. The primary endpoint was the percentage of patients with a reduction of 20% or more in IOP off medications at 12 months. This endpoint was reached by 89.2% of the 1-stent group, by 90.2% of the 2-stent group, and by 92.1% of the 3-stent group. The secondary endpoint (percentage of patients achieving an IOP  $\leq$ 15 mm Hg off medication) was reached by 64.9% of the 1-stent group, by 85.4% of the 2-stent group, and by 92.1% of the 3-stent group. Forty-two-month follow-up results for 109 patients were published by Katz et al. (2018). (54) Post-washout IOP was  $17.4\pm0.9$ ,  $15.8\pm1.1$  and  $14.2\pm1.5$  mmHg, for 1, 2, or 3 stents, respectively. The need for additional medication increased in single-stent eyes from 4 eyes at 12 months to 18 eyes at 42 months, suggesting a reduction in patency of the microstents over time. The need for additional medication did not increase from months 12 and 42 in multi-stent eyes. No between-group statistical comparisons were reported.

## Nonrandomized Studies

Sarkisian et al. (2023) published the results of an open-label, single-arm, pivotal study evaluating iStent infinite in patients with OAG uncontrolled by prior surgical or medical therapy. (55) The trial enrolled a total of 72 patients from 15 sites. The majority of patients had failed prior surgery (n=61) and the remainder were uncontrolled on medical therapy (n=11). At 12 months the proportion of patients achieving at least 20% reduction in IOP and receiving the same or fewer medications was 76.1% (95% CI, 66.2% to 86.1%). The mean reduction in IOP at 12 months was 5.9 mm Hg (standard error, 0.6; 95% CI, 4.8 to 7.1). No serious device-related adverse events were reported; however, blepharitis (4.2%), IOP increase requiring surgical intervention (4.2%), loss of best spectacle corrected visual acuity of 2 lines or more (8.3%), ocular surface disease (9.7%), and visual field loss of at least 2.5 dB were commonly reported adverse events. Stent migration and stent obstruction were each reported in 2 patients. Although this trial indicates positive outcomes with iStent infinite, the small sample size and lack of a control group are significant limitations.

## Section Summary: Microstent Implantation as a Stand-Alone Procedure

The evidence on microstents as a stand-alone procedure in patients with mild-to-moderate glaucoma that is controlled on medical therapy includes a nonrandomized study, RCTs and a systematic review of 3 heterogeneous RCTs. Two RCTs indicate that implantation of a microstent can reduce IOP at a level similar to ocular medications at 12-month follow-up. Reduction in medications is an important outcome for patients with glaucoma, both for the patients themselves and because lack of compliance can lead to adverse health outcomes. Whether microstents remain patent after 12 months is uncertain, and whether additional stents can subsequently be safely implanted is unknown. Some evidence on longer-term outcomes is provided by an RCT that compared implantation of a single iStent with multiple iStents. At longer-term (42-month) follow-up, the need for additional medication increased in eyes implanted with a single iStent but not with multiple iStents. The durability of multiple iStents is unknown. A fourth RCT compared implantation of the Hydrus microstent to 2 iStents. Outcomes from the Hydrus microstent were significantly better than 2 iStents, both statistically and clinically, for all outcome measures. The primary limitation of this study is that the duration of follow-up in the present publication is limited to 12 months. Longer-term follow-up from this study is continuing and will answer important questions on the durability of the procedure. Corroboration in an independent study and comparison with a medical therapy control group would also increase confidence in the results.

## **Summary of Evidence**

For individuals who have refractory open-angle glaucoma (OAG) who receive ab externo aqueous shunts, the evidence includes randomized controlled trials (RCTs), retrospective studies, and systematic reviews. Relevant outcomes are a change in disease status, functional outcomes, medication use, and treatment-related morbidity. RCTs assessing Food and Drug Administration (FDA) approved shunts have shown that the use of large externally placed shunts reduces intraocular pressure (IOP) to slightly less than standard filtering surgery (trabeculectomy). Reported shunt success rates show that these devices are noninferior to trabeculectomy in the long-term. The FDA approved shunts have different adverse event

profiles and avoid some of the most problematic complications of trabeculectomy. The FDA-approved shunts have different adverse event profiles and avoid some of the most problematic complications of trabeculectomy. Two trials have compared the Ahmed and Baerveldt shunts. Both found that eyes treated with the Baerveldt shunt had slightly lower average IOP at 5 years than eyes treated with the Ahmed, but the Baerveldt also had a higher rate of serious hypotony-related complications. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have refractory OAG who receive ab interno aqueous stents, the evidence includes systematic reviews, an RCT, nonrandomized comparative studies, and a single-arm study. Relevant outcomes are a change in disease status, functional outcomes, medication use, and treatment-related morbidity. The RCT found XEN45 to be noninferior to trabeculectomy. The nonrandomized comparative studies reported that patients receiving the stent experienced similar reductions in IOP and medication use as patients undergoing trabeculectomy. The single-arm studies, with 12-month follow-up results, consistently showed that patients receiving the stents experienced reductions in IOP and medication use. In addition, the FDA has given clearance to a gel stent based on equivalent IOP and medication use reductions as seen with ab externo shunts. Clearance for the stent was based on a review in which the FDA concluded that while there were technical differences between the stent and predicate devices (shunts), the differences did not affect safety and effectiveness in lowering IOP and medication use. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have mild-to-moderate OAG who are undergoing cataract surgery who receive aqueous microstents, the evidence includes RCTs and meta-analyses of RCTs. Relevant outcomes are a change in disease status, functional outcomes, medication use, and treatment-related morbidity. Implantation of 1 or 2 microstents has received FDA approval for use in conjunction with cataract surgery for reduction of IOP in adults with mild-to-moderate OAG currently treated with ocular hypotensive medication. When compared to cataract surgery alone, the studies showed modest but statistically significant decreases in IOP, and medication use through the first 2 years when stents were implanted in conjunction with cataract surgery. A decrease in topical medication application is considered to be an important outcome for patients and reduces the problem of non-compliance that can affect visual outcomes. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with mild-to-moderate OAG who are not undergoing cataract surgery who receive aqueous microstents as a stand-alone procedure, the evidence includes a nonrandomized trial, RCTs, and a systematic review of 3 heterogeneous RCTs. Relevant outcomes are a change in disease status, functional outcomes, medication use, and treatment-related morbidity. Several RCTs have evaluated the use of multiple microstents but comparators differed. Two RCTs indicate that implantation of a microstent can reduce IOP at a level similar to ocular medications at 12-month follow-up. Reduction in medications is an important outcome for patients with glaucoma. Whether microstents remain patent after 12

months is uncertain, and whether additional stents can subsequently be safely implanted is unknown. Some evidence on longer-term outcomes is provided by an RCT that compared implantation of a single iStent to implantation of multiple iStents. At longer-term (42-month) follow-up, the need for additional medication increased in eyes implanted with a single microstent but not with multiple microstents. The durability of multiple iStents is unknown. A fourth RCT compared implantation of the Hydrus microstent to 2 iStents. Outcomes from the Hydrus microstent were significantly better than 2 iStents, both statistically and clinically, for all outcome measures. The primary limitation of this study is that the duration of follow-up in the present publication is limited to 12 months. Longer-term follow-up from this study is continuing and will answer important questions on the durability of the procedure. Corroboration in an independent study and comparison with a medical therapy control group would also increase confidence in the results. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

#### **Clinical Input from Physician Specialty Societies and Academic Medical Centers**

Clinical input supports the use of aqueous shunts in patients with glaucoma uncontrolled by medication, as well as the use of a single microstent in patients with mild-to-moderate glaucoma undergoing cataract surgery to reduce the adverse events of medications and to avoid noncompliance.

#### **Practice Guidelines and Position Statements**

##### American Academy of Ophthalmology (AAO)

The AAO (2008) published a technology assessment on commercially available aqueous shunts, including the Ahmed, Baerveldt, Krupin, and Molteno devices, which was last reviewed for currency in 2014

. (5) The assessment indicated that, in general, IOP would settle at higher levels ( $\geq 18$  mm Hg) with shunts than after standard trabeculectomy (14-16 mm Hg). Five-year success rates of 50% were found for the 2 procedures, indicating that aqueous shunts are comparable with trabeculectomy for IOP control and duration of benefit (based on level I evidence; well-designed randomized controlled trials). The assessment also indicated that although aqueous shunts have generally been reserved for intractable glaucoma when prior medical or surgical therapy has failed, indications for shunts have broadened (based on level III evidence; case series, case reports, and poor-quality case-control or cohort studies). The AAO concluded that, based on level I evidence, aqueous shunts offer a valuable alternative to standard filtering surgery and cyclodestructive therapy for many patients with refractory glaucoma.

The AAO's (2016) preferred practice patterns on POAG indicated that the Academy considered laser trabeculoplasty as initial therapy in select patients or an alternative for patients who cannot or will not use medications reliably due to cost, memory problems, difficulty with installation, or intolerance to the medication. (56) The AAO stated that aqueous shunts have traditionally been used to manage refractory glaucoma when trabeculectomy has failed to control IOP or is unlikely to succeed, but these devices are being increasingly used in other

indications for the surgical management of glaucoma. The AAO also stated that micro-invasive glaucoma surgeries that are frequently combined with phacoemulsification have limited long-term data but seem to result in modest IOP reduction with postoperative pressures in the mid to upper teens. Although they are less effective in lowering IOP than trabeculectomy and aqueous shunt surgery, micro-invasive glaucoma surgeries may have a more favorable safety profile in the short term.

In 2020, the AAO updated its preferred practice pattern on POAG. (56) The document notes that aqueous shunts have traditionally been used to manage medically uncontrolled glaucoma when trabeculectomy has failed to control IOP or is deemed unlikely to succeed; however, the indications for using aqueous shunts have been broadening, and these devices are being increasingly used in the surgical management of glaucoma. The preferred practice pattern notes that "several studies have compared aqueous shunts with trabeculectomy" and that the "selection of aqueous shunts or trabeculectomy should be left to the discretion of the treating ophthalmologist, in consultation with the individual patient."

#### American Glaucoma Society

In 2020, the American Glaucoma Society published a position paper on microinvasive glaucoma surgery. (57) The Society supports efforts that facilitate patient access to these procedures, including more flexible regulatory pathways for new devices, expansion of the indications for already approved devices, and greater availability of information obtained by regulatory authorities.

#### National Institute for Health and Care Excellence

The National Institute for Health and Care Excellence (2017) updated guidance on trabecular stent bypass microsurgery for OAG. (58) The guidance stated that "Current evidence on trabecular stent bypass microsurgery for open-angle glaucoma raises no major safety concerns. Evidence of efficacy is adequate in quality and quantity."

The National Institute for Health and Care Excellence (2018) published guidance entitled "Microinvasive subconjunctival insertion of a trans-scleral gelatin stent for POAG." (59) The guidance states that evidence is limited in quantity and quality and therefore, the procedure should only be used with special arrangements and that patients should be informed of the uncertainty of the procedure.

#### **Ongoing and Unpublished Clinical Trials**

Some currently ongoing and unpublished trials that might influence this policy are listed in Table 19.

**Table 19. Summary of Key Trials**

NCT No	Trial Name	Planned Enrollment	Completion Date
<b><i>Ongoing</i></b>			

NCT05439161	Multicentric Evaluation of Best Corrected Visual Acuity of the XEN Implant Versus Classic Trabeculectomy in Open Angle Glaucoma Subjects	196	Apr 2025
NCT05411198 <sup>a</sup>	A Prospective, Multicenter Clinical Study to Evaluate the Safety and Effectiveness of Ab Extero Implantation of Glaucoma Gel Stent	65	Aug 2025
NCT04440527	Intraocular Pressure After Preserflo/Innfocus Microshunt vs Trabeculectomy: a Prospective, Randomised Control-trial (PAINT-Study)	70	Jul 2024
NCT04624698 <sup>a</sup>	iStent Inject Trabecular Micro-Bypass System New Enrollment Post-Approval Study	358	Jun 2026
NCT06066645 <sup>a</sup>	Multicenter, Randomized, Double-masked Trial to Evaluate the Safety and Efficacy of iDose® TR (Travoprost Intraocular Implant) in Conjunction With the Placement of iStent Infinite vs. iStent Infinite Alone in Subjects With Open-angle Glaucoma or Ocular Hypertension	150	Nov 2025
NCT06057051 <sup>a</sup>	A Prospective, Multicenter Study of the Glaukos® iStent Infinite Trabecular Micro-Bypass System Model iS3 in Subjects With Mild to Moderate Primary Open-angle Glaucoma	245	Aug 2027
NCT04635020 <sup>a</sup>	A Prospective Randomised Trial Comparing Selective Laser Trabeculoplasty (SLT) and iStent Trabecular Micro-bypass Stent Implantation Combined With Cataract Surgery in Exfoliation Glaucoma	285	Sep 2033
NCT05583591 <sup>a</sup>	Prospective, Randomized Controlled Study Comparing Combined Phacoemulsification With iStent Inject W Versus Hydrus for Mild to Moderate Open Angle Glaucoma (COMPETE)	390	Oct 2025
NCT05280366 <sup>a</sup>	A Prospective, Randomized, Multi-center Evaluation of the Safety and Effectiveness of the STREAMLINE®SURGICAL SYSTEM Compared to iStent Inject W® in Patients With Open-Angle Glaucoma	150	Jun 2026
NCT06289491 <sup>a</sup>	Randomized Trial of Hydrus Microstent Versus Goniotomy	243	Apr 2029

NCT04553523 <sup>a</sup>	The Hydrus® Microstent New Enrollment Post-Approval Study: A Prospective, Non-Randomized, Multicenter, Single Arm, Clinical Trial	545	Jun 2028
NCT05949242 <sup>a</sup>	Comparison of Clinical Outcomes in Patients Undergoing Cataract Surgery With OMNI Canaloplasty vs Cataract Surgery With OMNI Canaloplasty and Hydrus Stent	80	Oct 2024
NCT03904381 <sup>a</sup>	Efficacy and Safety of XEN® Gel Stent and Post-operative Management in Patients With Open Angle Glaucoma Compared to Classic Glaucoma Surgeries (Trabeculectomy and Sclerectomy) as Well as Other Minimally Invasive Glaucoma Surgery (MIGS)	100	Jan 2025
NCT05340647 <sup>a</sup>	NorMIGS - a Prospective Study of Micro-invasive Glaucoma Surgery	100	Jun 2028
<b><i>Unpublished</i></b>			
NCT02327312 <sup>a</sup>	Multicenter Investigation of Trabecular Micro-Bypass Stents vs. Laser Trabeculoplasty	91	Aug 2020
NCT04629521 <sup>a</sup>	An Observational Multicenter Clinical Study to Provide Additional Long-Term Follow-up Beyond 60 Months for Subjects Implanted With a CyPass Micro-Stent in the COMPASS Trial	54	Aug 2023
NCT04658095 <sup>a</sup>	A Prospective, Randomized, Multicenter Study To Compare The Safety And Effectiveness Of The OMNI® Surgical System And The iStent Inject In Pseudophakic Eyes With Open Angle Glaucoma. The TRIDENT European Trial	20	Aug 2022
NCT01841450 <sup>a</sup>	A Prospective, Randomized, Controlled, Parallel Groups, Multicenter 60 Post-Approval Study Of The Glaukos® iStent® Trabecular Micro-Bypass Stent System In Conjunction With Cataract Surgery.	360	Nov 2021
NCT01444040 <sup>a</sup>	A Prospective, Randomized Evaluation of Subjects With Open-angle Glaucoma, Pseudoexfoliative Glaucoma, or Ocular Hypertension Naïve to Medical and Surgical Therapy, Treated With Two Trabecular	196	Mar 2019

	Micro-bypass Stents (iStent <i>Inject</i> ) or Travoprost Ophthalmic Solution 0.004%		
NCT01461278 <sup>a</sup>	A Prospective, Randomized, Single-Masked, Controlled, Parallel Groups, Multicenter Clinical Investigation of the Glaukos® Suprachoroidal Stent Model G3 In Conjunction With Cataract Surgery	505	Mar 2020

NCT: National clinical trial

<sup>a</sup> Denotes industry-sponsored or co-sponsored trial.

## Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member's benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

<b>CPT Codes</b>	66179, 66180, 66183, 66184, 66185, 66989, 66991, 66999, 0253T, 0449T, 0450T, 0474T, 0671T
<b>HCPCS Codes</b>	C1783, L8612

\*Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.

## References

- Allison K, Patel DG, Greene L. Racial and Ethnic Disparities in Primary Open-Angle Glaucoma Clinical Trials: A Systematic Review and Meta-analysis. *JAMA Netw Open*. May 03 2021; 4(5):e218348. PMID 34003274
- Panarelli JF, Moster MR, Garcia-Feijoo J, et al. Ab-Extero MicroShunt versus Trabeculectomy in Primary Open-Angle Glaucoma: Two-Year Results from a Randomized, Multicenter Study. *Ophthalmology*. Mar 2024; 131(3):266-276. PMID 37769852
- Minckler DS, Vedula SS, Li TJ, et al. Aqueous shunts for glaucoma. *Cochrane Database Syst Rev*. Apr 19 2006; (2):CD004918. PMID 16625616
- Tseng VL, Coleman AL, Chang MY, et al. Aqueous shunts for glaucoma. *Cochrane Database Syst Rev*. Jul 28 2017; 7(7):CD004918. PMID 28750481
- Minckler DS, Francis BA, Hodapp EA, et al. Aqueous shunts in glaucoma: a report by the American Academy of Ophthalmology. *Ophthalmology*. Jun 2008; 115(6):1089-1098. PMID 18519069
- Zhang X, Wang B, Liu R, et al. The effectiveness of AGV, Ex-PRESS, or trabeculectomy in the treatment of primary and secondary glaucoma: a systematic review and network meta-analysis. *Ann Palliat Med*. Jan 2022; 11(1):321-331. PMID 35144423

7. Boland MV, Ervin AM, Friedman D, et al. Treatment for Glaucoma: Comparative Effectiveness. Comparative Effectiveness Review No. 60 (AHRQ Publication No. 12-EHC038-EF). Rockville, MD: Agency for Healthcare Research and Quality; 2012.
8. Gedde SJ, Schiffman JC, Feuer WJ, et al. Treatment outcomes in the Tube Versus Trabeculectomy (TVT) study after five years of follow-up. *Am J Ophthalmol.* May 2012; 153(5):789-803.e2. PMID 22245458
9. Kotecha A, Feuer WJ, Barton K, et al. Quality of life in the tube versus trabeculectomy study. *Am J Ophthalmol.* Apr 2017; 176:228-235. PMID 28161049
10. Swaminathan SS, Jammal AA, Kornmann HL, et al. Visual field outcomes in the tube versus trabeculectomy study. *Ophthalmology.* Sep 2020; 127(9):1162-1169. PMID 32327255
11. Swaminathan SS, Jammal AA, Medeiros FA, et al. Visual Field Outcomes in the Primary Tube Versus Trabeculectomy Study. *Ophthalmology.* Oct 2024; 131(10):1157-1163. PMID 38582154
12. Wang X, Khan R, Coleman A. Device-modified trabeculectomy for glaucoma. *Cochrane Database Syst Rev.* Dec 01 2015; (12):CD010472. PMID 26625212
13. Park J, Rittiphairoj T, Wang X, et al. Device-modified trabeculectomy for glaucoma. *Cochrane Database Syst Rev.* Mar 13 2023; 3(3):CD010472. PMID 36912740
14. Netland PA, Sarkisian SR, Moster MR, et al. Randomized, prospective, comparative trial of EX-PRESS glaucoma filtration device versus trabeculectomy (XVT study). *Am J Ophthalmol.* Feb 2014; 157(2):433-440.e3. PMID 24210765
15. de Jong LA. The Ex-PRESS glaucoma shunt versus trabeculectomy in open-angle glaucoma: a prospective randomized study. *Adv Ther.* Mar 2009; 26(3):336-345. PMID 19337705
16. de Jong L, Lafuma A, Aguade AS, et al. Five-year extension of a clinical trial comparing the EX-PRESS glaucoma filtration device and trabeculectomy in primary open-angle glaucoma. 2011; 5:527-533. PMID 21607021
17. Wagschal LD, Trope GE, Jinapriya D, et al. Prospective randomized study comparing Ex-PRESS to trabeculectomy: 1-year results. *J Glaucoma.* Oct-Nov 2015; 24(8):624-629. PMID 24247999
18. Gonzalez-Rodriguez JM, Trope GE, Drori-Wagschal L, et al. Comparison of trabeculectomy versus Ex-PRESS: 3-year follow-up. *Br J Ophthalmol.* Sep 2016; 100(9):1269-1273. PMID 26674779
19. Konopinska J, Byszewska A, Saeed E, et al. Phacotrabeculectomy versus phaco with implantation of the Ex-PRESS device: surgical and refractive outcomes-a randomized controlled trial. *J Clin Med.* Jan 22 2021; 10(3):424. PMID 33499300
20. Tokumo K, Okada N, Onoe H, et al. Ex-PRESS Implantation versus Trabeculectomy for Long-Term Maintenance in Patients with Open-Angle Glaucoma. *Clin Ophthalmol.* 2023; 17:2525-2537. PMID 37662650
21. Budenz DL, Barton K, Gedde SJ, et al. Five-year treatment outcomes in the Ahmed Baerveldt comparison study. *Ophthalmology.* Feb 2015; 122(2):308-316. PMID 25439606
22. Budenz DL, Feuer WJ, Barton K, et al. Postoperative complications in the Ahmed Baerveldt comparison study during five years of follow-up. *Am J Ophthalmol.* Mar 2016; 163:75-82.e3. PMID 26596400
23. Christakis PG, Kalenak JW, Tsai JC, et al. The Ahmed versus Baerveldt Study: five-year treatment outcomes. *Ophthalmology.* Oct 2016; 123(10):2093-2102. PMID 27544023

24. Christakis PG, Zhang D, Budenz DL, et al. Five-year pooled data analysis of the Ahmed Baerveldt comparison study and the Ahmed versus Baerveldt study. *Am J Ophthalmol.* Apr 2017; 176:118-126. PMID 28104418

25. Lim SY, Betzler BK, Yip LWL, et al. Standalone XEN45 Gel Stent implantation in the treatment of open-angle glaucoma: A systematic review and meta-analysis. *Surv Ophthalmol.* 2022; 67(4):1048-1061. PMID 35081414

26. Yang X, Zhao Y, Zhong Y, et al. The efficacy of XEN gel stent implantation in glaucoma: a systematic review and meta-analysis. *BMC Ophthalmol.* Jul 15 2022; 22(1):305. PMID 35836197

27. Sheybani A, Vera V, Grover DS, et al. Gel Stent Versus Trabeculectomy: The Randomized, Multicenter, Gold-Standard Pathway Study (GPS) of Effectiveness and Safety at 12 Months. *Am J Ophthalmol.* Aug 2023; 252:306-325. PMID 36972738

28. Schlenker MB, Gulamhusein H, Conrad-Hengerer I, et al. Efficacy, safety, and risk factors for failure of standalone ab interno gelatin microstent implantation versus standalone trabeculectomy. *Ophthalmology.* Nov 2017; 124(11):1579-1588. PMID 28601250

29. Wagner FM, Schuster AK, Emmerich J, et al. Efficacy and safety of XEN(R)-Implantation vs. trabeculectomy: Data of a "real-world" setting. *PLoS One.* 2020; 15(4):e0231614. PMID 32310972

30. Stoner AM, Capitena Young CE, SooHoo JR, et al. A comparison of clinical outcomes after XEN Gel Stent and EX-PRESS glaucoma drainage device implantation. *J Glaucoma.* Jun 01 2021; 30(6):481-488. PMID 34060508

31. Gabbay IE, Goldberg M, Allen F, et al. Efficacy and safety data for the Ab interno XEN45 gel stent implant at 3 Years: A retrospective analysis. *Eur J Ophthalmol.* May 02 2021: 11206721211014381. PMID 33938304

32. Le JT, Bicket AK, Wang L, et al. Ab interno trabecular bypass surgery with iStent for open-angle glaucoma. *Cochrane Database Syst Rev.* Mar 28 2019; 3:CD012743. PMID 30919929

33. Healey PR, Clement CI, Kerr NM, et al. Standalone iStent trabecular micro-bypass glaucoma surgery: a systematic review and meta-analysis. *J Glaucoma.* Jul 01 2021; 30(7):606-620. PMID 33596009

34. Samuelson TW, Katz LJ, Wells JM, et al. Randomized evaluation of the trabecular micro-bypass stent with phacoemulsification in patients with glaucoma and cataract. *Ophthalmology.* Mar 2011; 118(3):459-467. PMID 20828829

35. Craven ER, Katz LJ, Wells JM, et al. Cataract surgery with trabecular micro-bypass stent implantation in patients with mild-to-moderate open-angle glaucoma and cataract: two-year follow-up. *J Cataract Refract Surg.* Aug 2012; 38(8):1339-1345. PMID 22814041

36. Samuelson TW, Sarkisian SR, Lubeck DM, et al. Prospective, randomized, controlled pivotal trial of an ab interno implanted trabecular micro-bypass in primary open-angle glaucoma and cataract: two-year results. *Ophthalmology.* Jun 2019; 126(6):811-821. PMID 30880108

37. Hooshmand J, Rothschild P, Allen P, et al. Minimally invasive glaucoma surgery: Comparison of iStent with iStent inject in primary open angle glaucoma. *Clin Experiment Ophthalmol.* Sep 2019; 47(7):898-903. PMID 31034687

38. Al Yousef Y, Strzalkowska A, Hillenkamp J, et al. Comparison of a second-generation trabecular bypass (iStent inject) to ab interno trabeculectomy (Trabectome) by exact

matching. *Graefes Arch Clin Exp Ophthalmol*. Dec 2020; 258(12):2775-2780. PMID 32960322

39. Salimi A, Watt H, Harasymowycz P. Three-year outcomes of second-generation trabecular micro-bypass stents (iStent inject) with phacoemulsification in various glaucoma subtypes and severities. *J Glaucoma*. Mar 01 2021; 30(3):266-275. PMID 33105306

40. Matsuo M, Fukuda H, Buathong J, et al. Comparison of 1-year effectiveness between phaco-microhook ab-interno trabeculotomy and phaco-iStent trabecular micro-bypass stent in primary open-angle glaucoma with low-teen intraocular pressure. *Graefes Arch Clin Exp Ophthalmol*. Aug 19 2024. PMID 39160440

41. Fan Gaskin JC, Bigirimana D, Kong GYX, et al. Prospective, Randomized Controlled Trial of Cataract Surgery vs Combined Cataract Surgery With Insertion of iStent Inject. *Ophthalmol Glaucoma*. 2024; 7(4):326-334. PMID 38369058

42. Otarola F, Virgili G, Shah A, et al. Ab interno trabecular bypass surgery with Schlemms canal microstent (Hydrus) for open angle glaucoma. *Cochrane Database Syst Rev*. Mar 09 2020; 3:CD012740. PMID 32147807

43. Pfeiffer N, Garcia-Feijoo J, Martinez-de-la-Casa JM, et al. A randomized trial of a Schlemm's canal microstent with phacoemulsification for reducing intraocular pressure in open-angle glaucoma. *Ophthalmology*. Jul 2015; 122(7):1283-1293. PMID 25972254

44. Samuelson TW, Chang DF, Marquis R, et al. A Schlemm canal microstent for intraocular pressure reduction in primary open-angle glaucoma and cataract: the HORIZON study. *Ophthalmology*. Jan 2019; 126(1):29-37. PMID 29945799

45. Ahmed IIK, Fea A, Au L, et al. A prospective randomized trial comparing Hydrus and iStent microinvasive glaucoma surgery implants for standalone treatment of open-angle glaucoma: the COMPARE study. *Ophthalmology*. Jan 2020; 127(1):52-61. PMID 31034856

46. Montesano G, Ometto G, Ahmed IIK, et al. Five-Year Visual Field Outcomes of the HORIZON Trial. *Am J Ophthalmol*. Jul 2023; 251:143-155. PMID 36813144

47. Fea AM, Ahmed II, Lavia C, et al. Hydrus microstent compared to selective laser trabeculoplasty in primary open angle glaucoma: one year results. *Clin Experiment Ophthalmol*. Mar 2017; 45(2):120-127. PMID 27449488

48. Salimi A, Kassem R, Santhakumaran S, et al. Three-Year Outcomes of a Schlemm Canal Microstent (Hydrus Microstent) with Concomitant Phacoemulsification in Open-Angle Glaucoma. *Ophthalmol Glaucoma*. 2023; 6(2):137-146. PMID 36038108

49. Fea AM, Belda JI, Rekas M, et al. Prospective unmasked randomized evaluation of the iStent inject versus two ocular hypotensive agents in patients with primary open-angle glaucoma. 2014; 8:875-882. PMID 24855336

50. Vold SD, Voskanyan L, Tetz M, et al. Newly diagnosed primary open-angle glaucoma randomized to 2 trabecular bypass stents or prostaglandin: outcomes through 36 months. Dec 2016; 5(2):161-172. PMID 27619225

51. Berdahl J, Voskanyan L, Myers JS, et al. iStent inject trabecular micro-bypass stents with topical prostaglandin as standalone treatment for open-angle glaucoma: 4-year outcomes. *Clin Experiment Ophthalmol*. Aug 2020; 48(6):767-774. PMID 32311201

52. Lindstrom R, Sarkisian SR, Lewis R, et al. Four-year outcomes of two second-generation trabecular micro-bypass stents in patients with open-angle glaucoma on one medication. 2020; 14:71-80. PMID 32021070

53. Katz LJ, Erb C, Carceller GA, et al. Prospective, randomized study of one, two, or three trabecular bypass stents in open-angle glaucoma subjects on topical hypotensive medication. 2015; 9:2313-2320. PMID 26715834

54. Katz LJ, Erb C, Carceller Guillamet A, et al. Long-term titrated IOP control with one, two, or three trabecular micro-bypass stents in open-angle glaucoma subjects on topical hypotensive medication: 42-month outcomes. 2018; 12:255-262. PMID 29440867

55. Sarkisian SR, Grover DS, Gallardo MJ, et al. Effectiveness and Safety of iStent Infinite Trabecular Micro-Bypass for Uncontrolled Glaucoma. *J Glaucoma*. Jan 01 2023; 32(1):9-18. PMID 36260288

56. Gedde SJ, Vinod K, Wright MM, et al. Primary open-angle glaucoma preferred practice pattern. September 2020. Available at <<https://www.aoa.org>> (accessed December 21, 2024).

57. Fellman RL, Mattox C, Singh K, et al. American Glaucoma Society Position Paper: Microinvasive Glaucoma Surgery. *Ophthalmol Glaucoma*. Jan 2020; 3(1):1-6. PMID 32672638

58. National Institute for Health and Care Evidence (NICE). Trabecular stent bypass microsurgery for open-angle glaucoma [IPG575]. 2017; Available at: <<https://www.nice.org.uk>> (accessed September 25, 2024).

59. National Institute for Health and Care Excellence. Microinvasive subconjunctival insertion of a trans-scleral gelatin stent for primary open-angle glaucoma. [IPG612]. 2018; Available at: <<https://www.nice.org.uk>> (accessed September 25, 2024).

## **Centers for Medicare and Medicaid Services (CMS)**

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does not have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been developed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

## **Policy History/Revision**

<b>Date</b>	<b>Description of Change</b>
02/01/2025	Document updated with literature review. Coverage unchanged. Added references 1, 2, 4, 6, 11, 13, 20, 25-27, 40, 41, 46, 48, 55, 56; others updated.
01/01/2024	Reviewed. No changes.
05/15/2022	Document updated with literature review. Coverage unchanged. Added references 1, 7, 14, 20-22, 24, 29, 30 and 43.
02/01/2022	Reviewed. No changes.

06/15/2021	Document updated with literature review. The following change was made in Coverage: Added term "ab" to the existing policy statement "Implantation of 1 or 2 U.S. Food and Drug Administration approved ab interno stents in conjunction with cataract surgery may be considered medically necessary in patients with mild-to-moderate open-angle glaucoma treated with ocular hypotensive medication." Added references 19, 22-24, 26-28, 31 and 37; others updated and/or removed.
08/15/2020	Reviewed. No changes.
09/15/2019	Document updated with literature review. The following changes were made to Coverage 1) Separated ab interno and ab externo coverage statements. 2) Changed the use of ab interno aqueous stents to medically necessary when criteria are met; 3) Expanded coverage to allow for 1 "or 2" FDA approved interno stents when used in conjunction with cataract surgery in pts with mild to moderate open angle glaucoma treated with hypotensive medications; 4) Added a separate "ab interno" experimental, investigational, and/or unproven statement for all other conditions. Added references 1, 2, 7, 14, 15, 19-29, 34, 35, 39, 43, 44, 49, 52-56.
03/01/2018	Document updated with literature review. The following changes were made to Coverage 1) Removed Ab interno suprachoroidal microstent (i.e., CyPass Micro-Stent) from the experimental, investigational and/or unproven coverage statement. 2) Added language for XEN to state "Ab interno insertion of an aqueous drainage device into the subconjunctival space (i.e., XEN Glaucoma Treatment System) is considered experimental, investigational and/or unproven for all indications" 3) Added "Note 1: Refer to the Description Section for mild to moderate open-angle glaucoma definition".
07/01/2017	Document updated with literature review. The following was added to Coverage: Ab interno suprachoroidal microstent (i.e., CyPass Micro-Stent) are considered experimental, investigational and/or unproven for all indications.
07/15/2016	Reviewed. No changes.
03/15/2015	Document updated with literature review. No change to coverage.
07/01/2014	Document updated with literature review. The following was added to coverage: 1) Implantation of a single FDA-approved microstent in conjunction with cataract surgery may be considered medically necessary in patients with mild to moderate open-angle glaucoma currently treated with ocular hypotensive medication. 2) Use of a microstent for all other conditions is considered experimental, investigational, and/or unproven. Document title changed from Aqueous Shunts for Glaucoma. CPT/HCPCS code(s) updated.
08/15/2012	New medical document originating from SUR713.030, Surgical Treatments for Glaucoma that has been deleted.

